

Impact of Wheat-Barley Blending on Rheological, Textural and Sensory Attributes of Leavened Bread

Fiza Nazir¹ and Nayik GA^{2*}

¹Division of Post-Harvest Technology, SKUAST-Kashmir, India

²Department of Food Engineering and Technology, SLIET, Punjab, India

Abstract

The aim of the study was to develop a healthy alternative to wheat-leavened bread by using wheat-barley blended flour. The leavened breads were prepared by blending wheat and barley flour at different levels and varying the MSG concentration. A significant increase was observed in the extensibility of the dough with increase with increasing MSG concentration before fermentation while after fermentation it showed a significant decreasing effect. The highest value of color, taste and appearance was observed for T₂M₁. Results for the flavor of the breads revealed that the highest flavor score was observed for T₂M₂. The results showed that the overall acceptability score decreased with increasing barley flour and MSG level.

Keywords: Wheat flour; Barley flour; Leavened bread; MSG; Sensory analysis

Introduction

Barley (*Hordeum vulgare L.*) is an ancient and important cereal grain crop ranking fifth among all crops in dry matter production in the world. It is one of the most widely cultivated cereal crops that can provide valuable nutrients required by humans and domestic animals. The high adaptability of barley to various climates and growing conditions has led to its increased worldwide production. Barley is arguably the most widely adapted cereal grain species with production at higher latitudes and altitudes and farther in deserts than any other cereal crop. It is in extreme climates that barley remains a principal food source today, e.g. Himalayan nations, Ethiopia, and Morocco [1]. Whole barley grain consists of about 65% to 68% starch, 10% to 17% protein, 4% to 9% β-glucan, 2% to 3% free lipids and 1.5% to 2.5% minerals [2-4]. Human consumption of barley and barley-containing food products has been insignificant as compared to other cereal grains, the development of new processes and food products has been neglected and there has been little effort to define quality requirements for food uses. The significance of β-glucan and tocopherols for human nutrition is well known, but little is known about the functional properties of β-glucan for making food products. Some of the traits preferred for specific food applications are known through investigations on incorporating barley into wheat-based food products. Barley flour, prepared from pearled grain through hammer milling or roller milling, has been incorporated into wheat based products, including bread, cakes, cookies, noodles and extruded snack foods [5]. Wheat bread with barley flour added at 15% to 20% was acceptable in overall flavour, appearance and texture, but an increased proportion of barley flour caused a decrease in loaf volume, dull brown colour and hard crumb texture [6,7]

Numbers of technologies have been developed for the development of different kinds of bread like to design and optimize processing, storage, and transport conditions and methodologies. Many cereals and their products required for bread making are efficient in one way but at the same time are lacking in other properties as well. The demand for the specialty breads and the role of wheat bran, barley and commercial celluloses and other cereals and their products in bread making is increasing day by day. In order to overcome these demands, various methods like blending of different cereals are used for the development of breads with specialized properties. With this background, the present work for the development of leavened bread by using wheat-barley blended flour was undertaken to study the rheological attributes

of wheat-barley blended dough and to prepare breads from composite flours and evaluate the suitable level of barley flour supplementation.

Materials and Methods

The work was carried out in the department of food technology, Islamic University of Science and Technology Awantipora during the year 2010-2011. The wild barley was purchased from Ladakh and then milled in a mixer to obtain whole flour. The flour was stored in plastic air tight containers at refrigerated temperatures until used. Refined wheat flour, shortening, compressed yeast etc were purchased from local market of Srinagar. Composite flours were prepared as mentioned in Table 1. The formulations of the barley enriched breads were according to the Table 2. Breads were prepared from blended flour of wheat and barley according to the procedure shown in Figure 1. MSG was used at 0%, 0.3% and 0.5% level. The ingredients were weighed

Treatment	Concentration of MSG	Wheat flour (%)	Barley flour (%)	
T ₁	T ₁ M ₁	0.0%	100%	0%
	T ₁ M ₂	0.3%	100%	0%
	T ₁ M ₃	0.5%	100%	0%
T ₂	T ₂ M ₁	0.0%	90%	10%
	T ₂ M ₂	0.3%	90%	10%
	T ₂ M ₃	0.5%	90%	10%
T ₃	T ₃ M ₁	0.0%	80%	20%
	T ₃ M ₂	0.3%	80%	20%
	T ₃ M ₃	0.5%	80%	20%
T ₄	T ₄ M ₁	0.0%	70%	30%
	T ₄ M ₂	0.3%	70%	30%
	T ₄ M ₃	0.5%	70%	30%

Table 1: Wheat-barley composite flours used for bread formulations.

*Corresponding author: Nayik GA, Department of Food Engineering and Technology, SLIET, Punjab-148106, India, Tel: +91-9697470552, Fax: 01672-280057; E-mail: gulzarnaik@gmail.com


Received June 21, 2016; Accepted July 08, 2016; Published July 14, 2016

Citation: Nazir F, Nayik GA (2016) Impact of Wheat-Barley Blending on Rheological, Textural and Sensory Attributes of Leavened Bread. J Food Process Technol 7: 611. doi:10.4172/2157-7110.1000611

Copyright: © 2016 Nazir F, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Wheat flour/composite flour	150 g
Yeast	3.0 g
Sugar	6.0g
Salt	1.5g
Shortening	3.0g

Table 2: Wheat-barley composite bread formulations.



accurately and the yeast was activated in hot water. All the ingredients were mixed in a vessel and yeast was added while taking into account the amount of water. The dough was then placed in an incubator at 37°C for fermentation. Dough was taken out after 1 hour and then knocked back to remove the excess gases. The dough was again placed in incubator for fermentation and removed after 30 min, knocked back, rolled and moulded into pans and then allowed to ferment for another 35 min. The pans were then placed in baking oven at 220°C for 30-35min. The breads were taken out, cooled and then sliced. The breads were stored at room temperature.

Rheological and textural analysis

A Texture Analyzer was used to measure the extensibility and firmness (TA. HD. Plus, Stable Micro Systems, Godalming, Surrey, UK). For extensibility, the dough strip was placed onto the grooved region of the sample plate and the plate was inserted into the rig while holding down the spring loaded clamp lever. The handle was released slowly. The analysis was done to a distance of 75 mm, at a pre-test speed of 2.0 mm/s, test speed of 3.3 mm/s, post-test speed of 10.0 mm/s using a 5 Kg load cell. For firmness, the sample was removed from its place of storage and was placed centrally over the supports just prior to testing. The texture profile analysis was done to a distance of 30 mm at pre-test speed of 1.0 mm/s, test speed of 1.7 mm/s and a post- test speed of 10.0 mm/s using a 5 Kg load cell.

Determination of organoleptic characteristics

A panel of 10 judges evaluated the organoleptic characteristics of prepared breads. They assessed crust colour, appearance, flavour, texture, taste and overall acceptability, using a 9-point hedonic rating scale (9-Like extremely, 8-Like very much, 7-Like moderately, 6-Like slightly, 5-Neither like nor dislike, 4-Dislike slightly, 3-Dislike moderately, 2-Dislike very much, 1-dislike extremely). Tap water was provided to the panelists to rinse between evaluations.

Statistical analysis

The data was statistically analysed on a computer using design factorial in Completely Randomized Design (CRD) as suggested by Snedecor and Cochran [8].

Results and Discussion

Table 3 indicated that the mean values for extensibility statistically showed a significant increasing trend from 24.76 to 27.98 with addition of MSG. Further the highest mean extensibility was observed in T₁ (33.51) and lowest mean extensibility was observed in T₄ (18.46). These results are in accordance with the findings of Sternberg et al. [9] who observed increase in extensibility of dough due to the reducing action of MSG. MSG unexpectedly caused weakening of gluten structure for a relatively short period (before fermentation). The extensibility is primarily due to elasticity of glutenin proteins present in wheat, which may dilute with the addition of non-wheat flour resulting in decreased extensibility [10]. The mean values for extensibility showed a significant decreasing trend from 22.39 to 19.18 with addition of MSG (Table 4). The highest mean extensibility was observed in T₁ treatment (28.19) and lowest mean extensibility was observed in T₄ treatment (14.42). Similar results were found by Sternberg et al. [9] who reported that MSG caused strengthening of gluten network after fermentation due to the oxidative action. The decrease in extensibility with addition of barley flour is due to glutenin dilution.

Table 5 indicated that the mean values for firmness elucidated statistically showed a significant increasing trend from 26.60 to 44.73

Treatment	Extensibility			Treatment mean
	M ₁	M ₂	M ₃	
T ₁	32.10	33.01	35.42	33.51
T ₂	26.93	28.18	30.40	28.50
T ₃	22.82	24.07	26.00	24.30
T ₄	17.17	18.10	20.12	18.46
MSG mean	24.76	25.88	27.98	-

CD at 5%; Treatment mean (T) = 0.158; MSG mean (M) = 0.0132; T × M = 0.385; T₁ = Control bread (100% W.F); W.F = Wheat flour B.F = Barley flour; T₂ = 10% B.F: 90% W.F; T₃ = 20% B.F: 80% W.F; T₄ = 30% B.F: 70% W.F; M₁ = 0% MSG; M₂ = 0.3% MSG; M₃ = 0.5% MSG

Table 3: Extensibility (mm) of wheat-barley dough at different concentrations before fermentation.

Treatment	Extensibility			Treatment mean
	M ₁	M ₂	M ₃	
T ₁	30.12	28.34	26.10	28.19
T ₂	25.10	23.15	21.32	23.19
T ₃	19.40	17.42	16.10	17.64
T ₄	14.97	15.12	13.17	14.42
MSG mean	22.39	21.01	19.18	-

CD at 5%; Treatment mean (T) = 0.15; MSG mean (M) = 0.048; T × M = 0.152 T₁ = Control bread (100% W.F); W.F = Wheat flour; B.F = Barley flour; T₂ = 10% B.F: 90% W.F; T₃ = 20% B.F: 80% W.F; T₄ = 30% B.F: 70% W.F; M₁ = 0% MSG; M₂ = 0.3% MSG; M₃ = 0.5%MSG

Table 4: Extensibility (mm) of wheat-barley dough at different concentrations after fermentation.

Treatments	Firmness		
	Day 1	Day 2	Day 3
T ₁ M ₁	7.68	18.11	28.38
T ₁ M ₂	11.00	23.92	36.83
T ₁ M ₃	17.10	27.55	42.17
T ₂ M ₁	10.04	15.20	27.52
T ₂ M ₂	29.54	36.18	43.50
T ₂ M ₃	33.10	41.73	47.02
T ₃ M ₁	13.52	33.78	42.52
T ₃ M ₂	30.28	43.55	46.60
T ₃ M ₃	38.69	48.27	55.31
T ₄ M ₁	28.05	34.51	42.62
T ₄ M ₂	43.62	48.19	53.28
T ₄ M ₃	56.56	62.35	71.03

Table 5: Raw data for change in firmness (N) of bread at different time intervals and with addition of MSG.

Treatment	Firmness			Treatment mean
	Day1	Day2	Day3	
T ₁	11.93	23.19	35.78	23.63
T ₂	24.23	31.04	39.36	31.54
T ₃	27.49	41.84	48.14	39.16
T ₄	42.74	48.35	55.64	48.91
Storage mean	26.60	36.10	44.73	-

CD at 5%; Treatment mean (T) = 3.50; Storage mean (S) = 2.93; T × S = 7.38

Table 6: Change in firmness (N) of bread during storage (Temperature 13°C to 18°C and relative humidity 64% to 63%).

during three days of storage period. Further, the highest mean firmness was observed in T₄ treatment (48.91) and lowest mean firmness was observed in T₁ treatment (23.63). The results are in accordance with the findings of Goesaert et al. [11] who reported increase in firmness of bread during storage due to the transformation of gelatinised starch (amylopectin) network into an extensive, partially crystalline, permanent amylopectin network, with amylopectin crystallites acting as junction zones. The increase in firmness with addition of barley flour is due to gluten dilution. These results are in alignment with the

findings of Gill et al. [12] who further reported that β-glucan in barley flour, when added to wheat flour during bread making, could tightly bind to appreciable amounts of water in the dough, suppressing the availability of water for the development of the gluten network. An underdeveloped gluten network can lead to increased bread firmness.

Table 6 indicated that the mean values for firmness elucidated statistically showed a significant increasing trend from 25.15 to 45.07 during storage with the addition of MSG. Further, the highest mean firmness was observed in T₄ treatment (48.91) and lowest mean firmness was observed in T₁ treatment (23.63). These results are in accordance with the findings of Gill et al. [12] who observed increase in firmness of bread with addition of barley flour due to gluten dilution.

Organoleptic characteristics

Crust colour: The highest value of colour was observed for T₂M₁ (7.5) and the lowest value of colour was observed for T₄M₂ (5.4) (Table 7). As the level of barley was increased in blends, the crust colour of the breads changed from creamy white to dull brown. Similar results were reported by Gupta et al. [13] and Dingra et al. [6].

Appearance: The appearance score for the control breads decreased significantly upon increasing the blending level to 20 and 30% with barley. Among the blended breads, the highest appearance score was observed for T₂M₁ (7.1), whereas the lowest score was observed for T₄M₃ (5.0) (Table 7). These results are in quite similar with the findings of Gupta et al. [13].

Flavor: Results for the flavor of the breads revealed that the highest flavor score was observed for T₂M₂ (7.3), whereas the lowest score was observed for T₄M₁ (5.5). The flavor of barley-blended breads might be affected by the fibrous flavour of barley flour (Table 7). A similar decline in acceptability of breads due to strong flavor was observed by noticed by Dingra et al. [13].

Crust texture: The crust texture was related to the external appearance of the breads, i.e., smoothness or roughness of the crust. Crust texture score also decreased with increase in the substitution of barley and MSG in wheat flour as compared with the control sample. Among the blended bread, the highest score was observed for T₂M₁

Treatments	Colour	Texture	Taste	Flavor	Appearance	Overall acceptability
T ₁ M ₁	7.0	7.1	7.0	7.2	6.8	7.0
T ₁ M ₂	7.0	7.1	7.0	7.0	6.7	7.1
T ₁ M ₃	7.0	7.0	7.0	7.1	6.7	7.1
T ₂ M ₁	7.5	7.2	7.6	7.2	7.1	7.4
T ₂ M ₂	7.4	7.1	7.4	7.3	7.0	7.3
T ₂ M ₃	7.4	7.0	7.3	7.2	7.0	7.2
T ₃ M ₁	6.5	6.4	6.0	6.0	6.2	6.3
T ₃ M ₂	6.4	6.2	6.1	6.2	6.0	6.2
T ₃ M ₃	6.3	6.1	6.2	6.3	6.1	6.2
T ₄ M ₁	5.8	5.3	5.1	5.5	5.3	5.4
T ₄ M ₂	5.4	5.2	5.0	5.7	5.1	5.3
T ₄ M ₃	5.5	5.0	5.1	5.6	5.0	5.1

Table 7: Effect of blending on sensory quality of breads.

(7.2), whereas the lowest score was observed for T₄M₃ (5.0) (Table 7). Carson et al. [14] observed similar deterioration in the texture of wheat bread on supplementation.

Taste: Taste evaluation suggested that control and various supplemented breads had most satisfactory taste scores up to the 10% level. Results indicated that the highest taste score was observed for T₂M₁ (7.6), whereas the lowest score was observed for T₄M₂ (5.0) (Table 7). The decrease in taste score was because of the different flavor of flour blends [13].

Overall acceptability: The results showed that the overall acceptability score of all the supplemented breads at the 10% level was at par with the control (Table 7). Breads made from wheat and barley flour up to the 10% level were found acceptable, but at more than 10% substitution, the overall acceptability score was significantly reduced as compared with the control. In overall profile, the flavor of the breads was malty and fibrous at the 30% level of substitution. Based on the above results, breads containing 10% barley flour were found to be most acceptable by the sensory panelists.

Conclusion

The studies revealed that before fermentation the extensibility of the dough showed significant increase with addition of MSG and significant decrease with addition of barley flour. After fermentation the extensibility of the dough showed a significant decreasing trend with addition of barley flour and MSG. Firmness of bread showed a significant increasing trend with addition of barley, MSG and with storage. It was concluded from the present studies that breads made from 10% barley flour blend proved superior with respect to sensory characteristics and were better than other blended breads as far as firmness and extensibility was concerned. Further investigations are required to evaluate the compositional and the antioxidant properties of these breads.

Acknowledgement

The first author is thankful to Department of Food Technology, IUST Awantipora, Pulwama J&K, India for providing necessary facilities for lab work.

References

- Baik B, Ullrich C (2008) Barley for food: Characteristics, improvement and renewed interest. J Cereal Sci 48: 233-242
- Czuchajowska Z, Klamczynski A, Paszczyńska B, Baik BK (1998) Structure and functionality of barley starches. J Cereal Chem 75: 747-754.
- Izydorczyk MS, Storsley J, Labossiere D, MacGregor AW, Rossnagle BG, et al. (2000) Variation in total and soluble β -glucan content in hullless barley: Effects of thermal, physical, and enzymatic treatments. J Agri Food Chem 48: 982-989.
- Quinde Z, Ullrich SE, Baik BK (2004) Genotypic variation in colour and discolouration potential of barley-based food products. J Cereal Chem 81: 752-758.
- Newman RK, Newman CW (1991) Barley as a food grain. J Cereal Food World 36: 800-805.
- Dhingra S, Jood S (2004) Effect of flour blending on functional, baking and organoleptic characteristics of bread. Int J Food Sci Technol 39: 213-222.
- Malik H, Nayik GA, Dar BN (2015) Optimisation of process for development of nutritionally enriched multigrain bread. J Food Process Technol 7: 544.
- Snedecor GW, Cochran WG (1967) Statistical methods. The Iowa State University Press, USA.
- Sternberg G, George P (1976) Preparation of yeast leavened dough products. J Food Addit 44: 100-110
- Khatkar BS (2004) Dynamic rheological properties and breadmaking qualities of wheat gluten: Effects of urea and dithiothreitol. J Sci Food Agri 85: 337-341.
- Goesaert H, Slade L, Levine H, Delcour AJ (2009) Amylase and bread firming-an integrated view. J Cereal Sci 50: 345-352.
- Gill S, Vasanthan T, Oraikul B, Rossnagle B (2002) Wheat bread quality as influenced by the substitution of waxy and regular barley flours in their native and extruded forms. J Cereal Sci 36: 219-237.
- Gupta M, Bawa AS, Semwal AD (2011) Effect of barley flour blending on functional, baking and organoleptic characteristics of high-fiber rusks. J Food Process Preserv 35: 46-63.
- Carson L, Setser C, Sun XS (2000) Sensory characteristics Food Sci Technol 35: 465-471.

Citation: Nazir F, Nayik GA (2016) Impact of Wheat-Barley Blending on Rheological, Textural and Sensory Attributes of Leavened Bread. J Food Process Technol 7: 611. doi:10.4172/2157-7110.1000611