

## Impaired Cardiac Function in Cardiac-Specific SIRT1-Deficient Mice

Feng Cao\*

Chief Physician, Cardiology Department, Chinese PLA General Hospital, Beijing, P.R.China

\*Corresponding author: Feng Cao, Associate Director, Chief Physician, Cardiology Department, Chinese PLA General Hospital, 28# Fuxing Street, Beijing, 100853, P.R.China, Tel: 86-10-55499449; E-mail: [wind8828@gmail.com](mailto:wind8828@gmail.com)

Rec date: Nov 01, 2014; Acc date: Nov 02, 2014; Pub date: Nov 04, 2014

Copyright: © 2014 Cao F. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Description

Cardiac specific SIRT1 knockout mice were generated through the Cre/loxP system. Briefly, loxP-SIRT1 mice and Myh6-Cre mice were crossed to get SIRT1<sup>+/-</sup> mice. Then, SIRT1<sup>+/-</sup> female mice were mated with SIRT1<sup>+/-</sup> adult males to generate SIRT1<sup>-/-</sup> mice. RT-PCR revealed the gene type of SIRT1<sup>+/-</sup> and SIRT1<sup>-/-</sup> mice (A).

The low expressions of SIRT1 in SIRT1<sup>+/-</sup> and SIRT1<sup>-/-</sup> mice were confirmed by Western Blot (B). The heart/body weight was increased only in SIRT1<sup>-/-</sup> mice.

Echocardiography results (D) showed that cardiac function was impaired in SIRT1-deficient mice, with the decreased ejection fraction (EF) in SIRT1<sup>-/-</sup> mice (E).

