

Impairment of Pulmonary Vasoreactivity in Response to Endothelin-1 In Patients With Chronic Obstructive Pulmonary Disease (COPD)

Duong-Quy S^{1,2*}, Dao P³, Hua-Huy T¹, Le-Dong NN⁴ and Dinh-Xuan AT¹

¹Department of Respiratory Physiology, Cochin Hospital, University Paris Descartes, Sorbonne Paris Cité, Paris - France

²Bio-Medical Research Center, Lam Dong Medical College, Dalat – Vietnam

³Antony Hospital, Antony - France

⁴Department of Pulmonology, St. Elisabeth Hospital, Namur - Belgium

Abstract

Background: Pulmonary hypertension secondary to COPD is characterized principally by impairment of vasoreactivity and vascular remodeling. While the structural changes have been described in all stages of COPD, the impairment of vasoreactivity in response to ET-1, a potent vasoconstrictor, has not been well understood.

Materials and methods: Proximal pulmonary arteries were obtained from non-smokers, smokers with normal lung function, and COPD patients who underwent lung resection for other diseases (n=6 in each group). Pulmonary arterial contraction induced by ET-1 was assessed without or with the presence of ET-1 receptor antagonists (BQ-123 and BQ-788). Expressions of ET-1 receptors were measured by immunohistochemistry, western blot, and qRT-PCR.

Results: ET-1-induced pulmonary arterial contraction increased significantly in COPD patients in comparison with control subjects and non-COPD smokers (P<0.05) and the use of ET-1 receptor antagonists could not prevent this contraction. The hypercontraction was not associated with the increase of ET-1 and ET-1 receptors expression in pulmonary arterial vessels.

Conclusion: In patients with COPD, pulmonary arterial contraction induced by ET-1 has increased significantly and could not be prevented by using ET-1 receptor antagonist alone. This alteration might be related to the upregulation of other signalling pathways in pulmonary arterial vessels of COPD patients.

Keywords: COPD; ET-1; ET-1 receptor antagonists; BQ-123; BQ-788

Introduction

In patients with chronic obstructive pulmonary disease (COPD), the impairment of vasoreactivity in response to endothelium-derived constricting factor (endothelin-1, thromboxane) and relaxing factors (nitric oxide, prostacyclin) has not been well understood. This impairment might be due to endothelial dysfunction and might promote to the alteration of pulmonary vascular tone and pulmonary vascular hypertrophy in these patients [1]. Among these vasoreactivity factors, endothelin-1 (ET-1), the most potent endogenous vasoconstrictor peptide of endothelin family, produced by endothelial cells, has an important role in the modulation of vascular resistance.

Moreover, the abnormalities in endothelin-1 (ET-1) metabolism have been also reported in COPD patients. Previous studies showed that the concentration of ET-1 in plasma, in urinary excretion, and in saliva has increased significantly in COPD patients [2]. Further, at local level, the increase of endothelin expression has been founded in pulmonary vessels in patients with pulmonary hypertension secondary to COPD [3]. Although the effect of cigarette smoke on systemic vascular tone due to ET-1 by increasing the vasoconstriction has been well studied [4], the vasoreactivity of pulmonary vessels in response to ET-1 in healthy smokers and especially in smokers with COPD has not been well understood.

Generally, ET-1 exerts its vasoactive properties by binding to its specific receptor ET_A presenting in pulmonary arterial smooth muscle cells (PASCs), and ET_B expressing in endothelial cells (ECs) and in PASCs. Under normal conditions, ET-1 mediates Ca²⁺ releasing

from sarcoplasmic reticulum, (PKC-dependent inhibition of voltage-gated K⁺ (K_v) channels), and subsequent membrane depolarization and Ca²⁺ influx through voltage-dependent calcium channels [5-7]. This increase in intracellular calcium concentration ([Ca²⁺]_i) activates myosin light chain kinase (MLCK), leading to phosphorylation of myosin light chains and vasoconstriction [8, 9]. In chronic hypoxia exposure, where the ET-1 sensitivity and its receptor ET-A have been upregulated [10-12], ET-1-induced vasoconstriction is independently of intracellular calcium concentration and related mostly to Ca²⁺-sensitization [13]. Therefore, Ca²⁺-sensitization regulates vascular tone by increasing phosphorylation of myosin regulatory light chain (MLC₂₀) [14]. The use of ET-1 receptor antagonists may prevent the vasoconstriction in hypoxia-induced pulmonary hypertension.

The aims of this study were to evaluate the vasoreactivity of pulmonary arteries in smokers with and without COPD in response to exogenous ET-1 and its receptor inhibitors.

***Corresponding author:** Duong-Quy S, Cardiology – Respiratory Department, Cochin Hospital, Paris Descartes University, Paris – France, Tel: +33.679193377; E-mail: sduongquy.jfvp@gmail.com

Received May 28, 2015; Accepted July 20, 2015; Published July 22, 2015

Citation: Duong-Quy S, Dao P, Hua-Huy T, Le-Dong NN, Dinh-Xuan AT (2015) Impairment of Pulmonary Vasoreactivity in Response to Endothelin-1 In Patients With Chronic Obstructive Pulmonary Disease (COPD). J Vasc Med Surg 3: 213. doi:[10.4172/2329-6925.1000213](https://doi.org/10.4172/2329-6925.1000213)

Copyright: © 2015 Duong-Quy S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Methods

Subjects and tissue preparations

Subjects, who underwent lung resection for other diseases, enrolled in the study. Three groups of subjects were constituted: non-smokers (control subjects), healthy smokers (non-COPD smokers) and smokers with COPD (COPD patients). COPD were defined and classified by the international standard criteria [ratio of FEV₁ (forced-expiratory volume in one second)/FVC (forced vital capacity) <0.7] [15]. The study was approved by French Ethnical Committee and taken place in UPRES25511 – Paris Descartes University. All patients were informed about the aims of study and written informed consent was obtained from each subjects.

Proximal pulmonary arterial vessels were dissected precisely far away from local damaged tissues at operation room. Pulmonary arterial rings of 4 to 6 mm external diameter and 3 mm of length, cleaned of perivascular tissue, were dissected and immediately putted in Krebs-Henseleit solution at 4°C for vasoreactivity study within less than 30 minutes. The others dissected pulmonary arterial vessels were conserved at -80°C for proteins and RNA extractions or fixed in 4% formalin and embedded in paraffin for immunohistochemistry study.

Chemical reagents and antibodies

All chemical reagents used in the present study were purchased from Sigma-Aldrich (Sigma-Aldrich France; St Quentin En Fallavier, France) unless otherwise noted. Primary antibody anti-ET_A and ET_B were purchased from Assay Designs Inc (Ann Arbor, USA).

For all washing steps in immunohistochemical staining and Western blot, phosphate buffered saline (PBS, pH 7.4) was used with Tween 20 (0.05%), noted as PBS-T. Primary antibodies and HRP-conjugated secondary antibodies for Western blot were purchased from Santa Cruz Biotechnology Incorporation (Santa Cruz, CA, USA), biotinylated secondary antibodies for immunohistochemical staining from Vector Laboratories (Burlingame, CA, USA), unless differently mentioned. Antibodies used in western blotting were diluted in non-fat milk (5%), Tween-20 (0.05%) in PBS and those for immunohistochemistry in PBS.

Measurement of Pulmonary Vascular Contraction

Pulmonary arterial ring was suspended horizontally between two parallel stainless steel wires for the measurement of isometric tension by organ chambers (Emka Technologies, Paris, France). Pulmonary arterial ring was dipped in a fresh Krebs-Henseleit solution with the following composition: glucose 10 mM, pyruvate 2 mM, HEPES 10 mM, EDTA 0.03mM, NaCl 118 mM, KCl 4.7 mM, CaCl₂ 2.5 mM, KH₂PO₄ 1.2 mM, MgSO₄ 1.2 mM, NaHCO₃ 15 mM, and the solution was continuously oxygenated with 95% O₂-5%CO₂ (Air Liquide Santé; Paris, France) and maintained at 37°C.

The variations of force generated by the ring segment were measured and recorded by IOX-Software (Emka Technologies, Paris, France). Pulmonary arterial rings were stabilized under controlled condition at 1.0 g of resting tension (for 10 min before stretching progressively at 1.5 g to obtain the pre-tension values as determined by experimental assessments), and it was corresponded to optimal resting tension for isometric condition registry. After the initial equilibration period, the vessels were exposed to a maximally effective concentration of 80 mM KCl. When a stable tension plateau developed, the KCl was removed by several washes and a period of 30 min was allowed

for reequilibration. Then, pulmonary arterial rings were exposed to increasing and cumulative doses of ET-1 (10⁻¹⁰ to 10⁻⁷M), in the present of inhibitors of ET_A receptor (BQ-123 at 10⁻⁶M) or ET_B receptor (BQ-788 at 10⁻⁶M). Other pulmonary arterial rings of the same patients were incubated with receptor inhibitors during at least 1 hour before starting the measurement of ET-1 dose-response curve. The isometric contractions were normalized by expressing force developed per cross-sectional area (g/mm²), which took into account the variation in vascular sizes.

The integrity of pulmonary arterial endothelial cells was tested by the vasodilatation endothelium-dependent in response to 10⁻⁶M acetylcholine (Ach) at the end of ET-1 stimulation protocol and when the plateau of contraction curve had been obtained at 10⁻⁷M. The relaxation of more than 30% compared to the value before acetylcholine response was necessary to confirm the endothelial intact. The ET-1 response-curve had been presented as the percentage of maximal contraction induced by 80mM potassium chloride (KCl). For each patient, four pulmonary arterial rings were studied.

Immunohistochemistry for ET_A and ET_B receptors

For immunohistochemical staining of ET_A and ET_B receptors, after re-hydration by successive baths of decreasing ethanol concentrations, all pulmonary arterial sections had been submerged in citrate buffer (10 mM citrate, 0.05% Tween, and pH 6.0) and unmasked with heat treatment in micro-oven (750W × 3 times). Endogenous peroxidase activity was neutralized by incubating with oxygenate water of 3% during 15 minutes. The non-specific endogenous binding sites had been blocked by 2% BSA (bovine serum albumin) in 45 minutes. After washing steps, pulmonary arterial sections had been incubated for one hour with primary antibodies, which had been diluted to 1/50 for anti-ET_A and 1/25 for anti-ET_B in PBS-T of 0.05% and 1% BSA as titration essays. Negative-control sections were incubated in blocking buffer alone without primary antibody. The sections had been then incubated with biotinylated secondary antibody at 1/200 dilution within 45 minutes. After successive washing steps, pulmonary arterial sections were incubated with preformed avidin-peroxydase RTU Vectastain Kit (Vector Laboratory; Burlingame, USA) before revealed with metal enhanced DAB peroxidase substrate (3, 3'-diaminobenzidine) until the desired staining was obtained (4 minutes for ET_A and 5 minutes for ET_B). Tissue sections were then counterstained with hematoxylin-Gill solution and examined using light microscopy. ET_A and ET_B immunostaining expression was measured in endothelial and SMC cells of proximal pulmonary arteries. The staining expression was measured by using intensity score. At least ten counted images were performed for each patient.

Real time PCR for mRNA of ET-1, ET-1 conversion enzyme, and ET_A and ET_B receptors

Pulmonary arteries for real time PCR experiments were frozen in liquid nitrogen and stored at -80°C. The samples were homogenated and realized ARN extraction in RNase-free condition by using combined phenol-guanidine isothiocyanate buffer (Trizol[®], Life technologies, CA, USA). Each sample had been homogenated in Trizol at 4°C with Ultraturrax (T25, Staufen, Germany), centrifuged at 12000 × g within 10 minutes to eliminate tissue debris, and then added with chloroform (Merck, Darmstadt, Germany) to the lysate. After precipitated with isopropanol (Merck, Darmstadt, Germany) and centrifugated at 14000 × g (10 min), RNA pellet had been washed with 70% ethanol (Merck,

Hohenbrunn, Germany) and centrifugated at 12000 × g. Pellet had been then redissolve in RNase-free water and quantified with optical density at 260 nm.

After RNA isolation, reverse transcription was carried out in mixed solution of 7.5 μM hexanucleotide, 9 mM dithiothreitol, 20 U RNase inhibitor (Perkin-Elmer), 4 dNTP with the concentration of 220 μM for each (Boehringer), and 50 U Reverse Transcriptase (Superscript; Gibco BRL). It had been placed in PCR apparatus (Perkin-Elmer Gene Amp PCR system 2400, CA, USA) with following cycles: 10 min at 22°C, 1 hour at 42°C, and 5 min at 99°C. Real time PCR was performed with the real time PCR GeneAmp machine 7700. The GeneAmp machine 7700 sequence detection system monitored the binding of a fluorescent dye to double-strand DNA by real time detection of the fluorescence during each cycle of PCR amplification. Control samples without the reverse-transcription step and no added RNA were also included in each plate to detect any possible contamination. The housekeeping genes, HPRT (Hypoxanthine PhosphoRibosyl Transférase) was used as references. The real time PCR reaction was performed at a temperature of 50°C for 2 min, 95°C for 10 min, and the following 40 PCR cycles with 95°C for 15 s and 60°C for 1 min, and fluorescence was measured after amplification step.

The sequences of the primer pairs were designed as follows: ET-1: forward 5'-CTTCTGCCACCTGGACATCA- 3' and reverse 5'-GGCTTCCAAGTCCATACGGA-3'; ET_A receptor: forward 5'-TG-GTGTGCACTGCGATCTTC-3' and reverse 5'-GCAATTCT-CAAGCTGCCATTC-3'; ET_B receptor: forward 5'-TGGCG-TATTTCTTTGCAAGCT-3' and reverse 5'-AGGTTGAGGACGGT-GATCCC-3'; ECE-1: forward 5'-TGGGGGACCTTCAGCAACCT-3', reverse 5'-GGGTGCTGGAGTTGCTCTTG-3'; HPRT: forward 5'-TCAGGCAGTATAATCAAAGATGGT-3' and reverse 5'-AGTCTGGCTTATATCCAACACTTCG-3'.

At the end of each PCR run, the data were automatically analyzed by the system and an amplification plot was generated for each DNA sample. Quantitative real-time PCR values are expressed as the fold change of the target-gene expression relative to HPRT mRNA in each sample using the following formula: fold change = 2^{-ΔΔC_T}, where ΔC_T = C_{Target gene} - C_{HPRT} and -ΔΔC_T = ΔC_{Test sample} - ΔC_{Control}. The experimental approach was further validated by the observation that the differences between the C_T for the target gene and β-actin remained essentially constant for each starting DNA amount.

Western blot analysis for ET-1 receptors (ET_A, ET_B)

Frozen pulmonary arterial samples were homogenized in ice-cold RIPA buffer (Cell Signalling Technology, Danvers, USA). The homogenates were centrifuged at 21000 g at 4°C for 20 minutes to remove tissue debris. Protein contents of homogenated lysates were measured by BCA Protein assay kit (Pierce Biotechnology, Rockford, IL, USA). Equal amounts of lysates (30 μg of total protein content per sample), were subjected to electrophoresis on SDS-PAGE gels (12% for ET_A and ET_B), and then transferred to PVDF membranes (Immobilon-P, Millipore, MA, USA). Membranes were blocked in 5% non-fat milk, 0.1% Twen-20 in PBS for 1 hour at room temperature and then hybridized overnight at 4°C with primary antibodies diluted at 1/500. After being washed in PBS-T, membranes were incubated with HRP-linked goat anti-rabbit, diluted at 1/10000. Protein bands were developed on the film by using Enhanced ChemiLuminescence Plus reagent, according to the manufacturer's instructions (Amersham, Biosciences, Orsay, France). The protein density was quantified by using Image Software System (Genius 2). The equal amounts of proteins

loaded per well were verified by stripping the blots with Tris-HCl (pH 6.7) 62.5 mM, SDS 2%, and β-mercapto-ethanol 100mM for 30 minutes at 55°C. The membranes were reprobed with a mouse polyclonal anti-β-actin antibody (1/1000; sc-9104, Santa Cruz Biotechnology) and then incubated with HRP-linked goat anti-rabbit, diluted at 1/10000. Results were expressed as ratios of ET_A and ET_B/β-actin.

Statistical analysis

Statistical comparisons were performed with non parametric Kruskal-Wallis test and with analysis of variance (ANOVA) among more than three groups by using the SPSS 16.0 software (Chicago, IL). Data were presented as means ± SD. Differences were considered significant with P < 0.05.

Results

Clinical characteristics

Three groups of patients who met the inclusion criteria were included in the present study (6 subjects for each group). The clinical and functional characteristics of the patients are presented in Table 1. The tobacco consumption in COPD patients was higher than non-COPD smokers (P<0.05). The non-COPD smokers had a normal lung function with FEV₁ = 97 ± 5% and FEV₁/FVC=75 ± 4%. COPD patients had moderate airflow obstruction with FEV₁=73±15% and FEV₁/FVC=64±6%. All the patients of three groups had a normal gas blood exchange.

Vasoconstriction induced by ET-1

The results of contraction induced by ET-1 with increasing concentration (10⁻¹⁰-10⁻⁷ M) in organ bath chambers showed that in compare to Emax (contraction induced by 80 mM KCl) and with the high dose (10⁻⁸-10⁻⁷ M), the percentage of maximal contraction was significantly higher in COPD patients than controls and non-COPD smokers (P<0.05; Figure 1). Whereas, with the lose dose (10⁻¹⁰-10⁻⁹ M), there were not any significant differences between three groups in response to ET-1 (P>0.05; Figure 1).

Effect of ET_A and ET_B antagonists in ET-1-induced contraction


To determine the role of ET_A or ET_B receptor involved in the ET-1 induced-contraction, in the present study, the pulmonary arterial rings had been exposed to selective antagonist of ET_A receptor (BQ-123 at 10⁻⁶M) and ET_B receptor (BQ-788 at 10⁻⁶M) in organ bath chambers.

In control group, in the presence of BQ-123 and BQ-788, the percentage of pulmonary arterial contraction induced by ET-1 was

Variables	Control (non smoking) (n = 6)	Smokers non-COPD (n = 6)	COPD (n = 6)	P
Age, years	59 ± 7	50 ± 10	67 ± 12	NS ^{†,‡}
Male/Female	2/4	3/3	3/3	
Tobacco, pack-year	-	30 ± 13	41 ± 23	< 0.05 [‡]
FEV ₁ , % pred	97 ± 5	97 ± 5	73 ± 15	< 0.05 ^{†,‡}
FEV ₁ /FVC	86 ± 5	75 ± 4	64 ± 6	NS, < 0.05 ^{†,‡}
FVC, % pred	105 ± 17	104 ± 9	87 ± 22	NS ^{†,‡}
PaO ₂ , mmHg	86 ± 11	87 ± 17	83 ± 14	NS ^{†,‡}
PaCO ₂ , mmHg	38 ± 2	37 ± 2	37 ± 5	NS ^{†,‡}

(%pred : percentage of predicted value; FEV₁ indicates forced expiratory volume in one second; FVC, forced vital capacity; *: non-COPD smokers vs controls; †: COPD vs control subjects; ‡: COPD vs non-COPD smokers; NS: no significant)

Table 1: Clinical characteristics of study subjects.



significantly reduced in comparison without antagonists ($P<0.05$; Figure 2a). In non-COPD smokers, the contraction had been reduced significantly in the presence of BQ-123. Whereas, in the presence of BQ-788, the pulmonary arterial contraction induced by ET-1 had two phases: reduced the contraction at low dose (10^{-10} - 10^{-8} M), and increased the contraction at high dose (10^{-8} - 10^{-7} M); but there were not any significant differences in comparison with ET-1-induced contraction without BQ-788 ($P>0.05$; Figure 2b). In COPD patients, the reductions of pulmonary arterial contraction induced by ET-1 in the presence of BQ-123 and BQ-788 were not significantly different in comparison with ET-1-induced contraction without antagonists ($P>0.05$; Figure 2c).

Semi-quantitative and quantitative assessment of ET_A and ET_B receptors

Immunohistochemistry: The results of immunostaining ET_A and ET_B receptor expression in pulmonary arterial sections showed that there were no significant differences between non-COPD and COPD smokers for expression intensity per surface unit assessment and in comparison with control group ($P>0.05$; Figure 3a).

Western blot: The protein expression of ET_A and ET_B receptor measured by densitometry showed that there were no significant differences between three groups for protein levels ($P>0.05$; Figure 3b).



Real time quantity-PCR (RTq-PCR): Although the contractions of pulmonary arterial rings induced by ET-1 were different between non-COPD and COPD smokers, the results of RTq-PCR showed that there were no significant differences between three groups for the transcription of mRNA ($P>0.05$; Figure 3c).

The relative expressions of ET-1 and ET-1 enzyme conversion (EEC) were not significant different between three groups. Although the mRNA expression of ET-1 and EEC in non-COPD smokers were slightly higher than controls and COPD patients, but there were not any significant differences ($P>0.05$; data not shown).

Discussion

The aim of this study was to elucidate the role of ET_A and ET_B receptors in pulmonary arteries responding to ET-1 induced vasoconstriction in non-COPD smokers and in COPD patients. The results of present study showed that: (1) in COPD patients, the pulmonary vasoconstriction induced by ET-1 was significantly increased in comparison with non-smokers and non-COPD smokers; (2) ET-1-induced vasoconstriction was mediated by ET_A and ET_B in non-smokers and predominantly by ET_A in non-COPD smokers; (3) in COPD patients, vasoconstriction induced by ET-1 was not completely inhibited by selective ET-1 receptor antagonists.

Until now, the effect of cigarette smoking on the tonus of






Figure 2c: ET-1-induced contraction of pulmonary arterial rings with increased dose in the presence of ET_A (BQ-123) and ET_B (BQ-788) antagonists in COPD patients (4 measurement for each patient). Emax is represented as a contraction level induced by 80 mM of KCl. Values are means ± SE. NS: no significant differences with P>0.05 for BQ-123 and BQ-788 vs without antagonists.

the endothelium-derived contracting factors, ET-1 has an important role. The previous studies showed that the plasma concentration of ET-1 and the expression of ET_A receptor have been increased in chronic hypoxic exposure and in COPD with hypoxia, and it was associated with the alteration of ET-1-induced vasoconstriction [17-19]. The mechanism by which ET-1 is upregulated in COPD has not been completely understood. It might be partially mediated by oxidative stress due to cigarette smoke or by alveolar hypoxia when the disease is in progress. In the present study, although ET-1-induced pulmonary vasoconstriction had been found in all the three groups, the vasoconstriction in response to ET-1 at high doses ($\geq 10^{-8}$ M) was significantly higher in COPD patients than control subjects and non-COPD smokers (Figure 1). Whereas, ET-1-induced vasoconstriction in non-COPD smokers was slightly higher than control, but there were no significant difference (Figure 1). These results suggested that the contractility of pulmonary arteries with ET-1 was higher in COPD patients.

In the present study, the use of ET-1 antagonists reduced significantly the contraction induced by ET-1 in non-smokers (BQ-123 for ET_A and




Figure 3a: Expression of ET_A and ET_B receptors measured by immunohistochemistry in pulmonary arterial vessels in control, non-COPD smokers, and COPD patients (magnification $\times 10$). There were no significant differences between three groups (n=6 in each group). A: adventi; B: media. NS: no significant difference vs. control subjects with P>0.05.

pulmonary arterial vessels induced by ET-1 in COPD has not been completely clarified. In COPD, the pulmonary vascular tone has been changed progressively due to the disturbance of vasodilatation – vasoconstriction balance. This event starts at early stage of disease and becomes more severe in advanced stage patients where the alveolar hypoxia and vascular remodelling has been established [16]. Among




Figure 3b: Expression of ET_A and ET_B receptors measured by western blot in pulmonary arterial vessels in control, non-COPD smokers, and COPD patients. There were no significant differences between three groups (n=6 in each group). NS: no significant difference vs. control subjects with P>0.05.




Figure 3c: Expression of ET_A and ET_B mRNA quantified by RT-PCR in pulmonary arterial vessels in control, non-COPD smokers, and COPD patients. There were no significant differences between three groups (n=6 in each group). NS: no significant difference vs. control subjects with P>0.05.

BQ-788 for ET_B) and in non-COPD smokers (BQ-123 for ET_A) (Figure 2a-b). But the contractions were not different in COPD patients (Figure 2c). It confirmed that in non-smokers, the vasoconstriction responded to ET-1 was completely mediated by two receptor types (ET_A and ET_B) and the selective antagonists abolished this effect. However, in non-COPD smokers, ET_A receptor had a predominant role in ET-1-induced pulmonary vasoconstriction (Figure 2b). Particularly, in COPD patients, although the contraction induced by ET-1 at a high dose was significantly higher than non-smokers and non-COPD smokers, ET-1 receptor antagonists could not reduce significantly the contraction (Figure 2c). In addition, in the present study, the mRNA expression of ET-1 and its receptors ET_A and ET_B were not different between three groups (Figure 3a-c). It suggested that in COPD, the pulmonary arterial contraction in response to ET-1 was not related directly to the upregulation of ET_A and ET_B receptors. In our study, we could not measure the plasma concentration of ET-1 of study patients to assess the correlation between the level of mRNA-ET-1 expression in pulmonary arterial vessels and the plasma concentration of ET-1. As discussed above, the plasma concentration of ET-1 was increased and associated with the upregulation of ET_A expression and the alteration of vasoconstriction in COPD with hypoxemia. These data may explain why in our study, gene expression of ET-1 and its receptors was not increased significantly in COPD without hypoxemia.

In this study, the use of selective ET-1 receptor antagonists did not decrease the contraction of pulmonary arterial vessels induced by ET-1 in non-COPD smokers (for BQ788) and in COPD patients (for both BQ123 and BQ789) suggested that another pathway had been implicated in this phenomena. Under normal condition, vasoconstriction induced by ET-1 is mediated by Ca²⁺ influx through L-type Ca²⁺ channels and Ca²⁺ release from the sarcoplasmic reticulum [20]. In some conditions such as in hypoxia exposure, ET-1-induced contraction is independent of changes in [Ca²⁺]_i [21], and known as Ca²⁺ sensitization of the contractile apparatus.

In summary, the results of present study might be explained that in non-smokers (control subjects), ET-1-induced pulmonary arterial contraction was mediated by ET_A and ET_B receptors and this contraction was inhibited by the use of either ET_A or ET_B receptor antagonist. In non-COPD smokers, ET-1-induced pulmonary arterial contraction was mediated principally via ET_A receptor. Whereas, in COPD patients, either ET_A or ET_B receptor antagonist could not decrease significantly the contraction induced by ET-1. In these patients, the ET-1-induced pulmonary arterial contraction was not mediated via ET_A or ET_B receptor. In these patients, when stimulated with ET-1, the vasoconstriction might be mediated by another signaling pathway such as RhoA/Rho-kinase pathway. The activity of this pathway has been upregulated in healthy smoker and smokers with COPD [22, 23] and the use of Rho-kinase inhibitor might attenuated significantly the effect of ET-1 [24].

Conclusion

Although carried out in the limit of patient number, the present study demonstrated that in patients with COPD, the contraction of pulmonary arterial vessels induced by ET-1 was not completely mediated via ET-1 receptors. Especially, in these patients, the hypercontraction induced by ET-1 was not associated with the upregulation of the mRNA expression of ET-1. However, others studies must be done to clarify the role of other pathways in ET-1-induced pulmonary arterial vasoconstriction in patients with COPD.

References

1. Duong-Quy S (2013) Chronic Smoking and Vascular Disease: What Can we Hope for the Future? *J Vasc Med Surg* 1: 113.

2. Celik G, Karabiyikaylu G (1998) Local and peripheral plasma endothelin-1 in pulmonary hypertension secondary to chronic obstructive pulmonary disease. *Respiration* 65: 289-294.

3. Giaid A, Yanagisawa M, Langleben D, Michel RP, Levy R, et al. (1993) Expression of endothelin-1 in the lungs of patients with pulmonary hypertension. *N Engl J Med* 328: 1732-1739.

4. Kiowski W, Linder L, Stoschitzky K, Pfisterer M, Burckhardt D, et al. (1994) Diminished vascular response to inhibition of endothelium-derived nitric oxide and enhanced vasoconstriction to exogenously administered endothelin-1 in clinically healthy smokers. *Circulation* 90(1):27-34

5. Salter KJ, Wilson CM, Kato K, Kozlowski RZ (1998) Endothelin-1, delayed rectifier K channels, and pulmonary arterial smooth muscle. *J Cardiovasc Pharmacol* 31 Suppl 1: S81-83.

6. Shimoda LA, Sylvester JT, Booth GM, Shimoda TH, Meeker S, et al. (2001) Inhibition of voltage-gated K(+) currents by endothelin-1 in human pulmonary arterial myocytes. *Am J Physiol Lung Cell Mol Physiol* 281: L1115-1122.

7. Shimoda LA, Sylvester JT, Sham JS (2000) Mobilization of intracellular Ca(2+) by endothelin-1 in rat intrapulmonary arterial smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* 278: L157-164.

8. Somlyo AP, Somlyo AV (1994) Signal transduction and regulation in smooth muscle. *Nature* 372: 231-236.

9. Somlyo AP, Somlyo AV (2000) Signal transduction by G-proteins, rho-kinase and protein phosphatase to smooth muscle and non-muscle myosin II. *J Physiol* 522 Pt 2: 177-185.

10. Eddahibi S, Raffestin B, Braquet P, Chabrier PE, Adnot S (1991) Pulmonary vascular reactivity to endothelin-1 in normal and chronically pulmonary hypertensive rats. *J Cardiovasc Pharmacol* 17 Suppl 7: S358-361.

11. McCulloch KM, Docherty C, MacLean MR (1998) Endothelin receptors mediating contraction of rat and human pulmonary resistance arteries: effect of chronic hypoxia in the rat. *Br J Pharmacol* 123: 1621-1630.

12. Resta TC, Walker BR (1996) Chronic hypoxia selectively augments endothelium-dependent pulmonary arterial vasodilation. *Am J Physiol* 270: H888-896.

13. Shimoda LA, Sham JS, Shimoda TH, Sylvester JT (2000) L-type Ca(2+) channels, resting [Ca(2+)]_i, and ET-1-induced responses in chronically hypoxic pulmonary myocytes. *Am J Physiol Lung Cell Mol Physiol* 279: L884-894.

14. Somlyo AP, Somlyo AV (2003) Ca²⁺ sensitivity of smooth muscle and nonmuscle myosin II: modulated by G proteins, kinases, and myosin phosphatase. *Physiol Rev* 83: 1325-1358.

15. Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, et al. (2007) Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 176: 532-55.

16. Santos S, Peinado VI, Ramirez J, Melgosa T, Roca J, et al. (2002) Characterization of pulmonary vascular remodeling in smokers and patients with mild COPD. *Eur Respir J* 19: 632-638.

17. Eddahibi S, Raffestin B, Braquet P, Chabrier PE, Adnot S (1991) Pulmonary vascular reactivity to endothelin-1 in normal and chronically pulmonary hypertensive rats. *J Cardiovasc Pharmacol* 17 Suppl 7: S358-361.

18. McCulloch KM, Docherty C, MacLean MR (1998) Endothelin receptors mediating contraction of rat and human pulmonary resistance arteries: effect of chronic hypoxia in the rat. *Br J Pharmacol* 123: 1621-1630.

19. Resta TC, Walker BR (1996) Chronic hypoxia selectively augments endothelium-dependent pulmonary arterial vasodilation. *Am J Physiol* 270: H888-896.

20. Pollock DM, Keith TL, Highsmith RF (1995) Endothelin receptors and calcium signaling. *FASEB J* 9: 1196-1204.

21. Shimoda LA, Sham JS, Shimoda TH, Sylvester JT (2000) L-type Ca(2+) channels, resting [Ca(2+)]_i, and ET-1-induced responses in chronically hypoxic pulmonary myocytes. *Am J Physiol Lung Cell Mol Physiol* 279: L884-894.

22. Duong-Quy S, Dao P, Hua-Huy T, Guilluy C, Pacaud P, et al. (2011) Increased Rho-kinase expression and activity and human pulmonary endothelial dysfunction in smokers with normal lung function. *Eur Respir J* 37: 349-355.

23. Bei Y, Duong-Quy S, Hua-Huy T, Dao P, Le-Dong NN, et al. (2013) Activation of RhoA/Rho-kinase pathway accounts for pulmonary endothelial dysfunction in patients with chronic obstructive pulmonary disease. *Physiol Rep* 1: e00105.

24. Duong-Quy S, Bei Y, Liu Z, Dinh-Xuan AT (2013) Role of Rho-kinase and its inhibitors in pulmonary hypertension. *Pharmacol Ther* 137: 352-364.