

Improvement Bryc's Approximation to the Cumulative Distribution Function

Abderrahmane Malki*

Algeria Mathematical Department, EPST School of Algiers, Algeria

Abstract

We develop two news approximations to the cumulative distribution function. We begin by improving the accuracy of Cadwell's approximation. We reduce the accuracy from 0.006466 to 1.6635e-004. For the second approximation we reduce the accuracy of Bryc's approximation from 7.062e-004 to 2.072e-005. As a performance, we use a Maximum Absolute Error (M.A.E.). We recommend these two new approximations for their high accuracy.

Keywords: Cumulative distribution function; Maximum absolute error; Cadwell's approximation; Bryc's approximation

Introduction

Several domains of engineering, statistics or applied mathematics need the Cumulative Distribution Function (C.D.F). In statistics, the Probability Density Function (P.D.F) defined by:

$$\varphi(t) = \frac{e^{-\frac{t^2}{2}}}{\sqrt{2\pi}}; -\infty < t < +\infty \quad (1)$$

The Cumulative Distribution Function (C.D.F) given by:

$$\Phi(Z) = \int_{-\infty}^Z \varphi(t) dt \quad (2)$$

This C.D.F. does not have a closed form. Most books in probability and statistics insert tables of C.D.F. For each value of the variable Z non-in this tables, we need a computer for estimate $\Phi(Z)$ by elementary methods [1]. For this reason, in the literature we find many approximations to the C.D.F. with closed forms. In section two, we present Cadwell's approximation with his original form and we improve the accuracy of this approximation with four decimal places. In section three, we introduce an approximation with four decimals places and we improve his accuracy until five decimals places. In section four, we conclude our paper.

Improving Cadwell's Formula

In 1951, Cadwell presents his new formula for approximate the cumulative normal distribution [2]. This formula given by:

$$\Phi_{cadwell}(z) \approx \frac{1}{2} \left\{ 1 + \sqrt{1 - e^{-z^2 \left(\frac{2}{\pi} - \frac{3z^2}{\pi^2} \right)}} \right\} \quad (3)$$

The M.A.E. is:

$$E_{cadwell} = \max_{-6 \leq z \leq 6} |\Phi_{cadwell}(z) - \Phi(z)| \cong 0.00646 \quad (4)$$

Our new formula defined by:

$$\Phi_{Malki}(z) \approx \frac{1}{2} \left\{ 1 + \sqrt{1 - e^{-z^2 (0.6349114 - 0.0073962z^2)}} \right\} \quad (5)$$

$$E_{Malki} = \max_{-6 \leq z \leq 6} |\Phi_{Malki}(z) - \Phi(z)| = 1.663503182035564e-004$$

$$\text{We have then } E_{cadwell} \cong 38.87 \times E_{Malki} \quad (6) \text{ (Figure 1).}$$

The M.A.E. for Cadwell is about 40 times that Malki.

Improving Bryc's Formula

In, 2002, Bryc presents in his paper the following formula [3].

$$\Phi_{Bryc}(z) = 1 - \frac{z + 3.333}{2 \times 3.333 + 7.32z + \sqrt{2\pi z^2}} e^{-\frac{z^2}{2}} \quad (7)$$




Figure 1: Shows the curves of absolute error for Cadwell and our new formula Malki.

The maximum absolute error is:

$$E_0 = \max_z |\Phi_{Bryc}(z) - \Phi(z)| = 7.062660715300151e-004 \quad (8)$$

We can write formula (8) as

$$\Phi_{Bryc}(z) = 1 - \frac{\frac{z}{\sqrt{2\pi}} + \frac{3.333}{\sqrt{2\pi}}}{2 \times \frac{3.333}{\sqrt{2\pi}} + \frac{7.32}{\sqrt{2\pi}}z + z^2} e^{-\frac{z^2}{2}}$$

This formula has the form

$$\Phi_{Bryc}(z) = 1 - \frac{a z + b}{2 \times b + c z + z^2} e^{-\frac{z^2}{2}} \quad (9)$$

We improve the accuracy of this formula by our new formula

$$\Phi_{Malki2017}(z) = 1 - \frac{0.3656123z + 1.4578862}{2 \times 1.4578862 + 3.054682z + z^2} e^{-\frac{z^2}{2}} \quad (10)$$

*Corresponding author: Malki A, Algeria Mathematical Department, EPST School of Algiers, Algeria, Tel: +21321621841; E-mail: malkin010455@yahoo.fr

Received February 27, 2017; Accepted April 11, 2017; Published April 18, 2017

Citation: Malki A (2017) Improvement Bryc's Approximation to the Cumulative Distribution Function. Int J Swarm Intel Evol Comput 6: 151. doi: 10.4172/2090-4908.1000151

Copyright: © 2017 Malki A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 2: Shows the comparison of maximum absolute error for Bryc2001 and our new formula.

The maximum absolute error is:

$$E_1 = \max_z |\phi_{Malki2017}(z) - \Phi(z)| = 2.072294382748918e - 005 \quad (11)$$

Hence, we have a ratio

$$E_0 \cong 34.1 \times E_1 \quad (12) \text{ (Figure 2)}$$

Conclusion

This paper presents two approximations to the cumulative normal distribution and their improving approximations. The first approximation is two decimals places; we improve it from two decimals places to four decimals places. In the second approximation, we improve the Bryc's formula by our second new formula that reduce the accuracy is about 34 times. We recommend these two new approximations.

References

1. Zogheib B, Hlynka M (2009) Approximations of the standard normal distribution.
2. Cadwell JH (1951) The bivariate normal integral. Biometrika 38: 475-479.
3. Bryc W (2002) A uniform approximation to the right normal tail integral. Appl Math Comput 127: 365-374.

Citation: Malki A (2017) Improvement Bryc's Approximation to the Cumulative Distribution Function. Int J Swarm Intel Evol Comput 6: 151. doi: 10.4172/2090-4908.1000151

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>