

## In Vitro Glycation of the Pathogenic Prion Protein

Aly Moussa\*

National Agency for Sanitary Environmental, Laboratoire de Lyon, France

### National Agency for Environmental and Sanitary Security - Lyon Laboratory - France

The prion protein (PrP) possesses two glycosylation sites at the position 181 and 197. After electrophoresis the PrP is detected as a 3 isoforms; bi-glycosylated, mono-glycosylated and non-glycosylated. The interaction of ion, chemical or protein ligands with the prion protein leads to an increase of the molecular weight of the 3 isoforms either as aggregation or oligomerization [1].

Neurodegenerative disorders of the central nervous system are slowly developing, insidious conditions that contribute first to neuronal cell degeneration and later cell death. During the development of the spongiform encephalopathies in human and animals the surface cellular glycoprotein (PrP<sup>c</sup>) undergoes a misfolding conformation, become resistant to proteinase K and in detergents it accumulates as insoluble  $\beta$ -sheet containing aggregates [2,3].

Also was observed during the evolution of Alzheimer's disease (AD), Parkinson disease (PD) and Creutzfeldt-Jacob disease (CJD) an elevated levels of non-enzymatic protein glycation and formation of crosslinks via stable advanced glycation end products (AGEs). In addition, the AGEs also contributes to the generation of elevated release of intracellular reactive oxygen species (ROS) and nitrogen oxygen species, including the radical's superoxide and peroxy nitrite leading to development of oxidative stress, inflammation, and increased glycation of key intracellular as well extracellular proteins leading to the development of neurodegenerative disorders and progressively ending by death [4].

Here was investigated the effect of adding individual sugars to proteinase K treated pathogenic prion strains (PrPres) and heating at 100°C for variable time.



*In vitro* and at high temperature; after adding reducing sugars to PrPres it became attached to the amine group of the amino acid Lysine copies present on the prion protein through a non-enzymatic Maillard reaction. The quantity of the attached saccharide molecules will physically increase the molecular weight of the PrP peptide bands. Also the protein configuration can be changed due to the presence of sugar molecules leading to better exposure of certain epitopes and resulting in a better immune detection by their specific antibodies.

The results in Figure 1 showed the effect of adding 8  $\mu$ l of 1M lactose to 4  $\mu$ l of dilutions (P, 1:2, 1:4 and 1:8) of a proteinase K treated bovine brain suspension (PrPres), then adding 4  $\mu$ l Laemmli buffer to each of the 15 tubes, vortexing and heating during variable periods (0, 5, 10 and 20 min respectively). The tubes were then centrifuged for two min. at 10000 RPM. The supernatants were collected and a constant volume from each was deposited and run on 15% SDS acrylamide gel, transfer on nitrocellulose membrane and immunoblotted using an anti-PrP monoclonal antibody and peroxidase conjugated anti-mouse polyclonal serum [5].

The results in Figure 2 were obtained after adding to 4  $\mu$ l of proteinase K treated PrP<sup>c</sup> of previously confirmed cases from either ovine, bovine or mice adapted ovine strains 8  $\mu$ l of a 1 M solution of either galactose (Ga), glucose (G), lactose (La) or mannose (Ma) and finally to each tube 4  $\mu$ l of Laemmli buffer were added. The control

tubes (C) without any sugar were prepared by adding 4  $\mu$ l from the same prion strains to 12  $\mu$ l of Laemmli buffer. The mixtures were vortexed, heated for 15 min. at 100°C and centrifuged for two min. at 10000 RPM. The supernatants were collected and a constant volume from each was deposited and run on 15% SDS acrylamide gel, transfer on nitrocellulose membrane and immunoblotted using an anti-PrP monoclonal antibody and peroxidase conjugated anti-mouse polyclonal serum [5].

In Figure 3 was tested the capacity of 5 different prion strains to interact with mannose at 100°C during variable duration. To three sets




\*Corresponding author: Aly Moussa, National Agency for Sanitary Environmental, Laboratoire de Lyon, France, Tel: 336-174-609-52; E-mail: [moussa69@club-internet.fr](mailto:moussa69@club-internet.fr)

Received June 06, 2016; Accepted June 06, 2016; Published June 13, 2016

Citation: Moussa A (2016) *In Vitro* Glycation of the Pathogenic Prion Protein. J Chromatogr Sep Tech 7: e134. doi:10.4172/2157-7064.1000e134

Copyright: © 2016 Moussa A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



of 5 tubes each was added 8  $\mu$ l mannose and 4  $\mu$ l Laemmli buffer in each of the 15 tubes. to the tubes 1 of three set was added 4  $\mu$ l of an ovine field isolated strain, to the tubes 2 of three set was added 4  $\mu$ l of

another ovine field isolate, to the tubes 3 of three set was added 4  $\mu$ l of a bovine field prion strain, to the tubes 4 of three set was added 4  $\mu$ l of another bovine field isolate and to the tubes 5 of three set was added 4  $\mu$ l of a mouse brain suspension of a mouse adapted ovine strain. After vortexing the 5 tubes of the first set were not heated, those of the second set were heated for 10 min. at 100°C and those of the third set were heated for 20 min. at 100°C. The tubes were then centrifuged for two min. at 10000 RPM. The supernatants were collected and a constant volume from each was deposited and run on 15% SDS acrylamide gel, transfer on nitrocellulose membrane and immune blotted using an anti-PrP monoclonal antibody and peroxidase conjugated anti-mouse polyclonal serum.

### References

1. Moussa A (2016) Electrophoresis and western blot can detect the interaction of ion ligand with the pathogenic prion protein. J Chromatogr Sep Tech 7: e133.
2. Prusiner SB (1982) Novel Proteinaceous Infectious Particles Cause Scrapie. Science 216: 136-144.
3. Bolton DC, McKinley MP, Prusiner SB (1982) Identification of a protein that purifies with the scrapie prion. Science 218: 1309-1311.
4. Guix FX, Raga G, Bravo R, Nakaya T, De Fabritiis G, et al. (2009) Amyloid-dependent triose phosphate isomerase nitrotyrosination induces glycation and tau fibrillation. Brain 132: 1335-1345.
5. Moussa A, Coleman AW, Bencsik A, Leclere E, Perret F, et al. (2006) Use of streptomycin for precipitation and detection of proteinase K resistant prion protein (PrP<sup>Sc</sup>) in biological samples. Chem Commun (Camb) 9: 973-975.

**Citation:** Moussa A (2016) *In Vitro* Glycation of the Pathogenic Prion Protein. J Chromatogr Sep Tech 7: e134. doi:10.4172/2157-7064.1000e134

### OMICS International: Publication Benefits & Features

#### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

#### Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submit>