

In vitro Metabolic Stability Study of New Cyclen Based Antimalarial Drug Leads Using RP-HPLC and LC-MS/MS

Apoorva V Rudraraju¹, Mohammad F Hossain¹, Anjali Shrestha¹, Prince NA Amoyaw¹, Babu L Tekwani² and Faruk Khan MO^{1*}

¹College of Pharmacy, Southwestern Oklahoma State University, 100 Campus Drive, Weatherford, Ok 73096, USA

²National Center for Natural Products Research, University of Mississippi, University, MS 38677, USA

Abstract

Metabolic stability of the new antimalarial drug leads is determined using Human Liver Microsome (HLM) and specific cytochrome P450 enzyme (CYP2C8) taking the clinically used antimalarial drug chloroquine as a positive control. Experiment is done using standard methods. All the assays were conducted in 0.5 M phosphate buffer at pH 7.4. In general the metabolic reaction was initiated by adding 1 mM NADPH and 0.5 mg of enzyme. Incubations were done with time frequency of 0 hr, 1 hr, and 2 hrs at 37°C and the reactions were terminated by adding acetonitrile in the equal amounts of the assay mixture taken. The samples were centrifuged for 15 minutes at 10,000×g at 4°C and an aliquot of the supernatant was subjected to analysis using HPLC as well as LC-MS to confirm the masses of the drug and/or metabolite(s), if any. While chloroquine was found to be metabolized in a predictable manner by both HLM and CYP2C8, the drug leads were metabolically stable at similar experimental conditions. This study demonstrated that the new drug leads are worth conducting further preclinical evaluations.

Keywords: Metabolic stability; Preclinical study; Drug lead analysis; Drug discovery

Introduction

With the arrival of new combinational drug synthesis, the need for an expeditious evaluation of drug safety has become a paramount topic in drug discovery. The duration and impact of drug discovery revolving around drug absorption, distribution, metabolism and excretion will be evaluated according to their determined metabolic stability. The major organ for drug metabolism is liver that contains major drug metabolizing enzyme called Cytochrome P450 (CYP) system [1]. Drug metabolism refers to the susceptibility of compounds to biotransformation that depends on the presence of groups in the molecule those are open to enzyme catalyzed transformation [2]. Drug metabolism can be divided into two phases, Phase I and phase II. Phase I involves oxidation, reduction and hydrolysis reactions, which are catalyzed by the CYP and flavin containing monooxygenases (FMOs) whereas Phase II involves conjugation reactions catalyzed by metabolic enzymes like UDP-glucuronyltransferases (UGTs) and sulfotransferases [3,4].

Recognizing the metabolites of drugs is of paramount importance in drug discovery and development. The identification of drug metabolites in the early stages of the drug discovery is important in the development processes. The analytical tools like Liquid Chromatography-Mass Spectrometry (LC-MS) and HPLC play prominent role in these processes. Through this process of identification, the pharmacokinetic profiles can be assessed that are highly significant in detecting safety and efficacy of the drug leads before they are progressed to the clinical trials.

The investigation for metabolites take an advantage of the fact that majority of drug metabolites can be classified as predictable as they are formed from common accepted biotransformation reactions. However, there are many other illustrations of primary metabolites that are formed from uncommon reactions and are, therefore, not easily predictable. Molecular masses of predicted metabolites (*m/z* values) can be estimated based on mass shifts from the parent drug. For example, Chloroquine is the major antimalarial drug that used in the treatment and prophylaxis of malaria, the protonated molecular mass of the metabolite of chloroquine, desethylchloroquine (DCQ), is 292 *m/z* to that of the parent drug 320 *m/z*. Evaluation of some expected metabolites can be achieved by the acquisition of the complete MS spectrum using various MS instruments and also by Extracted Ion

Chromatography (EIC) [4,5]. Using LC/MS, Clarke (2001) defined broadly the approaches for the identification of metabolites in biological matrices. After identifying the unknown metabolite ion peaks, further investigation to achieve detailed information on the pathways for the structure evaluation is done by multistage product ion scans [6,7].

The main purpose of the present study was to determine the metabolic stability of a series of newly discovered cyclen bisquinoline antimalarial drug leads [8] and related compounds (Figure 1) by RP-HPLC and LC-MS techniques. Compound B was shown to be highly effective antimalarial agent both *in vitro* and *in vivo* and was found to work by inhibiting β -hematin formation [8]. This lead structure fulfilled few important criteria for a new drug lead: 1) it is a 4-aminoquinoline derivative that are the most trusted class of antimalarials, 2) active against chloroquine-resistant as well as multidrug resistant isolates of *Plasmodium falciparum*, 3) simple pharmacophore structure that will afford low cost manufacturing, and 4) it also showed oral efficacy in mice model. All these features warranted further preclinical studies with these leads and related newly synthesized compounds as shown in Figure 1. Compounds A, C, D and E were also shown to have potent *in vitro* antimalarial activity and also serve as drug leads (manuscript is under preparation). The present study was thus aimed at determining the *in vitro* metabolic stability and identifying potential metabolites by HPLC and LC-MS techniques using HLM and CYP enzymes. The study was designed based on the known metabolic pathway of the related clinically used drug chloroquine (CQ) [9-12] and thus utilized both pooled human liver microsomes as well as specific isozyme CYP2C8 (Figure 1).

***Corresponding author:** Faruk Khan MO, College of Pharmacy, Southwestern Oklahoma State University, 100 Campus Drive, Weatherford, Ok 73096, USA, Tel: 580-774-3064; E-mail: faruk.khan@swosu.edu

Received March 16, 2013; **Accepted** July 14, 2014; **Published** July 17, 2014

Citation: Rudraraju AV, Hossain MF, Shrestha A, Amoyaw PNA, Tekwani BL, et al. (2014) *In vitro* Metabolic Stability Study of New Cyclen Based Antimalarial Drug Leads Using RP-HPLC and LC-MS/MS. Mod Chem appl 2: 129. doi:10.4172/2329-6798.1000129

Copyright: © 2014 Rudraraju AV, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: Chemical structures of chloroquine and the main drug leads A, B, C, D, and E.

Materials and Methods

Materials

Chloroquine diphosphate purchased from Pfaltz & Bauer. NADPH, anhydrous with assay 93-100% (HPLC), was purchased from Sigma Aldrich. 0.5 M Potassium phosphate buffer pH 7.4 was purchased from BD Gentest. Human Liver Microsome (HLM) was purchased from BD Bioscience and Corning. Human cDNA expressing CYP 450 and 2C8 were purchased from XenoTech and Corning. In the solvent system methanol Triethylamine, DMSO, and acetonitrile were purchased from Fisher Scientific. Dibasic anhydrous sodium phosphate and phosphoric acid were purchased from Fisher Scientific. Deionized water used was further purified by filtration and degassing. The drug leads mentioned in the Figure 1 were synthesized.

Chromatographic conditions

HPLC Method 1: The HPLC system used was Agilent 1100 series (Hewlett-Packard-Strasse 8, 76337 Waldbronn Germany), equipped with a pump, Diode array detector with UV lamp operated at variable wavelengths, autosampler, and thermostat. Data Acquisition was performed using Open Lab CDS Chem Station Edition software package with A.01.02 version implemented in the chromatographic system. Detection at 250 nm and 280 nm was standard. The stationary phase consisted of Waters X-Bridge C-18 column (4.6 mm×150 mm, 5.0 μm particle size, pore volume 0.76 cm³/g) purchased from Waters Corporation (34 Maple Street, Milford, Massachusetts 01757-3696 U.S.A). It was operated at constant temperature of 25°C on both ends of the column.

The mobile phase consisted of a mixture of 0.04% formic acid in water (i), acetonitrile (ii) and methanol (iii) in a gradient elution mode as shown in Table 1 [9]. The flow rate was maintained at 1.0 ml/min.

LC-MS system used was Shimadzu prominence LC 20AT, equipped with Degasser (DGu- 20A5), Auto sampler (SIL-20AHT), Refractive index detector (RID-10A), Diode array detector (SPDM-20M20A), Fraction collector (FRC-10A) and a mass spectrometer (LC-MS 2020). Data Acquisition was performed by Lab Solutions Real Time Analysis Software, implemented in the chromatographic system.

HPLC Method 2: Chromatographic separation of CQ, drug lead B, and metabolites was successfully achieved on a Waters X-Bridge C-18 column (4.6 mm×250 mm, 5.0 μm particle size, part no. 186003117) purchased from Waters Corporation in an isocratic separation mode with mobile phase consisting of 0.1% of triethylamine in methanol

and 0.02 M dibasic sodium phosphate (anhydrous) at pH 3.5 adjusted with Phosphoric Acid in the ratio of (60:40, v/v). The flow rate was maintained at 1.0 ml/min, the column oven temperature at 40°C and the effluent was monitored at 325 nm (based on the λ). The peak purity data were obtained using Photodiode Array (PDA) detector in the sample chromatograms. The method was found linear over the concentration range of 3.5–200.1 μg/ml (R²=0.99) for drug lead B.

General procedure

The method used to determine the metabolism was a modified method obtained from BD Bioscience, which is summarized here. Metabolic stability study of a drug can be determined by treating it with the liver enzyme that correlates with the *in vivo* conditions. The assay mixture is prepared by the combination of substrate (0.01 mM to 0.5 mM), enzyme (0.5 mg), buffer (0.5 M), and 1 mM NADPH cofactor. The order of addition of the assay component also plays a major role in the stability study. Metabolism can be initiated by pre-warming and incubating the substrate, buffer, and the cofactor to 37°C and then adding liver microsomes or CYP2C8 to the mixture. It is customary that the assay mixture should be thoroughly mixed. After the preparation of the Total Assay Mixture (TAM), required volume of the sample is collected at different time intervals. The reaction is then terminated using acetonitrile in equal volumes as that of the sample solution collected. The samples are then transferred into the microcentrifuge tubes, vortexed for 2 minutes and then centrifuged at 10,000×g for 15 minutes. The purpose of centrifugation is to remove the protein. The sample solution is then separated from the protein pellet by taken the supernatants. These supernatant solutions are transferred into the HPLC auto sampler vials and are analyzed according to the analytical method.

Metabolic stability in HLM using HPLC Method 1

The HLM used was pooled and prepared from freshly frozen human tissues which were tested negative for pathogens using PCR. The HLM used in this study was comprised of 330 pmoles/mg of total P450 and 420 pmoles/mg cyt. b₅. It is a mixture of CYP1A2, 2A6, 2B6, 2C8, 2C9, 2C19, 2D6, 2E1, 3A4, 4A11, FMO, UGT1A1, UGT1A4, and UGT1A9 where the enzyme activity of CYP2C8, 3A4, and 2D6 was 82, 3200, and 110 pmoles/(mg×min) respectively. The volume of the enzyme per vial was 0.5 ml and the protein content was 20 mg/ml in 250 mM sucrose.

All the incubations were performed in duplicates. Therefore, all together there were 6 samples for each drug, one blank and 6 samples of the positive control. The total experiment with 5 drugs has about 65 samples. All the components used in the TAM are shown in Table

Time (Min)/solvent	0.04% formic acid in Water	Acetonitrile	Methanol
0	96	0	4
10	64	30	6
12	4	90	6

Table 1: Mobile phase for metabolic stability study: Gradient elution mode.

S. No.	Ingredients	Volume (in 1 mL of total assay mixture)	Final Concentration
1	Sample	10 μ l	0.05 mM
2	Buffer (0.5 M)	200 μ l	100 mM
3	DI Water	598 μ l	-
4	NADPH	167 μ l	1 mM
5	Enzyme	25 μ l	0.5 mg

Table 2: Total assay mixture using HLM.

S. No.	Ingredients	Volume (TAM)	Final Concentration
1.	Sample	180 μ l	0.5 mM
2.	Buffer	1275 μ l	0.2 M
3.	NADPH	300 μ l	1m M
4.	Enzyme	45 μ l	25 pmoles/ml

Table 3: Total assay mixture using CYP2C8.

S. No.	Ingredients	Sample		Standard	Blank
		Volume, 2 mL of TAM	Final Concentration	Volume, 2 mL of TAM	Volume, 2 mL of TAM
1	Substrate (1 mM)	20 μ l	0.01 mM	20 μ l	---
2	Buffer (0.5 M)	400 μ l	100 mM	400 μ l	400 μ l
3	NADPH	200 μ l	1 mM	---	200 μ l
4	DI Water	1355 μ l	---	1580 μ l	1375 μ l
5	Enzyme	25 μ l	0.5 mg	---	25 μ l

Table 4: Total assay mixture using HLM and CYP2C8.

2 with their volumes. All the ingredients in the TAM are taken in the order of their serial number (Table 2).

5 mM of the substrate stock solutions were prepared by dissolving in DMSO. The buffer used was 0.5 M potassium phosphate pH 7.4 (BD Biosciences Cat No. 451201). TAM was prepared by adding 10 μ l of the sample, 200 μ l of the buffer solution, 598 μ l of water, and 167 μ l of NADPH. It was then pre-incubated for 2 minutes in a shaking water bath and 25 μ l of the enzyme was added to initiate the reaction. The capped tube was mixed thoroughly by inverting a couple of times. Immediately, at time t=0 hr, 300 μ l of the TAM was taken into a centrifuge tube and the reaction was terminated by placing it on ice bath and adding 300 μ l of acetonitrile. The remaining TAM was warmed in a shaking water bath at 37°C for 2 hrs and the same procedure is followed for the samples at time t=1 hr and t=2 hr. The incubation mixtures were then centrifuged for 15 minutes at 10,000 \times g at 4°C. About 500 μ l aliquots of the supernatant were collected and were subjected to analysis using the HPLC conditions as shown in Table 1. After running HPLC the samples were frozen and then lyophilized to remove the solvent. The dried samples were then dissolved in methanol and water (50:50 v/v) for LC-MS analyses.

Metabolic stability in CYP2C8 using HPLC method 1

The CYP enzyme is comprised of a human CYP2C8 and human CYP-reductase co-expressed in *Escherichia coli*. The concentration of the CYP P450 is 1.0 nmol/ml and its protein concentration is 10.0 mg/ml. All the assays were conducted in 0.2 M phosphate buffer at pH 7.4. One blank for each sample A, B, C, D, and E was prepared by adding 180 μ l of the sample in 1620 μ l of the buffer. The total assay mixture contained 25 pmoles/ml of CYP2C8 and 0.5 mM of sample as shown in the table 3.

180 μ l of Sample, 1275 μ l buffer, and 300 μ l of 1 mM NADPH was taken into a reaction flask and was pre-incubated for 2 minutes. The reaction was initiated by adding 45 μ l enzyme. Incubations were done with increasing time (t=0 hr, 1 hr, 2 hrs) at 37°C. Immediately at time=0 hr, 300 μ l of the assay mixture was pipetted into a centrifuge tube and 300 μ l of acetonitrile was added to it to stop the reaction. All the samples were done in triplicates. Therefore, all together there were 9 samples for each drug, one blank and 9 samples of the positive control. The total experiment with 5 drugs had about 95 samples.

The remaining TAM was incubated for 1 hr and 2 hrs, same procedure of sampling was followed as mentioned in the HLM section. Then the samples were centrifuged for 15 minutes at 10,000 \times g at 4°C. An aliquot about 500 μ l of the supernatant fraction were collected and subjected to analysis using the above HPLC conditions as shown in Table 1. The retention times were noted and the amounts of drugs metabolized were calculated according to their percentage peak area using the equation: % peak area of the metabolite/(% peak area of the drug+% peak area of the metabolite). After running HPLC the samples were frozen and then lyophilized to remove the solvent. The dried samples were then dissolved in methanol and water (50:50 v/v) and tested for their mass. The mass of the drug and the metabolite were determined by using LC-MS.

Metabolic stability in HLM and CYP2C8 using HPLC method 2

To verify the metabolic stability results using HPLC method 1, HPLC method 2 was carried out using drug lead B, and CQ as a positive control. For that we prepared sample, standard and blank for our drug lead B and CQ in duplicates as mentioned in the following Table 4. All the ingredients in the TAM were taken in the order of their serial number (Table 4).

Sample stock solutions of Chloroquine diphosphate and Cylen Bisquinoline Hydrochloride Salt were prepared by dissolving in water. 0.5 M potassium phosphate pH 7.4 (BD Biosciences Cat No. 451201) was used as a buffer. Sample solutions were prepared by adding 20 μ l of substrate, 400 μ l of the buffer solution, 200 μ l of the NADPH solution and 1355 μ l of water. Standard solutions were prepared by adding 20 μ l of Substrate, 400 μ l of the buffer solution and 1580 μ l of water. Blank solutions were prepared by adding 400 μ l of the buffer solution, 200 μ l of the NADPH solution and 1375 μ l of water.

All the above solutions were then pre-incubated for 5 minutes in a shaking water bath at 37°C. Reaction was initiated by adding 25 μ l of the enzyme (from Corning HLM-452161 and CYP2C8-456252). These TAMs were mixed thoroughly by inverting a couple of times and returned to the water bath. Immediately, at time t=0 hr, 500 μ l of the TAM was taken into a centrifuge tube and the reaction was terminated by placing it on ice bath and adding 500 μ l of acetonitrile. The remaining TAMs were warmed in a shaking water bath at 37°C for 2 hrs and the same procedure was followed for the samples at time t=1 hr and t=2 hr. The incubation mixtures were then centrifuged for 15 minutes at 10,000 \times g at 4°C. About 500 μ l aliquots of the supernatant were collected and subjected to analysis using the HPLC method 2. After running HPLC the samples were frozen and then lyophilized to remove the solvent. The dried samples were then dissolved in methanol and water (50:50 v/v) for LC-MS analyses.

Results and Discussion

HPLC Method 1

Results using HLM: Under the provided experimental conditions




Figure 2: HPLC chromatogram of chloroquine at time A) 0 hour and B) 2 hour using HLM.




Figure 3: HPLC chromatogram of drug lead B at time A) 0 hour and B) 2 hour using HLM.

Drugs	Time t=1 hr			Time t=2 hr		
	RT	Metabolism	MS- (m/z)	RT	Metabolism	MS- (m/z)
A	8.136	NO	523	8.141	NO	523
B	5.737	NO	495	5.791	NO	495
C	4.869	NO	561	4.877	NO	561
D	5.818	NO	885	5.822	NO	885
E	5.822	NO	619	5.831	NO	619
CQ/DCQ	6.801/6.485	YES	320, 292	6.817/6.510	YES	320, 292

*Percentage of the metabolite formation was calculated according to the obtained percentage peak areas of the metabolite and the drug. The percent of CQ metabolized using HLM at time t=1 hr and 2 hrs was 5.12% and 6.146%, respectively.

Table 5: Data obtained from HPLC and MS for the drugs using HLM.

it was observed that CQ was metabolized in a predictable manner by HLM into the metabolite DCQ; contrarily all our experimental drug leads were metabolically stable. Not only was the metabolite for CQ identified in HPLC (Figure 2) but also in MS (Figure 7) reconfirming our results. Both drug (320 m/z) and metabolite masses (292 m/z) were observed in the LC-MS analysis of the samples. According to the area under curve ratios, CQ was converted to about 5.12% of its N-deethylated metabolite after 1 hour of incubation and 6.42% after 2 hours of incubation. On the other hand, the drug leads A (523m/z), B (495m/z), C (561m/z), D (885 m/z), and E (619 m/z) were found to be metabolically stable in both HPLC and LC-MS analyses. A sample chromatogram of CQ and compound B are shown in the Figures 2 and 3. The retention times and the peak areas of the drug leads A, B, C, D, E and CQ obtained at time t=1 hr and 2 hr are shown in the Table 5. At time t=0 min there was no metabolism with any of the drug samples. At time t=1 hr CQ showed some metabolism whereas the drug leads were stable. There was no evidence of any metabolite mass of any of the drug leads when they were analyzed by LC-MS. The molecular ion peaks (m/z) were similar to those of their molecular weights (Figures 2 and 3) (Table 5).

Results using CYP2C8: As shown in Figure 4, about 10.33% of CQ was transformed into DCQ after 1 hour of incubation and 12.26% after

2 hours of incubation, whereas the drug lead B was stable as shown in Figure 5. The drug leads A (523 m/z), B (495 m/z), C (561 m/z), D (885 m/z), and E (619 m/z) were metabolically stable at similar experimental conditions. The retention times of the drugs A, B, C, D, E and CQ obtained at time t=0 were 8.136, 5.971, 4.869, 5822, 5.817 and 6.998, respectively, and no metabolite peak was evident. The retention times of the drug leads and the CQ at time t=1 hr and t=2 hr is shown in the Table 6. All the samples were estimated in triplicates and the results shown are the average of the three trials (Figures 4 and 5) (Table 6).

With these results, it was confirmed that CYP2C8 has an important role in the metabolism of chloroquine and the drug leads were metabolically stable. The experiment was extended up to 24 hours for the drug leads and consequently no metabolism was observed. The drug masses were confirmed by MS (Figures 6-9).

HPLC Method 2

Results using HLM: Under provided experimental conditions using HPLC method 2, similar results were obtained, CQ was metabolized by HLM into the metabolite DCQ and contrarily the drug lead B was metabolically stable. The metabolite was identified by HPLC and also by LC-MS reconfirming the results shown in method 1. Both drug (320 m/z) and metabolite (292 m/z) masses were observed in the LC-MS analysis of the samples. According to the area under curve ratios, CQ was converted to about 4.29% of its N-deethylated metabolite after 1




Figure 4: HPLC chromatogram of chloroquine at time A) 0 hour and B) 2 hour using CYP2C8.




Figure 5: HPLC chromatogram of drug B at time A) 0 min and B) 2 hour using CYP2C8.

Drugs	Time t= 1 hr			Time t= 2 hr		
	RT	Metabolism	MS- (m/z)	RT	Metabolism	MS-(m/z)
A	8.126	NO	523	8.136	NO	523
B	5.973	NO	495	5.944	NO	495
C	4.875	NO	561	4.869	NO	561
D	5.819	NO	885	5.822	NO	885
E	5.821	NO	619	5.817	NO	619
CQ/DCQ	7.017/6.784	YES	320, 292	7.011/6.772	YES	320, 292

*The percent of CQ metabolized using CYP2C8 at time t=1hr and 2hrs was 12.26% and 10.33%, respectively.

Table 6: Data obtained from HPLC and MS for the drugs using CYP2C8.




Figure 6: MS spectrum showing chloroquine standard.




Figure 7: MS spectrum showing chloroquine and N-desethylchloroquine masses at t=2hr using HLM. Similar chromatogram was observed when the experiment was conducted using CYP2C8.




Figure 8: MS spectrum showing drug lead B standard.

hour of incubation and 6.61% after 2 hours of incubation. On the other hand, the drug leads B (495 m/z) was found to be metabolically stable in both HPLC and LC-MS analyses. A sample chromatogram of CQ and compound B are shown in Figure 10C and Figure 11C, respectively. The retention times and the peak areas of the drug lead B and CQ obtained at time t=1hr and 2hr are shown in the Table 7.

Results using CYP2C8: Under provided experimental conditions using HPLC method 2, similar results were obtained, CQ was metabolized by CYP2C8 into the metabolite DCQ and contrarily the drug lead B was metabolically stable. Using CYP2C8 about 2.23% of CQ was transformed into DCQ after 1 hour of incubation and 2.89% after 2 hours of incubation. The drug lead B (495 m/z) was metabolically stable at similar experimental conditions (Figure 11D). The retention times of the drug lead B and the CQ at time t=1 hr and t=2 hr are shown in the Table 8. All the samples were estimated in duplicates and the results shown are the average of the two trials (Figures 10 and 11) (Table 8).

According to the results obtained from HPLC method I and II, it was confirmed that both HLM and CYP2C8 has an important role in the metabolism of chloroquine whereas, all drug leads were metabolically stable to these enzyme.

Discussion

The *in vitro* metabolism of the new drug leads compared to the positive control chloroquine in the presence of HLM and cDNA expressing CYP2C8 have been demonstrated in this study. In the pharmacokinetic studies of CQ, Frisk-Holmberg [6,7] reported the single dose kinetics of CQ and its major metabolite DCQ in healthy subjects and identified that CQ is dealkylated into two main metabolites, N-desethylchloroquine (DCQ) and N-bis-desethylchloroquine (BDCQ). Identification of metabolites is done by LC-MS. Mass spectrometers help in separating the compounds from each other using the difference in mass-to-charge ratio (m/z) of ionized compounds.

In blood and plasma, the concentration of DCQ was detected up to 20 to 50% of those of the parent compound CQ, whereas the conversion to BDCQ was <10% [6,7]. It is known that CYPs are major drug metabolizing enzyme system. Ofori- Adjei and Ericsson [13] estimated that 30 to 50% of an administered dose of chloroquine is metabolized by the liver via cytochrome P450 enzymes. Identification of the major enzymes, associated with drug's metabolism therefore started by evaluating and identifying the role of CYP isoforms. Xue-Qing Li et al. [14] identified human CYPs that metabolize anti-parasitic drugs using three approaches, (a) Relative Activity Factor (RAF), (b) correlation analysis of their activity, and (c) inhibitors diagnostics. They reported that the major CYPs involved in the metabolism are CYP isoforms 2C8, 3A4, and 2D6. The RAF approach is used significantly to bridge the gap between liver microsomes and recombinant systems. It also estimates the individual P450 contributions to drug metabolism [15,16]. Using the same approach, Projean et al. [9] also reported that CYPs 2C8, 3A4, and 2D6 are the main isoforms catalyzing DCQ formation. This was the first investigation of CQ metabolism in HLM and they identified that CYP2C8 and CYP3A4 contributes collectively




Figure 9: MS spectrum showing drug lead B mass at t=2 hr using HLM. Similar chromatogram was observed when the experiment was conducted using CYP2C8.

Drugs	Time t=1 hr			Time t=2 hr		
	RT	Metabolism	MS- (m/z)	RT	Metabolism	MS- (m/z)
B	13.135	NO	495	13.10	NO	495
CQ/DCQ	5.460/4.480	YES	320, 292'	5.470/4.508	YES	320, 292'

*The percent of CQ metabolized using HLM at time t=1 hr and 2 hrs was 4.29% and 6.61%, respectively

Table 7: Data obtained from HPLC and MS for the drug using HLM.

Drugs	Time t=1 hr			Time t=2 hr		
	RT	Metabolism	MS- (m/z)	RT	Metabolism	MS- (m/z)
B	13.145	NO	495	13.129	NO	495
CQ/DCQ	5.486/4.533	YES	320,292'	5.432/4.517	YES	320, 292'

*The percent of CQ metabolized using CYP2C8 at time t=1 hr and 2 hrs was 12.26% and 10.33%, respectively

Table 8: Data obtained from HPLC and MS for the drugs using CYP2C8.




Figure 10: HPLC chromatograms of chloroquine. A) Blank (NADPH, Enzyme & Buffer), B) CQ (Control without Enzyme), C) CQ in HLM (t=2 hr), and D) CQ in CYP2C8 (t=2 hr).

more than 80% of the total CQ N-desethylation over a wide range of concentrations. Their study suggested that, in humans, the therapeutic concentration of CQ that is metabolized into DCQ is primarily via CYP2C8 and CYP3A4, whereas 2D6 plays a significant role when CQ concentrations are low. This study also shows that CYPs, 2C8, and 3A4 constitute low-affinity and high-capacity systems, whereas CYP2D6 has higher affinity but a significantly lower capacity. Among these CYP2C8 represents approximately 5% of the total hepatic CYPs and metabolizes drugs that are amides or weak bases with two hydrogen bond acceptors [11,17,18]. Kim et al. [10] also identified that CYP2C8 and 3A4/5 are involved in metabolism of CQ into desethylchloroquine in human liver microsomes. This was also consistent with the previous data obtained from Ducharme and Farinotti [19]. Previous studies show that the relative content of CYP 3A4, 2C8 and 2D6 in 60 human samples are 29%, 18%, and 2%, respectively [20]. Based on these literature evidence, HLM and CYP2C8 were obvious selection to study the *in vitro*

metabolism stability of the new drug leads due to the similarity of the structures and activity of the lead compounds with chloroquine. While chloroquine was metabolized by both HLM and CYP2C8 considerably, all these new drug leads were metabolically stable.

In the Figure 10, the HPLC chromatograms clearly showed that CQ was metabolized and there was an additional peak before the parent peak which was absent in blank and control sample, partially confirms the metabolism of chloroquine. Moreover, when the samples were tested with LC-MS (Figure 7), positive control chloroquine sample at time t=2 hrs gave two ion peaks, one at 320 m/z and another at 292 m/z, which confirms the metabolism of the drug.

In the Figure 11, the HPLC chromatograms clearly showed that drug lead B was not metabolized but the appearance of an additional peak before the parent peak in blank and control sample confirms the presence of impurity in drug lead B. The spectrum (Figure 9) obtained




Figure 11: HPLC chromatograms of drug lead B. A) Blank (NADPH, Enzyme & Buffer), B) Drug lead B (Control without Enzyme), C) Drug lead B in HLM (t=2 hr), and D) Drug lead B in CYP2C8 (t=2 hr).

from LC-MS showed the ion peak of the drug lead B at time t=2 hrs as 495 m/z, which confirms its molecular weight. Henceforth there is no metabolism of the drug lead B.

As discussed earlier, CYP enzymes provide information regarding the metabolites. Nebert and Russell [21] mentioned that there are 270 different CYP gene families, and 18 of those gene families were recorded in mammals that encrypt 57 CYP genes in humans. Out of all CYP enzymes, CYP2C8, 3A4, and 2D6 are involved in Chloroquine metabolism. This was proven in our experiment where CYP2C8 contributed to more than 10% of the total CQ *N*-desethylation over a 100 μ M concentration [22]. It has low affinity and high capacity in metabolizing the chloroquine. Further investigations would be conducted utilizing more CYP isozymes at different concentration ranges and time intervals to identify any possible metabolic pathways for these drug leads in near future as a part of our continuous preclinical analysis. This study demonstrated a convenient method development and also a metabolic stability of the drug leads against most important CYP isozyme that is involved in the metabolism of standard drug chloroquine.

Acknowledgment

This project was supported by the National Institute of General Medical Sciences of the National Institutes of Health through Grant Number 8P20GM103447.

References

- Casarett LJ, Doull J, Klaassen CD (2001) Parkinson A: Biotransformation of Xenobiotics. Casarett and Doull's Toxicology: The Basic Science of Poisons, McGraw-Hill Medical Publishing Division, New York, USA.
- Ariens EJ, Simonis AM (1982) Optimization of pharmacokinetics-an essential aspect of drug development by metabolic stabilization. Strategy in drug research. Keverling Buisman JA, editor. Elsevier Scientific Publishing Company, Amsterdam.
- Smith DA, Jones BC, Walker DK (1996) Design of drugs involving the concepts and theories of drug metabolism and pharmacokinetics. Med Res Rev 16: 243-266.
- Baranczewski P, Stańczak A, Sundberg K, Svensson R, Wallin A, et al. (2006) Introduction to *in vitro* estimation of metabolic stability and drug interactions of new chemical entities in drug discovery and development. Pharmacol Rep 58: 453-472.
- Tiller PR, Romanynshyn LA (2002) Liquid chromatography/tandem mass spectrometric quantification with metabolite screening as a strategy to enhance the early drug discovery process. Rapid Commun Mass Spectrom 16: 1225-1231.
- Frisk-Holmberg M, Bergqvist Y, Domeij-Nyberg B (1983) Steady state disposition of chloroquine in patients with rheumatoid disease. Eur J Clin Pharmacol 24: 837-839.
- Gustafsson LL, Walker O, Alván G, Beermann B, Estevez F, et al. (1983) Disposition of chloroquine in man after single intravenous and oral doses. Br J Clin Pharmacol 15: 471-479.

8. Khan MO, Levi MS, Tekwani BL, Khan SI, Kimura E, et al. (2009) Synthesis and antimalarial activities of cyclen 4-aminoquinoline analogs. *Antimicrob Agents Chemother* 53: 1320-1324.
9. Projean D, Baune B, Farinotti R, Flinois JP, Beaune P, et al. (2003) In vitro metabolism of chloroquine: identification of CYP2C8, CYP3A4, and CYP2D6 as the main isoforms catalyzing N-desethylchloroquine formation. *Drug Metab Dispos* 31: 748-754.
10. Kim KA, Park JY, Lee JS, Lim S (2003) Cytochrome P450 2C8 and CYP3A4/5 are involved in chloroquine metabolism in human liver microsomes. *Arch Pharm Res* 26: 631-637.
11. Lewis DF (2004) 57 varieties: the human cytochromes P450. *Pharmacogenomics* 5: 305-318.
12. Ducharme J, Farinotti R (1996) Clinical pharmacokinetics and metabolism of chloroquine. Focus on recent advancements. *Clin Pharmacokinet* 31: 257-274.
13. McChesney EW, Banks WF Jr, Sullivan DJ (1965) Metabolism of chloroquine and hydroxychloroquine in albino and pigmented rats. *Toxicol Appl Pharmacol* 7: 627-636.
14. Ofori-Adjei D, Ericsson O (1985) Chloroquine in nail clippings. *Lancet* 2: 331.
15. Li XQ, Björkman A, Andersson TB, Gustafsson LL, Masimirembwa CM (2003) Identification of human cytochrome P(450)s that metabolise anti-parasitic drugs and predictions of in vivo drug hepatic clearance from in vitro data. *Eur J Clin Pharmacol* 59: 429-442.
16. Crespi CL (1995) Xenobiotic-metabolizing human cells as tools for pharmacological and toxicological research. *Advances in Drug Research* 26: 179-235.
17. Störmer E, von Moltke LL, Shader RI, Greenblatt DJ (2000) Metabolism of the antidepressant mirtazapine in vitro: contribution of cytochromes P-450 1A2, 2D6, and 3A4. *Drug Metab Dispos* 28: 1168-1175.
18. Pelkonen O, Turpeinen M, Hakkola J, Honkakoski P, Hukkanen J, et al. (2008) Inhibition and induction of human cytochrome P450 enzymes: current status. *Arch Toxicol* 82: 667-715.
19. Musana AK, Wilke RA (2005) Gene-based drug prescribing: clinical implications of the cytochrome P450 genes. *WMJ* 104: 61-66.
20. Ducharme J, Farinotti R (1997) Rapid and simple method to determine chloroquine and its desethylated metabolites in human microsomes by high-performance liquid chromatography with fluorescence detection. *J Chromatogr B Biomed Sci Appl* 698: 243-250.
21. Shimada T, Yamazaki H, Mimura M, Inui Y, Guengerich FP (1994) Interindividual variations in human liver cytochrome P-450 enzymes involved in the oxidation of drugs, carcinogens and toxic chemicals: studies with liver microsomes of 30 Japanese and 30 Caucasians. *J Pharmacol Exp Ther* 270: 414-423.
22. Nebert DW, Russell DW (2002) Clinical importance of the cytochromes P450. *Lancet* 360: 1155-1162.