

Increased Serum Levels of TNF- α and IL-6 are not Related to HLA-Cw6 in Psoriasis Patients Correlation of Cytokine with HLA Cw6

Sangeeta Singh^{1*}, Gyanendra Kumar Sonkar², Shinjini Singh and Usha Singh³

¹Department of Biochemistry, King George's Medical University, Lucknow, India

²King George's Medical University, Lucknow, India

³Department of Pathology, Institute of Medical Sciences, Banaras Hindu University, Varanasi, India

Abstract

Psoriasis is a T cell mediated inflammatory skin disease. Complex network of cytokine induces and maintain the psoriatic lesion. Study was performed to evaluate the correlation between the level of TNF- α and IL-6, in the psoriasis patients with HLA-Cw6. Cytokine levels were assayed by sandwich ELISA and HLA-Cw6 typing was done by microcytotoxicity method. Serum TNF- α and IL-6 were significantly ($p < 0.001$ in both) expressed in psoriasis patients. TNF- α had significant positive correlation with IL-6 ($p = 0.56$, $p < 0.001$). We can conclude that TNF- α and IL-6 are found elevated in serum in psoriasis patients, compared to the control group. Proinflammatory cytokine such as TNF- α and IL-6 may cause inflammatory changes in microenvironment of psoriasis lesion. Gender of patient and age on onset of disease does not influence the expression of these cytokine in psoriasis cases. Results suggest that HLA-Cw6 did not influence these cytokines secretion.

Keywords: Psoriasis, TNF- α , IL-6, HLA-Cw6

Introduction

Psoriasis is characterized by two primary events hyper proliferation of keratinocyte and inflammatory dermal infiltration of mononuclear cells [1]. Previous studies had confirmed that altered T cells regulation along with complex network of cytokines are the main factors for induction and maintenance of various pathological stages of psoriasis [2-4]. Studies also demonstrated that proinflammatory cytokines TNF- α and IL-6 were markedly increased in the serum of psoriasis patients [5-9]. HLA antigen positive and negative individual differ in their ability to produce cytokines [10,11] or cytokines induce the expression of HLA antigen [12]. So present study was performed to evaluate correlation between the level of TNF- α and IL-6, in the psoriasis patients with HLA-Cw6.

Methods and Methods

In total of 60 psoriasis patients and 60 healthy controls were enrolled in this study. Neither patients nor control had present or past history of any skin disease or systemic or present history of infectious disease. Psoriasis patients who had not received any prior local or systemic treatment within three months were included in the study and were in active state of their disease.

About 5 ml of venous blood sample were collected from patients and controls in plain sterilized vials and 5 ml in heparinized vial for HLA-Cw6 typing. Serum sample was stored at -70°C until processed. Serum TNF- α and IL-6 were assayed using sandwich ELISA of Beckman Coulter, France. Typing for HLA-Cw6 antigen was performed by microcytotoxicity method of Terasaki and McClelland as described in detail by Mehra [13]. Antisera used for HLA-Cw6 typing was of BAG Company, Germany. The study was approved by ethical committee of the institution and informed consent was taken from all patients enrolled in the study.

Statistical Analysis

All data was analyzed using SPSS 14.0 (SPSS Inc., Chicago, IL, USA) computer statistics programme. Values were given as mean \pm Standard deviation (SD). Student t test used to compare mean, Pearson *Chi-square* used for comparing frequency and Pearson correlation was used to analysis of correlation. p values less than 0.05 were considered significant.

Results

A total of 120 subjects were enrolled in the study comprise of 60 psoriasis patients (median age 35 years) and 60 controls (median age 28 years). In patient group 35 males and 25 females were included and in control group 37 males and 23 females were included. Disease duration ranged from 0.5-30 years with the mean of 6.37 ± 6.35 years and age of onset ranged from 4- 69 years with the mean age of 30.48 ± 9.03 years. HLA-Cw6 was found positive in 20 cases (33.3%) of psoriasis while only six cases (10.0%) in control group. HLA-Cw6 was significantly expressed in patients with psoriasis as compared to control group with p value 0.002.

In controls mean value of serum TNF- α was 7.58 ± 4.19 pg/ml (range 2-15 pg/ml) while in psoriasis TNF- α was markedly increased (29.73 ± 17.31 pg/ml, range 9-74 pg/ml). Rise of TNF- α was highly significant ($p < 0.001$). Similarly serum IL-6 was markedly raised in psoriasis (36.88 ± 26.61 pg/ml, range 5-98 pg/ml) as compared to healthy control (5.56 ± 2.29 pg/ml, range 2-11 pg/ml) which was statically significant (Figures 1 and 2). In patients univariate analysis showed that serum TNF- α correlated significantly with IL-6 ($r = 0.26$, $p = 0.024$). HLA Cw6 showed positive correlation with family history ($r = 0.35$, $p = 0.003$) suggesting that HLA Cw6 was expressed in patients with positive family history psoriasis. No correlation was found among the cytokines and HLA Cw6 (Tables 1 and 2).

Discussion

Cytokines are essential for various pathological changes in the development of psoriasis. Cytokines form a complex and multidimensional network in psoriasis pathobiology, none of which alone can be considered to involve in disease causative mechanism [2].

***Corresponding author:** Sangeeta Singh, Women Scientist, Department of Biochemistry, King George's Medical University, Lucknow, India, Tel: 91 - (612) - 22232232; E-mail: sangeeta3.singh@gmail.com

Received February 06, 2016; **Accepted** April 20, 2016; **Published** April 27, 2016

Citation: Singh S, Sonkar GK, Singh S, Singh U (2016) Increased Serum Levels of TNF- α and IL-6 are not Related to HLA-Cw6 in Psoriasis Patients Correlation of Cytokine with HLA Cw6. Transcriptomics 4: 133. doi:[10.4172/2329-8936.1000133](https://doi.org/10.4172/2329-8936.1000133)

Copyright: © 2016 Singh S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

They influence keratinocyte proliferation, induce neutrophil and T cell chemotaxis, keep T cells in type 1 differentiation, enhance angiogenesis and upregulate adhesion molecules on endothelial cells, and stimulate the release of other chemokines. We found IL-6 and TNF- α to be elevated in our patient's group than in control group. Elevated serum levels of these cytokines were reported by most of the investigators [14-17]. TNF- α is produced by Langerhans cells, macrophages, monocytes, T cells and keratinocytes [18]. TNF- α is a proinflammatory cytokine and increases proliferation of keratinocytes [19]. TNF- α also induces

keratinocyte to produce IL-1, IL-6, IL-8 and adhesion molecules [6,20-22] all of which have proinflammatory properties. Autoantibody against TNF- α had also been reported in psoriasis patients [2]. Anti TNF- α therapy causes regression of the psoriatic lesions [4,19,23]. IL-6 is another proinflammatory cytokine which was included in our study. It is produced by keratinocytes, fibroblasts, endothelial cells and Th2 cells [24]. IL-6 is a major mediator of the host response to injury and infection. Some investigators reported that IL-6 enhances the activation, proliferation and chemotaxis of T lymphocytes in dermal infiltrate and also enhances proliferation and activation of B cells and macrophages [2,24]. Psoriatic keratinocyte are more sensitive to the growth-promoting effect of IL-6 than normal ones [2]. Increased level of IL-6 was also reported in psoriatic skin lesions [25,26]. It is speculated that the koebner phenomenon is likely to result from the increased activity of IL-6 and its receptor in psoriasis [27]. Antipsoriatic therapy including phototherapy, systemic and topical steroid reduced the increased level of IL-6 [1,25]. Elevated levels of TNF- α and IL-6 in psoriasis patients supports that they may cause inflammatory changes in microenvironment of psoriatic lesions and anti TNF therapy as well as anti-IL-6 therapy may be useful for treatment of psoriasis along with other drugs. TNF- α and IL-6 were positively correlated to each other suggesting that they influence the secretion of each other in psoriasis. Our results showed that TNF- α and IL-6 did not correlate with age of onset of psoriasis, gender, family history and HLA-Cw6. Our study for the first time correlated these parameters with cytokines in patients with psoriasis. Our results suggested that secretion of these cytokines are not age or gender biased. Association of HLA-Cw6 with psoriasis is well established. HLA-Cw6 expressing cells might affect the cytokine milieu in psoriasis and constitute an immune pathway important in psoriasis pathogenesis [28]. No correlation was found between elevated TNF- α and IL-6 with HLA-Cw6 antigen in our study suggests that HLA-Cw6 did not influence these cytokines secretion. Thus our study concludes that HLA-Cw6 may not influence the level of TNF- α and IL-6 in psoriasis patients. There may be some other factor(s) which elevation of these cytokines in psoriasis patients [29].




Figure 1: Showing serum TNF- α level in psoriasis patients and controls. Mean of TNF- α in psoriasis group and mean of TNF- α in control group.




Figure 2: Showing serum IL-6 level in psoriasis patients and controls. Mean of TNF- α in psoriasis group and mean of TNF- α in control group.

Cytokine	Group	Range (pg/ml)	Mean \pm SD (pg/ml)	p	CI (95%)	
					Lower	Higher
TNF- α	Psoriasis	9-74	29.75 \pm 17.31	<0.001*	16.96	26.07
	Control	2-15	7.58 \pm 4.19			
IL-6	Psoriasis	5-98	36.88 \pm 26.61	<0.001*	24.50	38.15
	Control	2-11	5.56 \pm 2.29			

*- significant (p <0.05)

Table 1: Serum level of TNF- α and IL-6 in patients of psoriasis and controls.

		TNF α	IL- 6	HLA-Cw6	Age of onset	Gender	Family History
TNF α	r	1	0.26	-0.41	0.05	-0.01	-0.03
	p		0.024*	0.377	0.362	0.467	0.399
IL- 6	r		1	0.11	-0.184	-0.23	-0.14
	p			0.211	0.080	0.042	0.144
HLA-Cw6	r			1	0.05	0.10	0.35
	p				0.356	0.234	0.003*
Age of onset	r				1	-0.004	0.09
	p					0.489	0.247
Gender	r					1	0.015
	p						0.467

*- significant (p <0.05)

Table 2: Correlation of cytokines with HLA-Cw6, age of onset, gender and family history.

References

- Ameglio F, Bonifati C, Fazio M, Mussi A, Trento E, et al. (1997) Interleukin-11 production is increased in organ cultures of lesional skin of patients with active plaque-type psoriasis as compared with nonlesional and normal skin. Similarity to interleukin-1 beta, interleukin-6 and interleukin-8. *Arch Dermatol Res* 289: 399-403.
- Bonifati C, Ameglio F (1999) Cytokines in psoriasis. *International Journal of Dermatology* 38: 241-251.
- Krueger J, Bowcock A (2005) Psoriasis pathophysiology: current concepts of pathogenesis. *Ann Rheum Dis* 64: ii30-ii36.
- Mehlis SL, Gordon KB (2003) The immunology of psoriasis and biologic immunotherapy. *J Am Acad Dermatol* 49: S44-S50.
- Bos JD, de Rie MA, Teunissen MB, Piskin G (2005) Psoriasis: dysregulation of innate immunity. *Br J Dermatol* 152: 1098-1107.
- Lowes MA, Bowcock AM, Krueger JG (2007) Pathogenesis and therapy of psoriasis. *Nature* 445: 866-873.
- Nickoloff BJ, Nestle FO (2004) Recent insights into the immunopathogenesis of psoriasis provide new therapeutic opportunities. *J Clin Invest* 113: 1664-1675.
- Prens EP, Benne K, van DJ, Bakkus M, Brakel K, et al. (1990) Interleukine-1 and interleukine- 6 in psoriasis. *J Invest Dermatol* 95: 121S.
- Lio D, Candore G, Romano GC, D'Anna C, Gervasi F, et al. (1997) Modification of cytokine patterns in subjects bearing the HLA-B8,DR3 phenotype: implications for autoimmunity. *Cytokines Cell Mol Ther* 3: 217-224.
- Mangalam AK, Veena T, David CS (2013) HLA class II molecules influence susceptibility versus protection in inflammatory diseases by determining the cytokine profile. *J Immunol* 190:513-518.

11. Johnson DR (2003) Locus-specific constitutive and cytokine-induced HLA class I gene expression. *J Immunol* 170: 1894-1902.
12. Mehra NK (1989) *Basic Methods in HLA-DNA Technology*. Technical Manual Published During the DBT Sponsored Training Workshop on HLA-DNA Technology. New Delhi: Sagar Publishers
13. Arican O, Aral M, Sasmaz S, Ciragil P (2005) Serum Levels of TNF-alpha, IFN-gamma, IL-6, IL-8, IL-12, IL-17 and IL-18 in Patients with Active Psoriasis and Correlation with Disease Severity. *Mediators Inflamm* 2005: 273-279.
14. Jacob SE, Nassiri M, Kerdel FA, Vincek V (2003) Simultaneous measurement of multiple Th1 and Th2 serum cytokines in psoriasis and correlation with disease severity. *Mediators Inflamm* 12: 309-313.
15. Pietrzak A, Koziol-Montewka M, Lecewicz-Toruń B, Krasowska D (2000) Is there any correlation between the total number of neutrophils in plasma and concentration of interleukin-8 in psoriatic patients? *Med Sci Monit* 6: 867-870.
16. Szepletowski JC, Bielicka E, Nockowski P, Noworolska A, Wasik F (2000) Increased interleukin-7 levels in the sera of psoriatic patients: lack of correlations with interleukin-6 levels and disease intensity. *Clin Exp Dermatol* 25: 643-647.
17. Krueger JG (2002) The immunologic basis for the treatment of psoriasis with new biologic agents. *J Am Acad Dermatol* 46: 1-23.
18. Pillai S, Bikle DD, Eessalu TE, Aggarwal BB, Elias PM (1989) Binding and biological effects of tumor necrosis factor alpha on cultured human neonatal foreskin keratinocytes. *J Clin Invest* 83: 816-821.
19. Cesare DC, Meglio PD, Nestle FO (2009) The IL-23/ Th 17 Axis in the Immunopathogenesis of Psoriasis. *J Invest Dermatol* 129: 1339-1350.
20. Grossman RM, Krueger J, Yourish D, Graneli-Piperno A, Murphy DP, et al. (1989) Interleukin 6 is expressed in high levels in psoriatic skin and stimulates proliferation of cultured human keratinocytes. *Proc Natl Acad Sci USA* 86: 6367-6371.
21. Makhatadze NJ (1998) Tumor necrosis factor locus: genetic organisation and biological implications. *Hum Immunol* 59: 571-579
22. Ware CF (2003) The TNF superfamily. *Cytokine Growth Factor Rev* 14: 181-184.
23. Chaudhari U, Romano P, Mulcahy L, Dooley L, Baker D, et al. (2001) Efficacy and safety of infliximab monotherapy for plaque-type psoriasis: a randomised trial. *Lancet* 357: 1842-1847
24. Nickoloff BJ (1991) The cytokine network in psoriasis. *Arch Dermatol* 127: 871-884.
25. Bonifati C, Solmone M, Trento E, Pietravalle M, Fazio M, et al. (1994) Serum interleukine-6 levels as an early marker of therapeutic response to UVB radiation and topical steroids in psoriatic patients. *Int J Clin Lab Res* 24: 122-123.
26. Mizutani H, Ohmoto Y, Mizutani T, Murata M, Shimizu M (1997) Role of increased production of monocytes TNF- α , IL-1 β and IL-6 in psoriasis: relation to focal infection, disease activity and responses to treatments. *J Dermatol Sci* 14: 145-153.
27. Toruniowa B, Krasowska D, Koziol M, Ksiazek A, Pietrzak A (1995) Serum levels of IL-6 in mycosis fungoides, psoriasis and lichen planus. *Ann N Y Acad Sci* 762: 432-434.
28. Mak R, Hundhausen C, Botti E, Laggner U, Grys K, et al. (2010) Demonstration of novel innate immune cells in psoriasis. *J Transl Med* 8: 18.
29. Murphy M, Philip K, Grant- Kels JM (2007) Histopathologic spectrum of psoriasis. *Clin Dermatol* 25: 524-528.

Citation: Singh S, Sonkar GK, Singh S, Singh U (2016) Increased Serum Levels of TNF-A and IL-6 are not Related to HLA-Cw6 in Psoriasis Patients Correlation of Cytokine with HLA Cw6. *Transcriptomics* 4: 133. doi:[10.4172/2329-8936.1000133](https://doi.org/10.4172/2329-8936.1000133)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700 Open Access Journals
- 50,000 editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>