

Indoor Air Pollution and Cardiovascular Health

Boris Z. Simkhovich*

Heart Institute, Good Samaritan Hospital, Division of Cardiovascular Medicine, Keck School of Medicine, University of Southern California, California, USA

Air pollution significantly increases cardiovascular morbidity and mortality [1-3]. During several past years, attention was drawn to Indoor Air Pollution (IAP) and its possible role as a risk factor for cardiovascular disease. National Human Activity Pattern Survey (sponsored by the Environmental Protection Agency) demonstrated that people spend most of their time indoors (up to 87%) [4]. While indoors, they are exposed to indoor air pollutants generated by cooking, combustion of biomass fuel (BMF, wood, charcoal, etc), use of electrical appliances, fireplaces, smoking and by ozone-associated chemical reactions involving household elements (e.g. linoleum, paint, adhesives, particle filters, ventilation ducts etc.) [5]. A significant source of Indoor Air Pollution (IAP) is smoking. Importantly, both mainstream and exhaled cigarette smoke generates fine particles (aerodynamic diameter $\leq 2.5 \mu\text{m}$, $\text{PM}_{2.5}$) [6]. Study of air quality in 66 US casinos revealed that in the smoking casinos means $\text{PM}_{2.5}$ concentration was $53.8 \mu\text{g}/\text{m}^3$ vs. $4.3 \mu\text{g}/\text{m}^3$ outside those casinos [7].

In addition to the indoor-specific pollutants, a great deal of IAP originates from outdoor sources, and then penetrates indoors. Polidori et al. demonstrated that in two retirement homes in the Los Angeles basin, the indoor measured Organic Carbon (OC) had on average 36-44% of outdoor-generated primary OC [8]. Penetration of outdoor pollutants indoors is characterized by infiltration efficiency (F_{inf}). F_{inf} represents the fraction of ambient particles that penetrate indoors and remains suspended. Study of home cohorts in six communities in Maryland, Illinois, North Carolina, California, New York and Minnesota found F_{inf} of 0.62 ± 0.21 for $\text{PM}_{2.5}$ [9]. These data indicate that indoor air contains considerable amount of ambient particulates from outdoors.

Significant adverse health effects are caused by ultrafine particles (aerodynamic diameter $< 100 \text{ nm}$, UFP). The main source of outdoor UFP is fuel combustion in motor vehicles, aircrafts and marine vessels. UFP were shown to actively penetrate buildings with closed windows (F_{inf} was 0.48 for particles $> 90 \text{ nm}$ in diameter). With opened windows F_{inf} for these particles increased significantly and comprised 0.72 [10].

The majority of the population in developing countries and residents in rural areas in developed countries extensively use Biomass Fuel (BMF) for cooking which leads to a significant release of air pollutants and decreases indoor air quality [11]. Improvement in open-fire stoves (adding enclosed combustion chamber and chimney for ventilation) in villages in San Marcos area in Guatemala, resulted in a greater than 2 times decrease in the indoor concentration of $\text{PM}_{2.5}$ (from 264 to $102 \mu\text{g}/\text{m}^3$), and was associated with 3.7 mm Hg decrease in systolic blood pressure (SBP) and 3.0 mm decrease in Diastolic Blood Pressure (DBP) in women [12].

In women living in six villages in northwestern Yunnan province in China, BMF combustion for cooking resulted in a median indoor concentration of $\text{PM}_{2.5}$ of $52 \mu\text{g}/\text{m}^3$ in summer and $105 \mu\text{g}/\text{m}^3$ during the winter. In female cohort older than 50 years, a 1-log- $\mu\text{g}/\text{m}^3$ increase in $\text{PM}_{2.5}$ (i.e. increase 10 times), was associated with 4.1 mm Hg increase in SBP and 1.8 mm Hg increase in DBP. Based upon a statistical model employed by the Asia Pacific Cohort Studies Collaboration,

the authors estimated that lowering SBP by 4 mm Hg among Asian women 50-59 years of age, would result in an 18% decrease in coronary artery disease and a 22% decrease in stroke, which would result in a staggering decrease in death rate by 230,900 deaths/year [13, 14]. Study in healthy young non-smoking students in Taipei (Taiwan) revealed a SBP increase by 1.85-2.64 mm Hg per interquartile range (IQR) of indoor coarse particles and by 1.94-2.99 mm Hg per IQR of indoor fine particles at 1-4-hour means. Increases of comparable magnitude were also found for DBP and heart rate [15].

More studies are needed to address the link between IAP and acute myocardial infarction, coronary artery disease, atherosclerosis, hypertension and congestive heart failure. However, data already published clearly indicate that IAP is a serious risk factor for cardiovascular disease. Health risk of IAP is underscored by the fact that disability-adjusted life-years (DALYs) lost due to IAP comprise 1,100 DALYs per 100,000 population [16]. On a positive note, there is evidence suggesting that measures aimed at fighting IAP, such as Clean Indoor Air Acts (banning indoor smoking) are effective in reducing the prevalence of cardiovascular disease [17].

References

1. Peters A, Dockery DW, Muller JE, Mittleman MA (2001) Increased particulate air pollution and the triggering of myocardial infarction. *Circulation* 103: 2810-2815.
2. Pope CA III, Muhlestein JB, May HT, Renlund DG, Anderson JL et al. (2006) Ischemic heart disease events triggered by short-term exposure to fine particulate air pollution. *Circulation* 114: 2443-2448.
3. Simkhovich BZ, Kleinman MT, Kloner RA (2008) Air pollution and cardiovascular injury: epidemiology, toxicology and mechanisms. *J Am Coll Cardiol* 52: 719-726.
4. Klepeis NE, Nelson WC, Ott WR, Robinson JP, Tsang AM et al. (2001) The national human activity pattern survey (NHAPS): a resource for assessing exposure to environmental pollutants. *J Expo Anal Environ Epidemiol* 11: 231-252.
5. Weschler CJ (2006) Ozone's impact on public health: contribution from indoor exposures to ozone and products of ozone-initiated chemistry. *Environ Health Perspect* 114: 1489-1496.
6. Sahu SK, Tiwari M, Bhangare RC, Pandit GG (2013) Particle size distribution of mainstream and exhaled cigarette smoke and predictive deposition in human respiratory tract. *Aerosol Air Quality* 13: 324-332.
7. Repace JL, Jiang RT, Acevedo-Bolton V, Cheng KC, Klepeis NE et al. (2011) Fine particle air pollution and secondary smoke exposures and risks inside 66 US casinos. *Environ Res* 111: 473-484.

*Corresponding author: Boris Z. Simkhovich, Heart Institute, Good Samaritan Hospital, 1225 Wilshire, Boulevard, Los Angeles, California 90017, USA, Tel: 213-977-4050; E-mail: simkhovi@usc.edu

Received November 19, 2013; Accepted November 20, 2013; Published November 23, 2013

Citation: Simkhovich BZ (2013) Indoor Air Pollution and Cardiovascular Health. *J Pollut Eff Cont* 1: e107 doi: [10.4172/jpe.1000e107](http://dx.doi.org/10.4172/jpe.1000e107)

Copyright: © 2013 Simkhovich BZ. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

8. Polidori A, Arhami M, Sioutas C (2007) Indoor/outdoor relationships, trends and carbonaceous content of fine particulate matter in retirement homes of the Los Angeles basin. *J Air Waste MA* 57: 366-379.
9. Allen RW, Adar SD, Avol E, Cohen M, Curl CL et al. (2012) Modeling the residential infiltration of outdoor PM_{2.5} in the multi-ethnic study of atherosclerosis and air pollution (MESA Air). *Environ Health Perspect* 120: 824-830.
10. Rim D, Wallace LA, Persily AK (2013) Indoor ultrafine particles of outdoor origine: importance of window opening area and fan operation condition. *Environ Sci Technol* 47: 1922-1929.
11. Kim KH, Jahan A, Kabir E (2011) A review of diseases associated with household air pollution due to the use of biomass fuels. *J Hazard Mat* 192: 425-431.
12. McCracken JP, Smith KR, Diaz A, Mittleman M, Schwartz J (2007) Chimney stove intervention to reduce long-term wood smoke exposure lowers blood pressure among Guatemalan women. *Environ Health Persp* 115: 996-1001.
13. Baumgartner J, Schauer JJ, Ezzati M, Lu L, Cheng C et al. (2011) Indoor air pollution in rural China. *Environ Health Perspect* 119: 1390-1395.
14. Asia Pacific Cohort Studies Collaboration (2003) Blood pressure indices and cardiovascular disease in the Asia Pacific region. A pooled analysis *Hypertension* 42: 69-75.
15. Lin LY, Lin CY, Chang KJ (2009) The effects of indoor particles on blood pressure and heart rate among young adults in Taipei, Taiwan. *Indoor Air* 19: 482-488.
16. Logue JM, Price PN, Sherman MH, Singer BC (2012) A method to estimate the chronic health impact of air pollutants in U.S. residences. *Environ Health Perspect* 120: 216-222.
17. Lippert WC, Gustat J (2012) Clean Indoor Air Acts reduce the burden of adverse cardiovascular outcomes. *Public Health* 126: 279-285.

Citation: Simkhovich BZ (2013) Indoor Air Pollution and Cardiovascular Health. J Pollut Eff Cont 1: e107 doi: 10.4172/jpe.1000e107

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>

