

Influence of Carrying Loads on Ratings of Perceived Exertion and Heart Rate during Walking

Siddhartha Sen* and Ajita

Punjabi University, Patiala, India

Abstract

Introduction: There is evidence that schoolbags may be a contributing factor for musculoskeletal complaints in schoolchildren. Several studies have reported an association between carrying heavily loaded schoolbags and musculoskeletal pain or discomfort.

Methods: 105 collegiate students were conveniently recruited in this study; heart rate (HR) and perceived exertion were measured immediately after the treadmill walking with the back pack, right side pack and left side pack. Subject has to walk for 20 min after that the HR was measured using pulse oximeter and perceived exertion was measured using Borg's RPE scale.

Results: It showed there is a significant increase in RPE with right side pack, with left side pack and with back pack after treadmill walking.

Conclusion: Ratings of Perceived Exertion (RPE) increased after treadmill walking with carrying load in right side pack, left side pack and back pack.

Keywords: Side pack; Back pack; Perceived exertion; Heart rate

Introduction

The backpack and side pack is one of the various forms of manual load carriage that provides flexibility in load carrying and is often used by the soldiers, school children as well as by the college students [1]. Backpack carriage is common among adults, collegiate students, schoolchildren and adolescents for daily transferring of personal belongings, laptops, books and stationeries to and from offices, colleges or schools. As the load of backpack is directly applied to the spine via the shoulder straps and external loading has been demonstrated to be associated with spinal disorders [2]. De Paula et al. [3] found that a significant number of students carry backpacks weighing more than 10% of their BWs most of the time. This heavy stress puts the student at an increased risk of injury [4].

There is evidence that schoolbags may be a contributing factor for musculoskeletal complaints in schoolchildren [5]. Several studies have reported an association between carrying heavily loaded schoolbags and musculoskeletal pain or discomfort [6]. Load carriage has been associated with an increased risk of musculoskeletal disorders in the back and upper and lower limbs in recreational hikers [7] with females suffering significantly higher injury rates than males when participating in the same hiking activities in outdoor education [8].

Backpack carried by collegiate students and their role in the development of musculoskeletal pain has been the subject of recent attention [9] and reducing backpack weight has been suggested as one prevention strategy to reduce hiking-related injury [10] with previous research recommending a 30% body weight (BW) as the maximum load for healthy adult males [11]. If it is necessary to carry loads more than 10% of BW, the modified backpack helps the integrity of the back and neck. Also, it helps to reduce muscular stresses on the back when the backpack exceeds the lifted weight of 10% of student's BW. Therefore, the use of the modified backpack is better than to use the commercial one [12].

Backpack straps often compress the anterior part of the shoulder, situated in the region over the brachial plexus, axillary artery and vein [13]. Therefore, if compression of these tissues occurs it may affect hand/arm

circulation and sensation. One study found that a third of backpackers report transient upper extremity paresthesias [14]. In some occupational categories, upper extremity and hand function is required to perform work while donning a heavy backpack. Therefore, reduced sensation and blood perfusion in the hand could impair task function.

Furthermore, several studies observed that carrying a heavy backpack causes a decrease in walking speed [15], a decrease in walking time [16], an increase in cardiorespiratory responses an increase loads on lumbar intervertebral discs [17], an increase in trunk forward lean [18] and an increase in foot-ground forces [19].

At present, however, community-based therapists are reporting an increase in the number of students requiring treatment for musculoskeletal injuries and discomfort [20]. This increase in the number of students as patients has been suggested to be related to students' increased use of computers. It is evident that a lot of research work has been done related to back pack and RPE but there is a paucity of research linking the involvement with side pack in their perceived exertion during walking, thus there is a need to do a work on that specific aspect. The purpose of the study is to find out the changes in perceived exertion after treadmill walking with backpack and side pack.

Methodology

Participants

105 collegiate student form Sardar Bhagwan Singh (PG) Institute of Biomedical Sciences and Research, Balawala, Dehradun were

*Corresponding author: Siddhartha Sen, Research Scholar, Punjabi University, Patiala, India, Tel: 919412985124; E-mail: siddhartha.pt@gmail.com

Received August 31, 2016; Accepted September 20, 2016; Published September 27, 2016

Citation: Sen S, Ajita (2016) Influence of Carrying Loads on Ratings of Perceived Exertion and Heart Rate during Walking. J Ergonomics 6: 176. doi: [10.4172/2165-7556.1000176](https://doi.org/10.4172/2165-7556.1000176)

Copyright: © 2016 Sen S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

conveniently recruited in this study. The detailed protocol of the research was explained and consent of every subject was taken prior to their inclusion in the study. The whole research was approved by the human subjects ethics review committee at the Punjabi University, Patiala, India. Healthy collegiate students of both male and female with age group of 17-25 years who carry their back pack/side pack at least two hours a day were included and subjects with history of any trauma and history of any cardiovascular conditions were excluded from the study.

A Custom made double-strap backpack was used in this study. An internal aluminum frame covered with rigid plastic material was used and position the center of gravity of the backpack approximately at the participant's T12 level [21]. Weights were attached to the custom made back pack about the midline of the backpack until the overall weight of the backpack was 10% [15] of the participant's BW. The lengths of the backpack straps were adjusted according to the participant's comfort at the beginning of the experiment to minimize the effect on the pulmonary function [22].

HR and perceived exertion were measured immediately after the treadmill walking with the back pack and side pack. Subject has to walk for 20 min after that the HR was measured using pulse oximeter and perceived exertion was measured using Borg's RPE scale. The data were recorded in back pack, right side pack and left side pack in different days.

Measurement of heart rate

Every subject was undergone treadmill walking to find out the muscular activity and exertion. Subject walked for 20 min at a speed of 4-5 km/hour with the right side pack then immediately after walking HR was measured using pulse oximeter. The pulse oximeter was placed in the index finger of the subjects and wait for readings. The digital screen will display the heart rate. The whole procedure was repeated with left side pack and back pack.

Measurement of Borg's Ratings of Perceived Exertion (RPE)

The Borg's RPE scale was used to assess whole body perceived exertion during treadmill exercise. Before the treadmill walking, all participants were read standardized instructions. [23] The low and high perceptual anchors for the RPE scale were established during the maximal treadmill walking exercise test. A rating of 6 (low anchor) was assigned to the lowest exercise intensity, whereas a rating of 20 was assigned to the highest exercise intensity [24]. The meaning of the perception of physical exertion was initially explained; this was defined as an intensity exertion, discomfort, stress, or fatigue that the individual feels during exercise. The following specific instructions were read to participants: "Please use this scale to translate into numbers your feelings of exertion while exercising. The numbers on the scale represent a range of feelings from very, very light to very hard. To help you select a number that corresponds to your feelings regarding the exercise, consider the following: When the exercise exertion feels very, very light, respond with a number 7. For example, you should respond with a number 7 when you are walking very slowly on the treadmill. When the exercise exertion feels very, very hard, respond with a number 19. For example, a response of 19 is appropriate when your feelings of exertion are the same as when you run on a treadmill almost as fast and hard as you can. If your exercise feelings are less intense than very, very light, respond with a number 6, and if your feelings are more intense than very, very hard, respond with a number 20. When you rate your overall exertion, be sure to select the number that most accurately represents your whole body's feelings. If the exercise exertion feels somewhere between very, very light (RPE 7) and very very hard (RPE 19) then you should give a number of between 7 and

19." The RPE values were recorded during the last 15 seconds of each minute throughout the treadmill walking test [25].

Data analysis and result

Paired sample t test was done to find out the changes of HR which was measured before and after treadmill walking and the result showed significant differences in all types of pack viz. right side pack, with left side pack and with back pack. When comparison of HR readings was done between different pack before and after treadmill walking, the result showed non-significant differences between right side pack and left side pack, and right side pack and back pack and significant differences between left side pack and back pack before treadmill walking. After treadmill walking results showed significant differences when compared between right side pack and left side pack, and left side pack and back pack (Tables 1 and 3 and Figures 1 and 3).

RPE also was measured before and after treadmill walking and the result showed significant differences with right side pack, with left side pack and with back pack. When comparison of RPE readings was done between different pack, the result showed significant differences between right side pack and left side pack, and right side pack and back pack before walking and showed significant differences between, right side pack and back pack and left side pack and back pack after treadmill walking (Tables 2, 4 and 5 and Figures 2 and 4).

Discussion

RPE was measured before and after treadmill walking and comparison was done between before and after treadmill walking in individuals with different pack, the results of the study showed significant differences with right side pack, with left side pack and with

	Minimum	Maximum	Mean	SD
Age	18.00	28	22.15	1.94
Height	150.00	188	164.54	7.73
Weight	40.00	78	65.23	7.53

Table 1: Demographic Statistics.




Figure 1: Mean Comparison of HR Readings of pre and post




Figure 2: Mean comparison of RPE (Rating of Perceived Exertion) of pre and post

	Mean ± SD		SEM	t-value	p-value	Significant
	Pre	post				
Right Side Pack	79.88 ± 4.58	96.86 ± 14.20	1.2451	13.638	0.0001	S
Left Side Pack	79.60 ± 4.31	101.16 ± 12.69	1.1494	18.758	0.0001	S
Back Pack	81.04 ± 6.22	96.72 ± 11.16	0.7147	21.932	0.0001	S

Table 2: Comparison of HR Readings of before & after treadmill walking with different packs.

	Mean ± SD		SEM	t-value	p-value	Significant
	Pre	post				
Right Side Pack	6.35 ± 0.48	10.66 ± 1.97	0.1713	25.178	0.0001	S
Left Side Pack	6.19 ± 0.39	10.74 ± 2.10	0.1815	25.074	0.0001	S
Back Pack	6.17 ± 0.37	10.09 ± 1.99	0.1710	22.943	0.0001	S

Table 3: Comparison of RPE (Rating of Perceived Exertion) Readings of right side and left side with different packs.

		SEM	t-value	p-value	Significant
pre	Right Side vs. Left side	0.5704	0.501	0.618	NS
	Right side vs. Back Pack	0.7780	1.493	0.138	NS
	Left side vs. Back Pack	0.6801	2.128	0.036	S
post	Right Side vs. Left side	1.745	2.460	0.016	S
	Right side vs. Back Pack	1.726	0.083	0.934	NS
	Left side vs. Back Pack	1.5681	2.830	0.006	S

Table 4: Comparison of HR between individuals with Different Pack in Pre reading.

		SEM	t-value	p-value	Significant
pre	Right Side vs. Left side	0.0543	2.979	0.004	S
	Right side vs. Back Pack	0.0502	3.602	0.0001	S
	Left side vs. Back Pack	0.0448	0.425	0.672	NS
post	Right Side vs. Left side	0.2230	0.342	0.733	NS
	Right side vs. Back Pack	0.2235	2.556	0.012	S
	Left side vs. Back Pack	0.2135	3.033	0.003	S

Table 5: Comparison of RPE between individuals with Different Pack in Pre reading.

exertion level and they feel a noticeable change in pain when carrying loads which supports with the findings of Devroey et al. [26] that loads of 0% is significantly different than loads of 10% BM. During the walking period, heart rates showed a significant difference, indicating that the student had reached a good exercise state at each workload. The trends of changing HR in this study were also found in other studies on loads carried by children [27]. Findings of Astrand and Rodahl [28] found that when a fit subject is exercising at less than 65% of maximum cardiovascular response is in a steady state.

Moreover, the increase in RPE during the treadmill walking can be explained by increments in perceived pulmonary efforts that are attributable to progressive hyperventilation. Indeed, leg effort should be near maximal from the very beginning of the exercise bout; conversely, ventilation would increase because of progressive metabolic acidosis and contribute markedly to RPE measures as the load was increased during the walking [29]. In addition, RPE has been correlated with several physiological indicators of exercise intensity such as oxygen consumption, blood lactate concentration, and HR during a variety of exercise protocols [30].

When comparison of RPE readings was done between different pack before and after treadmill walking, the result showed significant differences between right side pack and left side pack, and right side pack and back pack and non-significant differences between left side pack and back pack.

This might be because treadmill walking started with right side pack first then left side pack and lastly with back pack thus at the end the exertion was reached a steady level.

The HR was measured before and after treadmill walking and comparison was done between pre and post HR with different pack, the result showed significant differences in all types of pack viz. right side pack, with left side pack and with back pack that means HR was increased after treadmill walking with all types of packs. HR is the component has been commonly observed during static muscular exercise. Kilbom [31] has concluded in his review that the resulting increase in cardiac output during static contractions is mainly directed towards the peripheral parts of the body and only a small part is supplied to the myocardium. In this study standing while carrying a backpack, however, required no significant extra metabolic energy which is in agreement with other studies [32]. In most studies it has been assumed that the metabolic cost per kg load is not dependent on the total mass for loads carried centrally on the body [32].

The increased forward flexion of the spine will possibly increase the shear forces on the segments of the spinal column and it has been noted that sagittal and lateral shear should not be ignored in the assessment




Figure 3: Comparison of HR between individuals with Different Pack in pre and post




Figure 4: Comparison of RPE between individuals with Different Pack in pre and post

back pack, that means RPE was increased after treadmill walking with all types of pack. So when students carrying load that increases the

of lumbar load (Jager and Luttmann) [33]. This type of loading may be a factor contributing to chronic injuries and subsequent reports of low back pain. Although there was a strong trend towards higher discomfort in the poorer neck posture assumed for laptop use [34].

Conclusion

Based on our results and analysis, we can conclude that RPE increased after treadmill walking with carrying load in right side pack, left side pack and back pack.

References

1. Knapik J, Harman E, Reynolds K (1996) Load carriage using packs: A review of physiological, biomechanical and medical aspects. *Appl Ergon* 27: 207-216.
2. Korovessis P, Koureas G, Zacharatos S, Papazisis Z (2005) Backpacks, back pain, sagittal spinal curves and trunk alignment in adolescents: a logistic and multinomial logistic analysis. *Spine* 30: 247-55.
3. De Paula AJF, Silva JCP, Paschoarelli LC, Fujii JB (2012) Backpacks and school children's obesity: challenges for public health and ergonomics. *Work* 41: 900-906.
4. Mackie HW, Legg SJ, Beadlea J, Hedderley D (2003) Comparison of four different backpacks intended for school use. *Appl Ergon* 34: 257-264.
5. Negrini NS, Carabalona R (2002) Backpacks on! Schoolchildren's perceptions of load, associations with back pain and factors determining the load. *Spine* 27: 187-195.
6. Dianat I, Javadivala Z, Jafarabadi MA, Hashemi AA, Haslegrave CM (2013) The use of schoolbags and musculoskeletal symptoms among primary school children: are the recommended weight limits adequate? *Ergonomics* 56: 79-89.
7. Twombly SE, Schussman LC (1995) Gender differences in injury and illness rates on wilderness backpacking trips. *Wilderness Environ Med* 4: 363-376.
8. Leemon D, Schimelpfenig T (2003) Wilderness injury, illness, and evacuation: National Outdoor Leadership School's Incident Profiles. *Wilderness Environ Med* 14: 174-182.
9. Javadivala Z, Allahverdipour H, Dianat I, Bazargan M (2012) Awareness of parents about characteristics of a healthy school backpack. *Health Promot Perspect* 2: 166-172.
10. McIntosh SE, Leemon D, Visitacion J, Schimelpfenig T, Fosnocht D (2007) Medical incidents and evacuations on wilderness expeditions. *Wilderness Environ Med* 18: 298-304.
11. Haisman MF (1988) Determinants of load carrying ability. *Appl Ergon* 19: 111-121.
12. Mohamed Z, Ramadan, Al-Shayea AM (2013) A modified backpack design for male school children. *Int J Ind Ergonom* 43: 462-471.
13. Makela JP, Ramstad R, Mattila V, Pihlajamaki H (2003) Brachial plexus lesions after backpack carriage in young adults. *Clin Orthop Relat Res* 45: 205-209.
14. Boulware DR (2003) Backpacking-induced paresthesias. *Wilderness Environ Med* 14: 161-166.
15. Chow DH, Kwok ML, Au-Yang AC, Holmes AD, Cheng JC, et al. (2005) The effect of backpack load on the gait of normal adolescent girls. *Ergonomics* 48: 642-656.
16. Hong Y, Li JX, Fong DTP (2008) Effect of prolonged walking with backpack loads on trunk muscle activity and fatigue in children. *J Electromyography Kinesiol* 16: 990-996.
17. Neuschwander T, Cutrone BM, Cutrone S, Murthy G, Chambers H, et al. (2008) Typical school backpack loads significantly compress lumbar discs in children. In: *Proceedings of the NASS 23rd Annual Meeting. The Spine J* 8: S69.
18. Safikhani H, Fadilah T, Kamalden T, Amri SB, Ahmad M (2012) The effect of different backpack loading systems on trunk forward lean angle during walking among college students. *Eur J Sport Sci* 1: 1-5.
19. Pau M, Corona F, Leban B, Pau M (2011) Effects of backpack carriage on foot-ground relationship in children during upright stance. *Gait Posture* 33: 195-199.
20. Wilson K (1997) *Laptops a pain in neck*. Melbourne: Herald Sun.
21. Grimmer K, Dansie B, Milanese S, Pirunsan U, Trott P (2002) Adolescent standing postural response to backpack loads: a randomized controlled experimental study. *BMC Musculoskeletal Disorders* 3: 10.
22. Bygrave S, Legg SJ, Myers S, Llewellyn M (2004) Effect of backpack fit on lung function. *Ergonomics* 47: 324-329.
23. Robertson RJ (2001) Exercise testing and prescription using RPE as a criterion variable. *Int J Sport Psychol* 32: 177-188.
24. Noble BJ (1996) *Perceived Exertion*. Champaign, IL: Human Kinetics Books.
25. Elsangedy HM, Krinski K, Costa EC, Haile L, Fonteles AI, et al. (2013) The rating of perceived exertion is not different at the ventilatory threshold in sedentary women with different body mass indices. *J Exerc Sci Fit* 11: 102-106.
26. Devroey C, Jonkers I, de Becker A, Lenaerts G, Spaepen A (2007) Evaluation of the effect of backpack load and position during standing and walking using biomechanical, physiological and subjective measures. *Ergonomics* 50: 728-742.
27. Malhotra MS, Sen Gupta J (1965) Carrying of school bags by children. *Ergonomics* 8: 55-60.
28. Astrand PO, Rodahl K (1977) *Textbook of Work Physiology*. New York: McGraw-Hill.
29. Pereira G, Correia R, Ugrinowitsch C, Nakamura F, Rodacki A, et al. (2011) The rating of perceived exertion predicts intermittent vertical jump demand and performance. *J Sport Sci* 29: 927-932.
30. Eston R, Lambrick D, Sheppard K, Parfitt G (2008) Prediction of maximal oxygen uptake in sedentary males from a perceptually regulated, sub-maximal graded exercise test. *J Sport Sci* 26: 131-139.
31. Kilbom A (1976) Circulatory adaptation during static muscular contractions. A review. *Scand J Work Environ Health* 2: 1-13.
32. Pierrynowski MR, Winter DA, RW N (1981) Metabolic measures to ascertain the optimal load to be carried by man. *Ergonomics* 24: 393-399.
33. Jager M, Luttmann A (1992) The load on the lumbar spine during asymmetrical bi-manual materials handling. *Ergonomics* 35: 783-805.
34. Straker L, Miller J, Mangharam J, Bates M, Pollock C (1992) Performance effects of postural changes at a VDU work station.

Citation: Sen S, Ajita (2016) Influence of Carrying Loads on Ratings of Perceived Exertion and Heart Rate during Walking. *J Ergonomics* 6: 176. doi: 10.4176/2165-7556.1000176

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>