



iNKT Cells in Lupus Development

Yang JQ^{1,2*} and Singh RR²

¹Key Laboratory of National Health and Family Planning Commission on Parasitic Disease Control and Prevention, Jiangsu Institute of Parasitic Diseases, Jiangnan University, Wuxi, China

²Autoimmunity and Tolerance Laboratory, Division of Rheumatology, Department of Medicine, David Geffen School of Medicine at University of California Los Angeles, Los Angeles, USA

*Corresponding author: Jun-Qi Yang, Key Laboratory of National Health and Family Planning Commission on Parasitic Disease Control and Prevention, Jiangsu Institute of Parasitic Diseases, Jiangnan University, Wuxi, China, E-mail: yangjunqi@hotmail.com

Received date: December 2, 2016; Accepted date: December 10, 2016; Published date: December 20, 2016

Copyright: © 2016 Yang JQ, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Yang JQ, Singh RR (2016) iNKT Cells in Lupus Development. Lupus Open Access 1: e103.

Editorial

Invariant natural killer T (iNKT) cells are a new subset of T cells that bridge innate and acquired immunity and play important roles in both protective and regulatory responses in a variety of diseases [1,2]. Our group has worked on the role and the underlying molecular mechanisms of iNKT cells in lupus development for more than a decade and published a dozen of research articles in peer reviewed journals. Here, we briefly summarize and comment our findings.

Several promising and important findings are made on the features and roles of iNKT cells in three murine models of lupus. Firstly, we revealed for the first time that iNKT cells were deficient in both number and function in lupus-prone MRL-*lpr* mice. Moreover, repeated administration of α -GalCer, a glycolipid and natural activator of iNKT cells, expanded iNKT cell number and function in these mice [3]. This finding revealed that α -GalCer was able to induce iNKT cell expansion *in vivo*. Before that, NKT cells were generally believed to undergo apoptosis upon α -GalCer treatment. Consistent to this finding, MRL-*lpr* mice with CD1d-deficiency, which blocks iNKT cell development, exacerbated the frequency and severity of skin lesions [4]. Treatment with α -GalCer ameliorated the dermatitis in MRL-*lpr* mice [3]. Secondly, in an induced lupus model in BALB/c mice by hydrocarbon oil pristane (2,6,10,14-tetramethylpentadecane, TMP), iNKT cells were found to be functionally insufficient. CD1-deletion in these mice further increased the level of serum autoAbs and induced more severe glomerulonephritis [5]. In addition, repeated α -GalCer treatment of pristane-inoculated BALB/c mice prior to the onset of florid disease suppressed proteinuria [6]. Thirdly, (NZB \times NZW) F1 (BWF1) mice showed intrinsic deficiency of iNKT cell function [7]. Deficiency of CD1 in BWF1 mice further worsened the glomerulonephritis as compared to wild-type mice [8]. Similar findings are evident in β 2-microglobulin knockout BWF1 mice, where iNKT cells are also deficient as CD1 knockout mice [9]. Interestingly, activation of iNKT cells with α -GalCer in BWF1 mice at an early age by short term ameliorated the glomerulonephritis [10]. Moreover, iNKT cells can directly regulate autoreactive B cells in a contact and CD1-dependent manner. This regulation was related to the impaired IL-10 production from B cells, reduced number and function of marginal zone B cells and involved Fas pathway [11,12].

The above studies indicate the regulatory role of iNKT cells in lupus development. These findings should be helpful to delineate the pathogenesis of lupus and may have the potential leading to new

therapeutics for lupus, and perhaps other autoimmune diseases, based on manipulation of iNKT cells. Further in-depth studies are needed to explore the underlying molecular mechanisms how iNKT cells control lupus development.

References

1. Bendelac A, Savage PB, Teyton L (2007) The biology of NKT cells. *Annu Rev Immunol* 25: 297-336.
2. Wu L, Gabriel CL, Parekh VV, Van Kaer L (2009) Invariant natural killer T cells: innate-like T cells with potent immunomodulatory activities. *Tissue Antigens* 73: 535-545.
3. Yang JQ, Saxena V, Xu H, Van Kaer L, Wang CR, et al. (2003) Repeated α -galactosylceramide administration results in expansion of NK T cells and alleviates inflammatory dermatitis in MRL-*lpr/lpr* mice. *J Immunol* 171: 4439-4446.
4. Yang JQ, Chun T, Liu H, Hong S, Bui H, et al. (2004) CD1d deficiency exacerbates inflammatory dermatitis in MRL-*lpr/lpr* mice. *Eur J Immunol* 34: 1723-1732.
5. Yang JQ, Singh AK, Wilson MT, Satoh M, Stanic AK, et al. (2003) Immunoregulatory role of CD1d in the hydrocarbon oil-induced model of lupus nephritis. *J Immunol* 171: 2142-2153.
6. Singh AK, Yang JQ, Parekh VV, Wei J, Wang CR, et al. (2005) The natural killer T cell ligand α -galactosylceramide prevents or promotes pristane-induced lupus in mice. *Eur J Immunol* 35: 1143-1154.
7. Yang JQ, Kim PJ, Halder RC, Singh RR (2013) Intrinsic hyporesponsiveness of invariant natural killer T cells precedes the onset of lupus. *Clin Exp Immunol* 173: 18-27.
8. Yang JQ, Wen X, Liu H, Folyan G, Dong X, et al. (2007) Examining the role of CD1d and natural killer T cells in the development of nephritis in a genetically susceptible lupus model. *Arthritis Rheum* 56: 1219-1233.
9. Singh RR, Yang JQ, Kim PJ, Halder RC (2013) Germline deletion of beta2 microglobulin or CD1d reduces anti-phospholipid antibody, but increases autoantibodies against non-phospholipid antigens in the NZB/W F1 model of lupus. *Arthritis Res Ther* 15: R47.
10. Yang JQ, Kim PJ, Singh RR (2012) Brief Treatment with iNKT Cell Ligand α -Galactosylceramide Confers a Long-term Protection Against Lupus. *J Clin Immunol* 32: 106-113.
11. Yang JQ, Wen X, Kim PJ, Singh RR (2011) Invariant NKT cells inhibit autoreactive B cells in a contact- and CD1d-dependent manner. *J Immunol* 186: 1512-1520.
12. Wen X, Yang JQ, Kim PJ, Singh RR (2011) Homeostatic regulation of marginal zone B cells by invariant natural killer T cells. *PLoS One* 6: e26536.