

## Lipid Raft and Platelet SNARE Machinery

Fadi T Khasawneh and Zubair A Karim\*

Department of Pharmaceutical Sciences, College of Pharmacy, Western University of Health Sciences, Pomona, California, USA

Lipid rafts are defined as insoluble areas in the cell membrane, resistant to non-ionic detergents. These areas, which are also called detergent-resistant membranes (DRMs), and are enriched in glycosphingolipids, saturated phospholipids and cholesterol, have been identified in several cell types including platelets. Initially, they were believed to be responsible for the intercellular transport of glycosyl phosphatidylinositol (GPI)-anchored proteins to the apical surface in polarized cells [1,2]. However, over the decades, rafts have increasingly been recognized as membrane microdomains, and found to play a critical role in the control of several cellular activation processes. Thus, very divergent proteins such as Src family kinases, caveolins, palmitoylated proteins such as G proteins, GPI-anchored proteins such as Thy-1 and alkaline phosphatases, tetraspannin proteolipids and various signaling molecules have all been shown to be associated with lipid rafts. Different types of rafts coexist at the plasma membrane with functionally distinct lipid composition [3]. Furthermore, lipid rafts are not only found at the plasma membrane, but also as part of the internal membrane of granules, Golgi complex and even phagosomes [4,5]. Evidence for a functional role of lipid rafts in platelets is very recent: Gousset et al. [6] have shown that during cold-induced platelet activation, rafts cluster into larger aggregates, a reversible process depending on platelet activation. These authors showed raft aggregation to be dependent on the presence of cholesterol in the membrane, and further identified the presence of CD36 in DRMs. Using fluorescence microscopy of platelets being activated with thrombin and collagen, large fluorescent clusters of lipid rafts were formed, leading these investigators to conclude that raft aggregation is triggered by platelet activation, suggestive of a role for microdomains in platelet signaling [6].

An important physiological consequence of human platelets signaling is exocytosis, a process that involves secretory granule fusion with the platelet plasma membrane. This membrane fusion reaction is mediated by membrane proteins called Soluble *N*-ethylmaleimide sensitive factor attachment protein receptors (SNAREs). In platelets, vesicle-associated membrane protein (VAMP)-8/endobrevin (a vesicle, or v-SNARE) is required for each granule type [7]. Two target membrane SNAREs (*t*-SNAREs) are present: the syntaxin class (syntaxin-11 being functionally relevant) [8] and the SNAP-23/25 class (SNAP-23 being functionally relevant) [9-14]. Either immediately preceding or concurrent with membrane fusion, three cognate SNARE proteins on opposing membranes assemble, through hydrophobic interactions in their coiled-coil domains, into a parallel, four helical bundle [15-17]. This so-called *trans* complex, containing one copy each of SNAP-23/syntaxin/VAMP, is minimally required for membrane fusion [18] and on fusion, the SNAREs exist in the same membrane in a thermodynamically stable *cis* configuration [19]. This *cis* SNARE complex has to be disassembled by the adapter proteins *N*-ethylmaleimide-sensitive fusion protein and  $\alpha$ -SNAP in a process called SNARE priming [20,21], thereby allowing SNAREs to be reused in another round of membrane fusion. However, what the function of NSF and  $\alpha$ -SNAP is in platelets is still unknown, and how the SNARE recycling takes place in platelets is also unknown. The SNAREs form a *trans*-bilayer complex, which facilitates fusion of the granule and plasma membranes for cargo release. Although SNAREs are essential for fusion, how, when, and where they assemble into fusogenic complexes represent key secretion control points. Thus, SNARE complex assembly is determined by numerous SNARE regulators (i.e.,

Munc18s and Munc13s) and post-translational modifications (i.e., acylation and phosphorylation) [15].

The SNARE proteins have also been reported to associate with lipid rafts in other secretory cells [22,23]. SNAP-23 and SNAP-25 associate with membranes by palmitoylation [24-28], and it is likely that this post-translational modification is responsible for the association of these proteins with lipid rafts. By contrast, members of the syntaxin and synaptobrevin/VAMP family are generally not palmitoylated, and their mechanism of association with rafts remains unknown. In addition, it is unclear to what extent individual SNARE subunits or binary/ternary SNARE complexes associate with lipid raft membranes or whether the nature of SNARE complexes associated with rafts is altered during the process of regulated exocytosis.

Karim et al. [8] have been examining the membrane distribution of individual SNARE proteins and SNARE complexes that regulate granule exocytosis from human platelets (unpublished). The role of lipid rafts in platelet exocytosis is still unclear. Nonetheless, it is interesting that SNAP-23 appears to localize to rafts yet for biophysical reasons, rafts appear to be an unlikely site for membrane fusion events given the rigidity of their structure. It seems possible that phosphorylation could be an exclusion signal that prevents a portion of the acylated SNAP-23 from associating with the rafts.

Finally, it is very important to notice that cholesterol is at the core of the platelet lipid-raft function. Could cholesterol-lowering drugs directly affect platelet function, and could their action be via an effect on the platelet rafts? Certainly statins have been shown to reduce platelet-dependent thrombus formation [29], and such effects could explain the rapid therapeutic benefits of statin therapy [30]. Much further research lipid rafts is warranted and also it is important to understand how lipid rafts are involved in platelet exocytosis.

### References

1. Simons K, van Meer G (1988) Lipid sorting in epithelial cells. *Biochemistry* 27: 6197-6202.
2. Van Meer G, Stelzer EH, Wijnaendts-van-Resandt RW, Simons K (1987) Sorting of sphingolipids in epithelial (Madin-Darby canine kidney) cells. *J Cell Biol* 105: 1623-1635.
3. Schnitzer JE, McIntosh DP, Dvorak AM, Liu J, Oh P (1995) Separation of caveolae from associated microdomains of GPI-anchored proteins. *Science* 269: 1435-1439.
4. Gkantiragas I, Brügger B, Stüven E, Kaloyanova D, Li XY, et al. (2001) Sphingomyelin-enriched microdomains at the Golgi complex. *Mol Biol Cell* 12: 1819-1833.
5. Dermine JF, Duclos S, Garin J, St-Louis F, Rea S, et al. (2001) Flotillin-1-

\*Corresponding author: Zubair A Karim, Department of Pharmaceutical Sciences, College of Pharmacy, Western University of Health Sciences, Pomona, California 91766, USA, Tel: 909-469-8204; E-mail: [zkarim@westernu.edu](mailto:zkarim@westernu.edu)

Received March 27, 2014; Accepted March 28, 2014; Published April 08, 2014

Citation: Khasawneh FT, Karim ZA (2014) Lipid Raft and Platelet SNARE Machinery. *J Glycomics Lipidomics* 4: e119. doi:10.4172/2153-0637.1000e119

Copyright: © 2014 Khasawneh FT, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

- enriched lipid raft domains accumulate on maturing phagosomes. *J Biol Chem* 276: 18507-18512.
6. Gousset K, Wolkers WF, Tsvetkova NM, Oliver AE, Field CL, et al. (2002) Evidence for a physiological role for membrane rafts in human platelets. *J Cell Physiol* 190: 117-128.
  7. Ren Q, Ye S, Whiteheart SW (2008) The platelet release reaction: just when you thought platelet secretion was simple. *Curr Opin Hematol* 15: 537-541.
  8. Ye S, Karim ZA, Al Hawas R, Pessin JE, Filipovich AH, et al. (2012) Syntaxin-11, but not syntaxin-2 or syntaxin-4, is required for platelet secretion. *Blood* 120: 2484-2492.
  9. Ren Q, Barber HK, Crawford GL, Karim ZA, Zhao C, et al. (2007) Endobrevin/VAMP-8 is the primary v-SNARE for the platelet release reaction. *Mol Biol Cell* 18: 24-33.
  10. Karim ZA, Zhang J, Banerjee M, Chicka MC, Al Hawas R, et al. (2013) IκB Kinase phosphorylation of SNAP-23 controls platelet secretion. *Blood* 121: 4567-4574.
  11. Chen D, Bernstein AM, Lemons PP, Whiteheart SW (2000) Molecular mechanisms of platelet exocytosis: role of SNAP-23 and syntaxin 2 in dense core granule release. *Blood* 95: 921-929.
  12. Chen D, Lemons PP, Schraw T, Whiteheart SW (2000) Molecular mechanisms of platelet exocytosis: role of SNAP-23 and syntaxin 2 and 4 in lysosome release. *Blood* 96: 1782-1788.
  13. Lemons PP, Chen D, Whiteheart SW (2000) Molecular mechanisms of platelet exocytosis: requirements for alpha-granule release. *Biochem Biophys Res Commun* 267: 875-880.
  14. Weber T, Parlati F, McNew JA, Johnston RJ, Westermann B, et al. (2000) SNAREpins are functionally resistant to disruption by NSF and alphaSNAP. *J Cell Biol* 149: 1063-1072.
  15. Snyder DA, Kelly ML, Woodbury DJ (2006) SNARE complex regulation by phosphorylation. *Cell Biochem Biophys* 45: 111-123.
  16. Poirier MA, Xiao W, Macosko JC, Chan C, Shin YK, et al. (1998) The synaptic SNARE complex is a parallel four-stranded helical bundle. *Nat Struct Biol* 5: 765-769.
  17. Sutton RB, Fasshauer D, Jahn R, Brunger AT (1998) Crystal structure of a SNARE complex involved in synaptic exocytosis at 2.4 Å resolution. *Nature* 395: 347-353.
  18. Weber T, Zemelman BV, McNew JA, Westermann B, Gmachl M, et al. (1998) SNAREpins: minimal machinery for membrane fusion. *Cell* 92: 759-772.
  19. Otto H, Hanson PI, Jahn R (1997) Assembly and disassembly of a ternary complex of synaptobrevin, syntaxin, and SNAP-25 in the membrane of synaptic vesicles. *Proc Natl Acad Sci U S A* 94: 6197-6201.
  20. Mayer A, Wickner W, Haas A (1996) Sec18p (NSF)-driven release of Sec17p (alpha-SNAP) can precede docking and fusion of yeast vacuoles. *Cell* 85: 83-94.
  21. Barnard RJ, Morgan A, Burgoyne RD (1997) Stimulation of NSF ATPase activity by alpha-SNAP is required for SNARE complex disassembly and exocytosis. *J Cell Biol* 139: 875-883.
  22. Chamberlain LH, Burgoyne RD, Gould GW (2001) SNARE proteins are highly enriched in lipid rafts in PC12 cells: implications for the spatial control of exocytosis. *Proc Natl Acad Sci U S A* 98: 5619-5624.
  23. Gil C, Soler-Jover A, Blasi J, Aguilera J (2005) Synaptic proteins and SNARE complexes are localized in lipid rafts from rat brain synaptosomes. *Biochem Biophys Res Commun* 329: 117-124.
  24. Gonzalo S, Linder ME (1998) SNAP-25 palmitoylation and plasma membrane targeting require a functional secretory pathway. *Mol Biol Cell* 9: 585-597.
  25. Gonzalo S, Greentree WK, Linder ME (1999) SNAP-25 is targeted to the plasma membrane through a novel membrane-binding domain. *J Biol Chem* 274: 21313-21318.
  26. Koticha DK, Huddleston SJ, Witkin JW, Baldini G (1999) Role of the cysteine-rich domain of the t-SNARE component, SYNDET, in membrane binding and subcellular localization. *J Biol Chem* 274: 9053-9060.
  27. Veit M, Söllner TH, Rothman JE (1996) Multiple palmitoylation of synaptotagmin and the t-SNARE SNAP-25. *FEBS Lett* 385: 119-123.
  28. Vogel K, Cabaniols JP, Roche PA (2000) Targeting of SNAP-25 to membranes is mediated by its association with the target SNARE syntaxin. *J Biol Chem* 275: 2959-2965.
  29. Thompson PD, Moyna NM, White CM, Weber KM, Giri S, et al. (2002) The effects of hydroxy-methyl-glutaryl co-enzyme A reductase inhibitors on platelet thrombus formation. *Atherosclerosis* 161: 301-306.
  30. Wierzbicki AS, Poston R, Ferro A (2003) The lipid and non-lipid effects of statins. *Pharmacol Ther* 99: 95-112.

Citation: Khasawneh FT, Karim ZA (2014) Lipid Raft and Platelet SNARE Machinery. *J Glycomics Lipidomics* 4: e119. doi:10.4172/2153-0637.1000e119

### Submit your next manuscript and get advantages of OMICS Group submissions

#### Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

#### Special features:

- 350 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: [www.editorialmanager.com/omicsgroup](http://www.editorialmanager.com/omicsgroup)

