

## Macrobenthic Community Structure as A Bio-Indicator for the Assessment of Coastal Water Pollution In Greater Noakhali-Bangladesh

Md. Jahangir Sarker\*, Mehedi Hasan Tanmay, Farhana Rahman, Md. Shamsul Alam Patwary and Nazmun Nahar Rima

Department of Fisheries and Marine Science, Noakhali Science and Technology University, Noakhali-3814, Bangladesh

### Abstract

Present study was aimed for the partial judgement of coastal water pollution and accordingly the abundance, diversity as well as the species composition of macrobenthic communities on the sediment of 3 selected stations were carried out following standard methods during January to March, 2015 in Noakhali, Bangladesh. 14 families under 4 major groups yielded a total of 5481 ind./m<sup>2</sup> with a mean density of 609 ± 445 ind./m<sup>2</sup> during the study. The highest and the lowest number of taxa were identified in January (10) and February (6) respectively during the study period. On the other hand, monthly diversity profile of microbenthic communities showed clear difference in the diversity of all stations which is compatible to the other published results of macrobenthic animal as well. Based on the calculated results from Shannon-Wiener index of microbenthic species diversity (H'), the present study might be considered as heavily polluted in February (H' = 0.9202) and moderately polluted in January (H' = 1.514) and in March (H' = 1.571). On the other hand, The Margalef's index of benthic macro-invertebrates at station S1 (0.4501) was lowest in February and at station S1 (1.096) was highest. Therefore, the research output revealed that the abundance and diversity of macrobenthic animal community might be good indicators to assess the aquatic environmental pollution as the selected study area fall under the category of more or less polluted.

**Keywords:** Environmental pollution; Macrobenthic animal; Bangladesh

### Introduction

Macrobenthos are commonly used as bio-monitors to detect pollution impacts in estuaries [1] for their exclusive and unique characteristics. They are abundant, easy to collect and very diverse with representatives from many different phyla [2] utilizing many different habitats and feeding strategies [3-6]. These assemblages respond predictably to pollution [7,8], integrates the stress over months to years and relatively sedentary in nature [5,9]. Besides this, macrobenthic communities are eaten by other higher tropical organisms like fin and shell fishes as food and recycle the organic matters and debris like an ecological engineer [10]. They also provide a linkage between substratum and sea bed and water column predators [11]. Macrobenthos are an important role playing community in aquatic ecosystem because they mineralize, promote and mix the oxygen flux into the sediment which recycle the organic matter [12]. Benthic community determines the amount of nutrients release of the sediments [13]. Biological interactions, such as predation and competition affect the macrobenthic community structure by acting on recruitment, survival, or migration of organisms [14-18]. The characteristics of the life cycle of the species and the influence of temporal fluctuation of abiotic factors, such as environmental temperature or salinity can also change macrobenthic communities in a cyclic pattern over time [19-22]. Physical and chemical factors affect the distribution and abundance of macrobenthic community like contaminations of sediments environment, current of the water organic contents of the sediments, depth, rapid sedimentation and toxicity of sediments causes shifts of macrobenthic communities towards lower abundance [23].

Although much studies on macrobenthic community has been carried out globally however there is scanty published documents on macrobenthic community of coastal area in Bangladesh. There are few works on Hatia and Nijhum Dweep Island but the attached part of Hatia with main land of Noakhali district remain untouched. So this study was aimed for the documentation of the abundance and composition of microbenthic community of the branch of lower part of the mighty

Meghna River. Therefore the objective of the present study was to know the monthly abundance of macrobenthic community, to know the species composition and diversity of macrobenthic communities in estuarine waters and to assess the status of pollution in Chairman Ghat, Noakhali.

### Materials and Methods

Sediment and water samples were collected from the intertidal zone of the estuary of a branch of the Meghna River at Chairman Ghat during January, February and March, 2015 from 3 sampling stations namely S1 (Station 1) (22° 30' 48.3876" N, 91° 5' 6.6078" E), S2 (Station 2) (22° 31' 5.2278" N, 91° 5' 26.4768" E) and S3 (Station 3) (22° 31' 34.1868" N, 91° 5' 41.5566" E). An Ekman dredge (mouth opening of 0.0225 m<sup>2</sup>) was used to collect sediment samples with triplicate fashion from each station. Subsurface water samples were collected from three sites during high tide condition for measuring water temperature (°C), salinity (ppt), pH, DO (dissolved oxygen, ppm) and transparency (cm). Temperature (°C) was measured using a centigrade thermometer in study area where water salinity, pH, transparency and DO were measured *in situ* by using Refractometer (INDEX, Model No. REF 201), Digital pen pH meter (HANNA Instrument, Model No. H196107), Secchi disc (20 cm diameter) and DO meter (LUTRON, Model No. DO-5509) respectively.

**\*Corresponding author:** Md. Jahangir Sarker, Department of Fisheries and Marine Science, Noakhali Science and Technology University, Noakhali-3814, Bangladesh, Tel: +8801733910237/+880-321-71487; Fax: +880-321-62788; E-mail: [swaponj@yahoo.com](mailto:swaponj@yahoo.com)

Received May 02, 2016; Accepted May 25, 2016; Published May 31, 2016

**Citation:** Sarker J, Tanmay MH, Rahman F, Patwary MSA, Rima NN (2016) Assessment of Coastal Water Pollution In Greater Noakhali-Bangladesh. J Coast Zone Manag 19: 427. doi:[10.4172/2473-3350.1000427](https://doi.org/10.4172/2473-3350.1000427)

**Copyright:** © 2016 Sarker J, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Sampled sediments were sieved (through 500 µm mesh) to retain macrobenthos and then preserved immediately in the plastic container with other residues in 10% formalin solution. It was then labeled and for further analysis transferred to the laboratory. Small amount of “Rose Bengal” was added to make the macro organism visible. Benthic macrofauna was sorted manually putting on a tray under sufficient light and enumerated up to major taxa. Magnifying glass and microscope (Model No. XSZ 21-05DN, China) often were used for the identification of macrobenthos. Abundance of macrobenthic fauna was calculated in individual per m<sup>2</sup>. Sample was then preserved in 70% alcohol (Ethanol). Statistical analysis was performed using SPSS (Version 20) and PAST [24]. Occurrence of macro-invertebrates occurrence (N) was calculated using the following formula [25],

$$N = \frac{O}{a.s} \times 10000$$

Here,

s = Number of sample.

a = Mouth opening area of Ekman dredge in cm<sup>2</sup>, and

O = Actual counted number of macro-invertebrates per sampled area,

The dominance (D) index [26] was determined by using following formula:

$$D = \sum_{i=1}^s \left( \frac{ni}{n} \right)^2$$

Here,

n = Total individual number.

ni = Individuals of species number i.

The Simpson Index (1-D) was measured by the following formula [27],

$$(1-D) = \frac{1}{\sum_{i=1}^s Pi^2}$$

Here,

P = Proportion (n/N) of individuals of one particular species found (n) divided by the total number of individuals found (N),

S = Number of species.

Shannon-Wiener Index (Species diversity, H') were calculated according to Wilhm and Dorris [28] formula,

$$H' = - \sum_{i=1}^s Pi \ln Pi$$

Here,

Pi = ni/N = Proportion of individuals of the total sample of the i<sup>th</sup> species.

N = Individual number of all the species in total,

ni = Individuals number of the i<sup>th</sup> species, and

S = Number of species.

The Margalef's Index (D') was calculated by the following index [29],

$$D' = \frac{S-1}{\ln N}$$

Here,

N = Individuals number in total of sample

S = Species number in sample

ln = Normal log

Equitability (J) was measured by using the following formula [26],

$$J = \frac{H'}{\log S}$$

Here,

S = Species number in a population.

H' = Shannon and Weiner Index

## Results and Discussion

The measured water quality parameter is recorded in Table 1. The highest water temperature was recorded (32°C) in station S3 in March with mean of 30 ± 0.96 while the lower was observed in January (22.33 ± 0.47). Higher salinity was measured in February (10.33 ± 1.25) being the highest at station S3 (12 ppt) comparing to other two months while DO was lowest in this month (4.0 ± 0.29 ppm) in relation to January (6.8 ± 0.24 ppm) and March (5.3 ± 0.73 ppm). PH showed no major variation in January (7.67 ± 0.31), February (7.77 ± 0.21) or in March (7.28 ± 0.70) while transparency was recorded very low in all 3 month of the present study area in January (3.17 ± 1.03 cm), February (4.17 ± 0.85 cm), and in March (1.5 ± 0.71cm). All the parameters varied with the seasonal variation accordingly. Salinity and transparency

Month	Station	Temperature (°C)	Salinity (ppt)	Dissolve Oxygen (ppm)	pH	Transparency (cm)
January	S1	22	5	7.1	7.5	4.5
	S2	22	4	6.5	7.4	3
	S3	23	6	6.8	8.1	2
	Mean ± SD	22.33 ± 0.47	5.00 ± 0.82	6.8 ± 0.24	7.67 ± 0.31	3.17 ± 1.03
February	S1	27	9	3.6	7.8	3
	S2	27	10	4.3	7.5	5
	S3	28	12	4.1	8	4.5
	Mean ± SD	27.33 ± 0.47	10.33 ± 1.25	4.0 ± 0.29	7.77 ± 0.21	4.17 ± 0.85
March	S1	30	5	6.2	7.28	3
	S2	31	6	6.8	7.55	2
	S3	32	8	5.3	8.75	1.5
	Mean ± SD	30 ± 0.96	5 ± 1.41	5.3 ± 0.73	7.28 ± 0.70	1.5 ± 0.71

Table 1: Water quality parameter measured from three stations during January, February and March (2015).

may fluctuated due to the water availability in different month from January to March because usually March is more close to the monsoon season where January falls under Winter season in this area. A Pearson's correlation was run to determine the relationship between these 5 measured parameters (temperature, salinity, DO, pH and transparency) from three stations (S1, S2 and S3) along with abundance of macrofauna found during January, February and March, 2015 from the present study (Table 2). There was a very strong negative correlation between salinity and DO ( $r = -0.859, p > 0.01$ ).

A total number of 14 taxa (families) of macrofauna under 4 class identified from January to March yielded 5481 ind./m<sup>2</sup> with a mean density of  $609 \pm 445.32$  ind./m<sup>2</sup> from all stations (Table 3). The highest density of macrobenthic animal was found 919 ind./m<sup>2</sup> in station S1 during March while the lowest was 326 ind./m<sup>2</sup> at station S3 during February (Table 3). Lumbrineridae almost dominated each months in all stations contributing 49% (Figure 1) of the total macrofauna while Goniadidae and Nereididae were also abundant in all months with a mean of ( $296 \pm 139.33$  ind./m<sup>2</sup>), ( $71 \pm 42.35$  ind./m<sup>2</sup>) and ( $127 \pm 88.37$  ind./m<sup>2</sup>) with a total of (2667 ind./m<sup>2</sup>), (637 ind./m<sup>2</sup>) and (1141 ind./m<sup>2</sup>) respectively (Figure 2). In regard of the rest 11 families, their attendance were scattered in different months at different stations. Nephtyidae was found only in March while Terebridae was found in January only at station S3 (15 ind./m<sup>2</sup>) and Oedicerotidae was only at station S1 (44 ind./m<sup>2</sup>) in March. Variation of microbenthic density might be due to the seasonal variations of environmental parameters [30]. Observed

macrobenthic diversity indices were positively correlated with salinity and pH ( $p < 0.05$ ) and negatively correlated with clay and organic matter [31]. Although species level identification of microbenthic animal provides better understanding of environmental impact assessment [32], present results with the family of microbenthic individual is also acceptable [33].

Quantitative distribution of intertidal macrobenthic community structure of the Chairman Ghat has been furnished in Table 4. The fauna comprised a total 4 groups including Polychaete, Gastropoda, Bivalvia and Arthropods where Polychaete was dominant in all the stations occupying the highest percentage (Figure 3) in all three months during all the study period that yield a total (4815 ind./m<sup>2</sup>) with a mean value of ( $535 \pm 175.63$  ind./m<sup>2</sup>) comprising 87.85% of total abundance. Gastropoda was found only at station S3 (15 ind./m<sup>2</sup>) and Bivalvia at station S1 (15 ind./m<sup>2</sup>) and at S3 (30 ind./m<sup>2</sup>) in January. Arthropods were common in all month after the polychaete where polychaete was 83% and Arthropods was 7% in January, polychaete was 96% and Arthropods was 1% in February and polychaete was 84% and Arthropods was 8% in March (Figure 3). In all the comparative study of benthic group it is clear that the Polychete is the most abundant and dominant group in all three month. The density of macrobenthos group found in 3 station during 3 months of study period from Chairman Ghat are tabulated in Table 5 that shows the abundance in individual per square meter (ind./m<sup>2</sup>), their station wise percentage and the rank of abundance in 3 months. It shows that Polychete is ranked in number

Pearson Correlation (r)	Temperature	Salinity	DO	pH	Transparency	Abundance
Temperature	-	0.356	-0.37	0.3	-0.349	0.301
Salinity	-	-	-0.859**	0.4	0.398	-0.207
DO	-	-	-	0	-0.265	-0.141
pH	-	-	-	-	-0.472	-0.293
Transparency	-	-	-	-	-	-0.01
Abundance	-	-	-	-	-	-

\*\*Correlation is significant at the 0.01 level (1-tailed)

Table 2: Pearson correlation (r) among the measured parameters during the present study period.

Month	January			February			March			Mean $\pm$ SD	Total	Percentage	Rank
	JS1	JS2	JS3	FS1	FS2	FS3	MS1	MS2	MS3				
Lumbrineridae	148	207	222	563	504	193	370	222	237	$296 \pm 139.33$	2667	48.65	1
Goniadidae	59	30	15	119	148	30	59	74	104	$71 \pm 42.35$	637	11.62	3
Nephtyidae	0	0	0	0	0	0	104	30	15	$16 \pm 32.34$	148	2.7	5
Nereididae	281	193	74	74	30	0	163	104	222	$127 \pm 88.37$	1141	20.81	2
Polynoidae	30	15	0	0	0	0	0	0	0	$5 \pm 9.88$	44	0.81	12
Syllidae	0	0	0	0	0	0	0	0	30	$3 \pm 9.31$	30	0.54	13
Sabellidae	0	0	0	0	15	0	0	0	0	$2 \pm 4.66$	15	0.27	15
Capitellidae	15	0	0	30	0	44	44	0	0	$15 \pm 18.48$	133	2.43	7
Terebridae	0	0	15	0	0	0	0	0	0	$2 \pm 4.66$	15	0.27	14
Isaeidae	15	44	0	0	0	0	0	0	0	$7 \pm 14.16$	59	1.08	8
Mytilidae	15	0	30	0	0	0	0	0	0	$5 \pm 9.88$	44	0.81	9
Oedicerotidae	0	0	0	0	0	0	44	0	0	$5 \pm 13.97$	44	0.81	10
Mysidae	0	30	0	0	0	15	0	0	0	$5 \pm 9.88$	44	0.81	11
Portunidae	30	0	0	0	0	0	89	30	0	$16 \pm 28.32$	148	2.7	6
Unidentified	30	15	44	15	0	44	44	59	59	$35 \pm 19.75$	311	5.68	4
<b>Total</b>	<b>622</b>	<b>533</b>	<b>400</b>	<b>800</b>	<b>696</b>	<b>326</b>	<b>919</b>	<b>519</b>	<b>667</b>	<b><math>609 \pm 445.32</math></b>	<b>5481</b>	<b>100</b>	
<b>Family Found</b>	<b>8</b>	<b>6</b>	<b>5</b>	<b>4</b>	<b>4</b>	<b>4</b>	<b>7</b>	<b>5</b>	<b>5</b>		<b>14</b>		


JS1, JS2 and JS3 = S1, S2 and S3 in January respectively; FS1, FS2 and FS3 = S1, S2 and S3 in February respectively; MS1, MS2 and MS3 = S1, S2 and S3 in March respectively.

Table 3: Abundance of macrobenthic family (ind./m<sup>2</sup>) at all stations during the study period.

1 in all stations while other groups fluctuated within different stations in different months where sometime some groups were totally absent. Similar findings was reported near the study area [34] where Polychaeta was dominant [10,35].


### Diversity indices

Diversity Indices of different station characterized the present



study (Table 6). Different diversity indices showed significant difference between the months (Figure 4). Higher values of the diversity indices were observed in January and March while in February the values were lower than other two months. The highest value for Dominance (D) was in February (0.5438). The Simpson (1-D) value was highest (0.728) in March and lowest (0.4562) in February. The Shannon (H') diversity index is another important one. In the present investigation it ranged from 0.7797 at station S2 in February to 1.648 at station S1 in March where 1.514, 0.9202 and 1.571 was for January, February and March respectively. The Margalef's index of benthic macro-invertebrates at station S1 (0.4501) was lowest maintaining the value 0.6689 in February for in total and at station S1 (1.096) was highest maintaining the value 1.234 in January for in total while that value was 0.9246 in March. While the value of Equitability (J) index of benthic macro-invertebrates at station S2 (0.5452) was lowest maintaining the value 0.4183 in February for in total and at station S2 (0.7602) was highest maintaining the value 0.6016 in March for in total while that value was 0.4546 in January respectively.

Shannon-Weaver diversity index was defined [36] as H' and the highest and lowest value of H' indicated rich diversity with healthier environment and poor diversity with polluted environment respectively. Water bodies with macro-benthos Shannon-Wiener diversity index <1 are classified as heavily polluted, 1-3 is moderately polluted and >3 clean is environment [28,37-40]. So it can be said that all the selected stations in the study area was heavily polluted in February (H' = 0.9202) while pollution was moderate in January (H' = 1.514) and March (H' = 1.571). In one hand the suitability of habitat is reflected by the higher diversity values for the organism while it is reported that high species diversity is correlated with longer food chain, complex food web and more stable community



Month	January			February			March			Mean ± SD	Total	Rank
Station	JS1	JS2	JS3	FS1	FS2	FS3	MS1	MS2	MS3			
Polychaete	533	444	311	785	696	267	741	430	607	535 ± 175.63	4815	1
Gastropoda	0	0	15	0	0	0	0	0	0	2 ± 4.66	15	5
Bivlvia	15	0	30	0	0	0	0	0	0	5 ± 9.88	44	4
Arthropods	44	74	0	0	0	15	133	30	0	33 ± 42.92	296	3
Unidentified	30	15	44	15	0	44	44	59	59	35 ± 19.75	311	2
<b>Total</b>	<b>622</b>	<b>533</b>	<b>400</b>	<b>800</b>	<b>696</b>	<b>326</b>	<b>919</b>	<b>519</b>	<b>667</b>	<b>609 ± 176.74</b>	<b>5481</b>	

JS1, JS2 and JS3 = S1, S2 and S3 in January respectively; FS1, FS2 and FS3 = S1, S2 and S3 in February respectively; MS1, MS2 and MS3 = S1, S2 and S3 in March respectively.

**Table 4:** Group wise total abundance distribution of macrobenthos in all stations in Chairman Ghat observed during the present study.






Figure 3: Composition of macrobenthic animal (group) observed during the study period (J- in January, F- in February and M- in March).

Station	Groups	Ind./m2	Percentage (%)	Station wise Rank
JS1	Polychaete	533	85.74	1
	Gastropoda	0	0	5
	Bivalvia	15	2.38	4
	Arthropods	44	7.15	2
	Unidentified	30	4.76	3
	<b>Total</b>	<b>622</b>	<b>100</b>	
JS2	Polychaete	444	83.39	1
	Gastropoda	0	0	4
	Bivalvia	0	0	5
	Arthropods	74	13.9	2
	Unidentified	15	2.78	3
<b>Total</b>	<b>533</b>	<b>100</b>		
JS3	Polychaete	311	77.78	1
	Gastropoda	15	3.7	4
	Bivalvia	30	7.41	3
	Arthropods	0	0	5
	Unidentified	44	11.11	2
<b>Total</b>	<b>400</b>	<b>100</b>		

JS1, JS2 and JS3 = S1, S2 and S3 in January respectively.

Table 5a: Density and station wise rank of macrobenthic groups based on percentage found during January of the study period.

Station	Groups	Ind./m2	Percentage (%)	Station wise Rank
FS1	Polychaete	785	98.15	1
	Gastropoda	0	0	3
	Bivalvia	0	0	4
	Arthropods	0	0	5
	Unidentified	15	1.85	2
	<b>Total</b>	<b>800</b>	<b>100</b>	
FS2	Polychaete	696	100	1
	Gastropoda	0	0	2
	Bivalvia	0	0	3
	Arthropods	0	0	4
	Unidentified	0	0	5
<b>Total</b>	<b>696</b>	<b>100</b>		

on the other hand [41]. Depending on the number of species Margalef index (D') shows variation having no limit on tis value. So it can be postulated that the present study area is moderately

FS3	Polychaete	267	81.8	1
	Gastropoda	0	0	4
	Bivalvia	0	0	5
	Arthropods	15	4.54	3
	Unidentified	44	13.63	2
	<b>Total</b>	<b>326</b>	<b>100</b>	

FS1, FS2 and FS3 = S1, S2 and S3 in February respectively.

Table 5b: Density and station wise rank of macrobenthic groups based on percentage found during February of the study period.

Station	Groups	Ind./m2	Percentage (%)	Station wise Rank
MS1	Polychaete	741	80.6	1
	Gastropoda	0	0	4
	Bivalvia	0	0	5
	Arthropods	133	14.51	2
	Unidentified	44	4.84	3
	<b>Total</b>	<b>919</b>	<b>100</b>	
MS2	Polychaete	430	82.78	1
	Gastropoda	0	0	4
	Bivalvia	0	0	5
	Arthropods	30	5.71	3
	Unidentified	59	11.42	2
	<b>Total</b>	<b>519</b>	<b>100</b>	
MS3	Polychaete	607	91.07	1
	Gastropoda	0	0	3
	Bivalvia	0	0	4
	Arthropods	0	0	5
	Unidentified	59	8.88	2
	<b>Total</b>	<b>667</b>	<b>100</b>	

MS1, MS2 and MS3 = S1, S2 and S3 in March respectively.

Table 5c: Density and station wise rank of macrobenthic groups based on percentage found during March of the study period.




Figure 4: Monthly Values of different diversity indices of the present study.

polluted according the results calculated from Margalef index which is compatible with the findings of other research [34].

## Conclusion

14 families under 4 major groups/taxa yielded a total of 5481 ind./m<sup>2</sup> with a mean density of 609 ± 445.32 ind./m<sup>2</sup> from all stations. The Shannon-Wiener index indicate that all the stations selected was heavily polluted in February (H' = 0.9202) while pollution was moderate in January (H' = 1.514) and in March (H' = 1.571). The Margalef Species Richness Index (D') value was highest in January (1.234) and lowest in February (0.6689) where March (0.9246) maintained the medium value. Which reviles that the area falls under present study was moderately polluted.

Month	Station	Taxa (S)	Individuals	Dominance (D)	Simpson (1-D)	Shannon (H')	Margalef (D')	Equitability (J)
January	JS1	8	592.58	0.305	0.695	1.507	1.096	0.7245
	JS2	6	518.51	0.3127	0.6873	1.374	0.7999	0.7666
	JS3	5	355.54	0.4445	0.5555	1.092	0.681	0.6787
	Total	10	1466.65	0.3044	0.6956	1.514	1.234	0.6576
February	FS1	4	785.18	0.5472	0.4528	0.8703	0.4501	0.6278
	FS2	4	696.29	0.5709	0.4291	0.7797	0.4583	0.5625
	FS3	4	281.47	0.5069	0.4931	0.943	0.5319	0.6802
	Total	6	1762.95	0.5438	0.4562	0.9202	0.6689	0.5136
March	MS1	7	874.06	0.2485	0.7515	1.648	0.8859	0.8468
	MS2	5	459.25	0.3195	0.6805	1.335	0.6526	0.8296
	MS3	5	607.4	0.3183	0.6817	1.275	0.6241	0.7921
	Total	8	1940.74	0.272	0.728	1.571	0.9246	0.7557
All Station		14	5170.35	0.3327	0.6673	1.509	1.52	0.5717

JS1, JS2 and JS3 = S1, S2 and S3 in January respectively; FS1, FS2 and FS3 = S1, S2 and S3 in February respectively; MS1, MS2 and MS3 = S1, S2 and S3 in March respectively.

**Table 6:** Different diversity Indices at different stations observed during the study period.

## References

- Smith RW, Weisberg SB, Cadien D, Dalkey A, Montagne D, et al. (2001) Benthic response index for assessing infaunal communities on the southern California Coastal Shelf. *Ecological Applications* 11: 1073-1087.
- Snelgrove PVR (1998) The biodiversity of macrofaunal organisms in marine sedi-ments. *Biodiversity and Conservation* 7: 1123-1132.
- Rhoads DC (1974) Organism-sediment relations on the muddy sea floor. *Oceanography and Marine Biology Annual Review* 12: 263-300.
- Fauchald K, Jumars PA (1979) The diet of worms: a study of polychaete feeding guilds. *Oceanography and Marine Biology Annual Review* 17: 193-284.
- Weisberg SB, Ranasighe JA, Schaffner LC, Diaz RJ, Dauer DM, et al. (1997) An estuarine index of biological integrity (B-IBI) for Chesapeake Bay. *Estuaries* 20: 149.
- Little C (2000) *The Biology of Soft Shores and Estuaries*. Oxford University Press, New York, NY: 158.
- Pearson TH, Rosenberg R (1978) Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. *Oceanography and Marine Biology Annual Review* 16: 229-311.
- Hart CW, Fuller SLH (1979) *Pollution Ecology of Estuarine Invertebrates*. Academic Press, New York: 406.
- Paul JF, Scott KJ, Campbell DE, Gentile JH, Strobel CS (2001) Developing and applying a benthic index of estuarine condition for the Virginian Biogeographic Province. *Ecological Indicators* 1: 83-99.
- Asadujaman M, Belal Hossain M, Shamsuddin M, Amin MA, Azam AKM (2012) Occurrence and Abundance of Macrobenthos of Hatiya and Nijhum Dweep Islands, Bangladesh. *Middle-East Journal of Scientific Research* 11: 184-188.
- Gray JS, Elloitt M (2009) *The Ecology of Marine Sediments (2ndedn)*. The Oxford University Press, Oxford, UK: 256.
- Lind OT (1979) *Handbook of common method in limnology (2ndedn)*. The C.V. Mosby Company, St. Louis: 136-145.
- Newrkla P, Gunatilaka A (1982) Benthic community metabolism of three Austrian pre-alpine lakes of different tropic conditions and its oxygen dependency. *Hydrobiologia* 92: 531-536.
- Woodin SA (1974) Polychaete abundance patterns in a marine soft-sediment environment: the importance of biological interactions. *Ecological Monographs* 44: 171-187.
- Peterson CH, Andre SV (1980) An experimental analysis of interespecific competition among marine filter feeders in a soft-sediment environment. *Ecology* 61: 129-139.
- Brenchley GA (1982) Mechanisms of spatial competition in marine soft-bottom communities. *Journal of Experimental Marine Biology and Ecology* 60: 17-33.
- Thrush SF, Pridmore RD, Hewitt JE, Cummins VJ (1992) Adult infauna as facilitators of colonization on intertidal sandflats. *Journal of Experimental Marine Biology and Ecology* 159: 253-265.
- Knox GA (2000) *The ecology of sea shores*. CRC Press, New York: 555.
- Day JW, Hall CAS, Kemp WM, Yañez-Arancibia A (1989) The estuarine bottom and benthic subsystem. *Estuarine Ecology*. Wiley & Sons, New York: 338-376.
- Souza JBR, Gianuca NM (1995) Zonation and seasonal variation of the intertidal macrofauna on a sandy beach of Parana State, Brazil. *Scientia Marina* 59: 103-111.
- Veloso VG, Cardoso RS, Fonseca DB (1997) Spatio-temporal characterization of intertidal macrofauna at Prainha beach (Rio de Janeiro State). *Oecologia Brasiliensis* 435: 213-225.
- Das Neves LP, da Silva P de SR, Bemvenuti CE (2008) Temporal variability of benthic macrofauna on Cassino beach, southernmost Brazil. *Iheringia, Serie Zoologica* 98: 36-44.
- Pearson TH (1970) The benthic ecology of Loch Linnhe and Loch Eil, a Sea-Loch system on the west coast of Scotland .The physical environment and distribution of the macrobenthic fauna. *J Exp Mar Biol Ecol* 5: 1-34.
- Ryan PD, Harper DAT, Whalley JS (1995) *Palstat, Statistics for palaeontologists*. Chapman & Hall (now Kluwer Academic Publishers).
- Welch PS (1948) *Limnology II edition*. Mc graw Hill book Company, New York.
- (1999) In: Harper DAT (ed.), *Numerical Palaeobiology*. John Wiley & Sons.
- Simpson EH (1949) Measurement of diversity. *Nature* 163: 688.
- Wilhm JL, Dorris TC (1966) Species diversity of benthic macroinvertebrates in a stream receiving domestic and oil refinery effluents. *Am. Midland Nat* 76: 427-449.
- Margalef R (1968) *Perspectives in Ecological Theory*. University of Chicago Press, Chicago, IL: 111.
- Islam Shafiqul M, Sikdar Nurul Azim M, Al-Imran M, Hossain Belal M, Mallick Debbroto (2013) Intertidal Macrobenthic Fauna of the Karnafuli Estuary: Relations with Environmental Variables, *World Applied Sciences Journal* 21: 1366-1373.
- Hossain Belal M, Marshall David J (2014) Benthic infaunal community structuring in an acidified tropical estuarine system. *Aquatic Biosystems* 10: 11.
- Narayanaswamy BE, Nickell TD, Gage JD (2003) Appropriate levels of taxonomic discrimination in deep-sea studies: species vs family. *Marine ecology. Progress series* 257: 59-68.
- Warwick RM (1988) Analysis of community attributes of the macrobenthos of Frierfjord/ Langesundfjord at taxonomic levels higher than species. *Marine Ecology Progress Series* 46: 167-170.
- Sarker JM, Patwary SAM, Uddin AMMB, Hasan M Md, Tanmay MH, et al. (2016) Macrobenthic Community Structure - An Approach to Assess Coastal Water Pollution in Bangladesh. *Fish Aquac J* 7: 157.
- Hossain Belal M, Amin AMN, Asadujaman M, Rahman S (2013) Analyses of Macrobenthos of Hatiya and Nijhum Dweep Islands at Higher Taxonomic Resolution. *Journal of Fisheries and Aquatic Science* 8: 526-534.

36. Wilhm JL (1975) Biological indicators of pollution, In: River ecology, Whitton BA (ed.), Studies in Ecology, Blackwell Sci Publ, London 2: 375-402.
37. Islam MS (2008) Phytoplanktonic diversity index with reference to Mucalinda Serovar, Bodh-Gaya. Proceedings of the TaaI' 2007: The 12th World Lack Conference: 462-463.
38. Tanimu Y, Bako SP, Adakole JA, Tanimu J (2011) Phytoplankton as bioindicators of water quality in saminaka reservoir, Northern Nigeria. Proceedings of the International Symposium on Environmental Science and Technology, Dongguan, Guangdong Province, China: 318-322.
39. Stub R, Appling JW, Hatstetter AM, Hass IJ (1970) The effect of industrial waste of Memphis and Shelby country on primary planktonic producers. Bioscience 20: 905-912.
40. Mason CF (1988) Biology of Fresh Water Pollution. Longman scientific and technical.
41. Margalef R (1956) Information Y diversidad especiifi caenlas comunidades de organismos. Invest Pesg 3: 99-106.

**Citation:** Sarker J, Tanmay MH, Rahman F, Patwary MSA, Rima NN (2016) Assessment of Coastal Water Pollution In Greater Noakhali-Bangladesh. J Coast Zone Manag 19: 427. doi:[10.4172/2473-3350.1000427](https://doi.org/10.4172/2473-3350.1000427)

#### OMICS International: Publication Benefits & Features

##### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

##### Special features:

- 700+ Open Access Journals
- 50,000+ Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: [www.omicsonline.org/submission/](http://www.omicsonline.org/submission/)