

Major Health Challenges of Dairy Cattle in Hawassa Town SNNPRS, Ethiopia

Nigussu Fasil^{2*}, Terefe Simon Jutta¹ and Dessie Sheferaw²

¹Department of Parasitology and Pathology, Jigjiga University, Jigjiga, Ethiopia

²School of Veterinary Medicine, Hawassa University, Ethiopia

Abstract

The aim of this study was to assess major health challenges of dairy cattle in Hawassa town, Ethiopia, which occurred on November 2014 up to April 2015. A single visit-multiple subject formal survey technique was used to collect data from 20 dairy farming which were selected at random and were interviewed using pre-tested, structured questionnaire which indicated that over all prevalence in this study was 33.6% (n=269) cattle were found affected by either one or more of health challenges. It appeared from the study that LSD (30.1%), mastitis (20.4%), hypocalcaemia (17.5%), repeated breeding (13%), RFM (10%) and parturient paresis (6.7%) were identified as the most frequently occurring diseases. Results of the major dairy cattle disease in the study area ranked LSD as number one disease occurred in different dairy farms, followed by mastitis (20.4%) and hypocalcaemia (17.5%). The degree of association of risk factors was assessed and parity, age, farm scale, and management system found to be directly associated. As the result shows age with dystocia directly associated and have good significances with the $\chi^2=12.479$ and p-value =0.002, and LSD associated with farm scale but not significant, with the $\chi^2=4.705$ and (p>0.05) and also abortion associated with management system but have no significance (p>0.05). This particular study indicated major health challenges which included hypocalcaemia, ketosis, abortion, RFM, parturient paresis, repeated breeding, diarrhea, bloat, and anestrus, uterine prolapsed, vaginal prolapsed, dystocia and LSD were one of the major reproductive and metabolic disorders responsible for the low reproductive performance of dairy cows.

Keywords: Age; Breed; Cattle; Dairy farm; Hawassa; Health

Abbreviations: DVM: Doctor of Veterinary Medicine; P-Value: Expected Prevalence; RFM: Retained Fetal Membrane; SPSS: Stastical Package for Social Science; LSD: Lumpy Skin Disease; χ^2 : Chi-square test.

Introduction

Ethiopia has the largest livestock population being the first in Africa countries and the 10th in the world and holds large potential for dairy development due to its large livestock population and Urban and peri-urban livestock production constitutes an important sub-sector of the agricultural production system [1]. In Ethiopia, livestock represents a major national resource and form an integral part of the agricultural production system [2]. The livestock sector in general and the dairy industry in particular do not provide the expected contribution to the national income despite their large numbers due to several factors. The development of the dairy sector in the country is hindered by a number of technical, institutional and socio-economic constraints. The growth in milk production has been slow and the annual milk production is estimated to be 1,089,488,251 liters [3] which doesn't meet even the domestic demand for dairy products. As a result the country imports large volumes of dairy products per annum to meet the domestic demand. In 2005, for instance, the country imported 457,260 kg of milk (liquid and powder) which is equivalent to 3,026,724 Birr [3].

Despite the huge number of cattle and their dairy industry the productivity is low due to the constraints of disease, scarcity of feed, inefficient and insufficient AI, veterinary services nutrition, poor management, lack of marketing facilities and opportunity, inadequate animal health services, uncoordinated development programs between various levels of government institutions and /or non-government organizations and poor performance of indigenous breeds. These constraints result in health challenges of dairy cattle [4].

Major health challenges of dairy cattle were consists of ketosis, hypocalcaemia, metritis, retained fetal membranes, LSD, bloating, mastitis and uterine prolapsed. All of these diseases are related to

one another, with complicated cause and effect mechanisms in place. Numerous studies [5-10] had shown that postpartum diseases can affect the length of calving interval, the number of days open, and the reproductive efficiency in general. These diseases can also affect the overall productivity of dairy cows by reducing milk yield. Studies conducted so far in Ethiopia [11-15] revealed poor reproductive performance of dairy cows in the tropics. For feasible intervention, the poor reproductive performance of dairy cows should warrant investigation on the types and magnitudes of the existing postpartum problems. These constraints also result in poor reproductive performance of dairy cattle. Among the major problems that have a direct impact on reproductive performance of dairy cows are: abortion, retention of the fetal membrane (RFM) and metritis. These results in considerable economic losses to the dairy industry due to slower uterine involution, reduced reproductive rate, prolonged inter conception and calving interval, cost of medication, drop in milk production, reduced calf drop, and early depreciation of potentially useful cows [16,17].

Although, the major health problems are greatly responsible for high economic loss in the dairy industry, there is scarcity of reliable information regarding the reproductive performances of dairy cows in subsistence dairy farms in the tropics, particularly in Ethiopia. Information pertaining to reproductive performance and interacting factors is of paramount importance primarily to the livestock owners and also to the extension agents, veterinarians and researchers. Moreover, it

***Corresponding author:** Nigussu Fasil, Assistant Professor, School of Veterinary Medicine, Hawassa University, Ethiopia, Tel: +251913304082; E-mail: fasilkedeseagne@yahoo.com

Received August 26, 2015; **Accepted** July 28, 2016; **Published** August 02, 2016

Citation: Fasil N, Jutta TS (2016) Major Health Challenges of Dairy Cattle in Hawassa Town SNNPRS, Ethiopia. J Vet Sci Technol 7: 367. doi:10.4172/2157-7579.1000367

Copyright: © 2016 Fasil N, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

can assist in the development of strategies and prioritization of possible intervention options for performance improvement [18]. Dairy cattle require minerals in their diet for optimal productivity. These are derived from the feed and fodder. The input of minerals through feed and water must balance their output through feces, urine and milk to maintain the animal's health. If the output exceeds input, the animals meet out their normal requirements by mobilization from its body reserves for a shorter period. But continuous imbalances develop into productivity related problems.

Nutritional imbalances, deficiencies, or erratic management of feeding programs for dairy cows can create large numbers and various types of health problems generally categorized as metabolic diseases. Most periparturient abnormalities have some metabolic element as a component of the cause of clinical disease. The metabolic disturbance of milk fever can be measured through low serum calcium concentrations. Negative energy balance, fat mobilization and subsequent elevations in ketone body concentrations play a contributing role in the expression of fatty liver syndrome, clinical ketosis, and abomasal displacement. A negative energy balance may also increase the risk of retained placenta, metritis, and mastitis through impaired immune function.

Therefore, the objectives of present study were;

- To identify major health challenges of dairy cattle found in different dairy farms.

- To assess risk factors that cause to occurrence of health challenges in dairy cow.

Materials and Methods

Description of the study area

The cross sectional study was conducted from November, 2014 up to April, 2015, in Hawassa, capital city of the Southern Nations Nationalities and Peoples Regional State (SNNPRS), which is one of the high potential areas for milk production in Southern Ethiopia. It is located 275 km south of Addis Ababa along the Addis Ababa - Moyale highway. Hawassa is situated at an altitude of 1750 m above sea level and according to an estimate, it lies between 6°83' to 7°17' N and 38°24' to 38°72' E. Hawassa receives an average annual rainfall of 955 mm with mean annual temperature of 20°C [3] (Figure 1).

Study design and method of sampling

The cross-sectional study design was needed to determine the prevalence of hypocalcaemia, parturient paresis, ketosis, bloat, mastitis, uterine prolapsed, vaginal prolapsed, dystocia, abortion, anestrus, diarrhoea, pyometra, retained fetal membrane, LSD and its risk factors that predisposes to this major health challenges. The study was a questionnaire data collection and analysis to establish the prevalence and to identify the major health challenges in the selected




Figure 1: The regional map of SNNPRS.

farms. Farms purposively selected based their accessibility number of dairy populations in the area. Animals were selected by using simple random methods.

Study animals

The exotic, local and cross breeds of cows were used in this study. Addition to this extensively managed cross breeds that high grade Holstein Fresian, predominantly Holstein-indigenous zebu cross breeds and indigenous local zebu lactating cows in dairy farms were taken to this study. The history of the animals like age, parity, farm scale, stage of lactation, lactation number, milk yield, body condition, nutritional condition, symptoms observed by the owner, past and present history regarding other illness, and have different production system (extensive, semi-intensive and intensive type of management) were recorded in selected farms in Hawassa town.

Data collection

Questionnaires survey method: Structured questionnaire was prepared and used to collect information from 20 dairy farm owners by regular farm visit interview and recorded lists about major health challenges of their dairy cattle on individual level were studied. The questionnaires were checked for clarity of the questions prior the interview. Prior the interview, respondents were briefed to the objective of the study. Following that, the actual questions and questionnaires were presented. Accordingly, information about the parity, breed, feeding system, production system, and type of feed, health care and major health challenges such as hypocalcaemia, ketosis, abortion, mastitis, parturient paresis, anoestrus, uterine and vaginal prolapse, bloat, repeat breeding and LSD were collected on individual cattle level. From 20 selected dairy farms five farms have well recorded data list was needed to this study. In order to identify dairy farms considered in the current study, an initial list of dairy farms in the city was obtained from Hawassa City Administration Agricultural and Rural Development Office.

Data analysis

The data was presented using the descriptive statistics, the retrospective data results was entered to a Microsoft Excel sheet 2007 and analyzed using a software SPSS' version 20. Different factors include age, breed, productivity, parity, management, farm scale and body condition score that were considered during the study period were analyzed using the Chi-square technique. The possible association of dystocia, abortion, and LSD with age, management, and farm scale respectively was also tested using these techniques. In all chi-square test application, probability of $p < 0.05$ was considered statistically significant.

Results

In the cross sectional study of questionnaire LSD, mastitis, hypocalcaemia, and repeated breeding were found to be the major health challenges occur in Hawassa town dairy farm, containing (30.1%), (20.4%), (17.5%) and (13%), respectively. Other health problems observed with lower prevalence include retain fetal membrane (10%), parturient paresis (6.7%), diarrhea (6.3%), dystocia (5.6%), anestrous (5.6%), ketosis (5.2%), abortion (5.2%), bloat (4.5%), payometra (3.7%), uterine prolapse (3.7%) and vaginal prolapse (2.2%) accounting were obtained from study.

From the total of 269 dairy cattle, 129 (47.9%) were managed semi-intensive, 38 (14.1%) were intensively and 102 (37.9%) were extensively and of which 250 (92.9%) were cross breed and the rest 19 (7.1%) were local breed. Most of the animal were adult age group encountered, 222

(82.5%), the rest of them were 23 (8.6%) young and 24 (8.9%) old age group. The parity of the animal is few, moderate, and high, which was in number showed 194 (72.1%), 62 (23.0%) and 13 (4.8%) respectively.

Major health challenges identified in dairy farms

The major health challenges identified were LSD (30.1%), mastitis (20.4%) and hypocalcaemia (17.5%), as summarized on Table 1, bellow. In this study 33.6% (n=269) cattle were found affected by either one or more of health challenges.

Association of risk factors with health challenges of dairy cattle

In this study among risk factors age, farm scale, parity and management system were considered to assess its association with the occurrence of the health challenges as shown on the tables below (Table 2).

Table 3 showed that there is statistically significant ($p < 0.05$) variation with regard to age and dystocia. Highest prevalence was found in cattle with age group adult and followed by age group young while the least in cows with age group old. The prevalence rate of LSD at farm scale level is higher in small farm scale (46.1%) than in medium (26.3%) and large (33.3%) farm scale.

As shown on the Tables 4 and 5 above, statistically no significant difference ($P > 0.05$) was found in the prevalence of health challenges with respect to management system.

Discussion

In the present study (33.6%) of dairy cattle in the study areas were affected by either one or major health challenges based on questionnaires to the owners. Major health challenges are LSD, mastitis, hypocalcaemia, and repeated breeding were found to be the major health challenges occur in Hawassa town dairy farm, containing (30.1%), (20.4%), (17.5%) and (13%), respectively. Recent survey which assesses the risk factors and financial impacts of LSD in selected districts of North-eastern Ethiopia (Tigray and Afar Regional States) conducted by Birhanu [19] reported a higher herd prevalence of (51%) and (37%) was recorded in Afar and Tigray Region respectively which is higher result from present study (30.1%), this variation might be due to environment, breed of animal and management system. The prevalence rate of mastitis (20.4%) recorded in this study is higher than to the reports done in different dairy farms in Hawassa town (4.9%) by

Challenges	Frequency	Percentage (%)
Hypocalcaemia	47	17.5
Payometra	10	3.7
Parturient paresis	18	6.7
Diahorrea	17	6.3
Ketosis	14	5.2
Bloat	12	4.5
Mastitis	55	20.4
Dystocia	15	5.6
Abortion	14	5.2
RFM	27	10
Uterine prolapsed	10	3.7
Vaginal prolapsed	6	2.2
Anestrous	15	5.6
Repeated breeding	35	13
LSD	81	30.1

Table 1: Frequency and Prevalence rate of major health problems of dairy Cattle in Hawassa town.

Risk factors	No. of cattle	Percentage
Age		
Young (<4 years)	23	8.6
Adult (4-8 years)	222	82.5
Old (>9 years)	24	8.9
Parity		
Few (<3 birth)	194	72.1
Moderate (3-5 birth)	62	23.0
High (>5 birth)	13	4.8
Productivity		
No	1	4
Low	157	58.4
Medium	94	34.9
High	17	6.3
Breed		
Local	19	7.1
Cross	250	92.9
Body condition		
Poor	78	29.0
Medium	138	51.3
Good	53	19.7
Management system		
Extensive	103	38.3
Semi-intensive	128	47.6
Intensive	38	14.1
Farm scale		
Small	26	9.7
Medium	171	63.6
Large	72	26.8

Table 2: Association of risk factors with health challenges of dairy cattle.

Challenge	Age				χ ²	P- value
	Young	adult	Old	Total		
Dystocia						
-ve	18	213	23	254	12.479	0.002
+ve	5	9	1	15		
Total	23	222	24	269		

Table 3: Association of age group with health challenge.

Challenges	Farm scale				χ ²	P- value
	Small	Medium	Large	Total		
LSD						
-ve	14	126	48	188	4.705	0.095
+ve	12	45	24	81		
Total	26	171	72	269		

Table 4: Association of farm scale with health challenge.

Challenges	Management system				χ ²	P-value
	Extensive	Semi-intensive	intensive	Total		
Abortion						
-ve	99	122	34	255	2.615	0.270
+ve	4	6	4	14		
Total	103	128	38	269		

Table 5: Association of management system with health challenge.

Nibret et al. [20]; in and around Mekelle (6.55%) by Wudu [21] and in central high lands of Ethiopia (6.6%) by Mungube et al. [22]. While it was comparable in the reports done in and around Sebeta (16.1%) by Hunderra et al. [23]; in Dire Dawa Administrative Council and Eastern Hararghe Zone (19.8%) by Birhanu [24]; in two major state owned dairy farms at Rapi and Debre Zeit, Ethiopia (21%) by Workineh et al. [25]; but it was lower in the research finding in selected areas of southern Ethiopia (37%) by Kerro and Tareke [26].

The prevalence rate of hypocalcaemia (17.5%) recorded in this study is lower than the result (30.2%) reported by Samuel et al. [27] this difference may be management, type of breed and study population. Most of the literatures suggest that when the incidence of milk fevers increases above (10%) in their third or latter lactation, considerations

should be given to a specific control program [28]. Therefore, these results indicated that control methods are required to avoid loss due to milk fever. Milk fever is caused by a severe deficiency of metabolizable calcium ion in the circulation. This could be attributed due to several risk factors [29]. The risk factors identified in this study include milk yield, parity and breed of the cows.

The higher prevalence of repeated breeding (13%) found in the present study is in close agreement with (13%) reported by Micheal [30] in and around Hawassa and (13.08%) by Adane et al. [31] but higher than (11.4%) reported by Hadush et al. [32] from central Ethiopia. Repeated breeding can be caused by a number of factors, including sub-fertile bulls, endocrine imbalance, malnutrition, reproductive tract infections and poor management practices such as wrong time of insemination or faulty heat detection, inappropriate semen handling and insemination techniques [33]. In addition to these, communal use of bull for natural services also considered as contributing factor. Hence the difference between the findings of the current study and previous reports may be attributed to the above-mentioned factors.

Other health problems observed with lower prevalence include retain fetal membrane (10%), parturient paresis (6.7%), diarrhea (6.3%), dystocia (5.6%), anestrus (5.6%), ketosis (5.2%), abortion (5.2%), bloat (4.5%), payometra (3.7%), uterine prolapse (3.7%) and vaginal prolapse (2.2%) were obtained.

The prevalence rate of RFM (10%) in recent study is similar with the (8.6%) reported by Molalegn and Shiv [34] and lower than (14.2%) reported by Mamo [35] and 19.2% by Gashaw et al. [36]. The variation in the incidence of RFM attributed to variations in predisposing factors to which the animals are subjected to nutritional status and management problems such as lack of exercise. Dystocia that accounted (5.9%) of the problems is an important predisposing factor for occurrence of RFM. Previous report (5.79%) on the prevalence of dystocia by Mamo [35] in small holder dairy cows in and around Debre Zeiet fairly agrees to the prevalence of (5.6%) obtained in this study. However, the current finding is lower than the prevalence of (7.7%) reported by Dawit and Ahmed [37], and higher than those (3.8%) of Gashaw et al. [36]. This variation in the occurrence of dystocia due to the fact that it is influenced by the factors such as, age and parity of the dam as well as breed of the sire. Inseminating cows with semen collected from large sized bulls without taking into account the size and age of cows is an important factor in precipitating dystocia [38].

The prevalence of anestrus observed in this study (5.6%) was lower than the results of Hadush et al. [32], who reported (12.9%) in dairy cattle in Debre Zeit and Befekadu [39], who reported (24%) in cross breed dairy cows in central high lands of Ethiopia. This variation might be due to the age, faulty heat detection, and breed and management system differences.

The prevalence rate of abortion (5.2%) recorded in this study was higher than the result (2.2%) reported by Bekana et al. [34] in Nazret, but is lower than the (9.0%), (13.9%) and (14.6%) reported by Dawite and Ahmed [37], Molalegn and Shiv [34] and Hunduma [40] respectively. The lower prevalence rate of abortion may be attributed to the increasing practice of AI in the study area where the semen is collected from bulls free from brucellosis, in addition breed, management system specially feeding and sanitation. Study methodology and geographical location differences are all sources of differences in prevalence of abortion [36].

The prevalence rate of vaginal prolapse (2.2%) recorded in this study is lower than the result (5.2%) reported by Kidusan [41] but is higher than the (1.2%) reported by Dawite and Ahmed [37]. This

variation might be due to management system (feeding), abortion, RFM, dystocia and breed of animals.

The prevalence rate of ketosis (5.2%) recorded in this study is lower than the result (11.2%) reported by Mulat et al. [42] in and around Addis Ababa. The highest prevalence of ketosis was recorded in the month of January. This indicates that ketosis occurred mostly during the winter season as the animals are usually housed and there is scarce of pasture [43].

There was both reproductive and metabolic problems (49%) and (29.4%) respectively. This result indicates that it is higher value than the prevalence (43.3%) reported by Adane et al. [31] in urban and per urban area of Hossana, (44.3%) by Hadush et al. [32] in central Ethiopia and (40.3%) by Dawite and Ahmed [37] in Northeast Ethiopia. This difference might be due to sample size, production system, study methodology and breed of animals as well as environmental factors.

The higher prevalence rate of health challenges in cross breed animals 92.9% (n=250) than local breed 7.1% (n=19) may be due to the fact that cross breeds are less adapted to tropical conditions of high temperature and humidity, disease and low feed quality than zebu cattle [44-46] making them more susceptible than indigenous zebu. In addition to this the risk factors like age group, parity, management system; farm scale, body condition and productivity are great effect up on major health challenges of dairy cattle.

Conclusion and Recommendation

The result found in the current study was an agreement with the other studies those metabolic and reproductive problems which have adverse effects on the health, production and reproduction indices of the dairy cow. Hypocalcaemia, ketosis, bloat, mastitis, uterine and vaginal prolapse, LSD, abortion, dystocia, RFM, anestrus and diarrhoea were major health challenge in the dairy farm. Age, breed, parity, and management systems were the most important predisposing risk factor for the various health challenges. Development of practical management strategies to cope with the negative effects associated with reproductive and health problems on dairy cattle is critical in this study area. Therefore, the following recommendation will be forwarded:

- Providing an adequate amount of a properly formulated and delivered ration,
- Providing a clean, comfortable and minimal-stress environment.
- There should be proper animal management, cleanliness and good hygiene on dairy farms.
- Improving veterinary services with respect to adequate vaccination.
- There should be Routine and periodical examination of cows during postpartum period.
- Screening, sanitation, serious follow up and health care are very important.

Acknowledgements

To begin with, I would like to express my thanks to my academic advisor, Dr. Nigussu Fasil, who had made professional advices and critical comments, without which my study would have not reached to this stage of completion. My thanks also extend to my co-advisors, Dr. Desse Shiferaw. They all contributed invaluable advices and comments to my study from the step of inception up to the completion of this DVM, Thesis research work. My family, particularly my dad Simon Juta, wishes long life, additionally my golden sisters Minitwab Simon, Tsedal Simon and my diamond brother Ababayehu Simon, I'm very much grateful to your ever

supports to all my educational works. Prior to all, "GOD IS GREAT!!" is my ever saying to all great deeds he did to me.

References

1. Tegegne A, Tadesse M, Yami A, Mekasha Y (2000) Market-oriented urban and peri-urban dairy systems. *Urban Agriculture Magazine* 1: 23-24.
2. IFPRI (2006) Atlas of the Ethiopian Rural Economy. International Food Policy Research Institute (Washington, DC), Central Statistical Agency (Addis Ababa), Ethiopian Development Research Institute (Addis Ababa).
3. BoFED (2007) Bureau of Finance and Economic Development, Southern Nations, Nationalities and People's Regional State (SNNPR).
4. Drug Administration and Control Authority (2006) Standard veterinary treatment guide lines for Veterinary Clinics. 1st edition. Pp: 3-13.
5. Borsberry S, Dobson H (1989) Periparturient diseases and their effect on reproductive performance in five dairy herds. *Vet Rec* 138: 217-219.
6. Gröhn Y, McCulloch C, Salonien H (1990) Epidemiology of mammary gland disorders in multiparous Finnish Ayrshire cows. *Prev Vet Med* 8: 241-252.
7. Rajala PJ, Gröhn YT (1998) Effects of dystocia, retained placenta, and metritis on milk yield in dairy cows. *J Dairy Sci* 81: 3172-3181.
8. Ferguson J, Galligan D (2000) Assessment of Reproductive Efficiency in Dairy Herds. *Educ Proc Vet* 22: 150-158.
9. LeBlanc SJ, Duffield TF, Leslie KE, Bateman KG, Keefe GP, et al. (2002) Defining and diagnosing postpartum clinical endometritis and its impact on reproductive performance in dairy cows. *J Dairy Sci* 85: 2223-2236.
10. LeBlanc SJ (2008) Postpartum uterine disease and dairy herd reproductive performance: a review. *Vet J* 176: 102-114.
11. Negussie E, Brannang E, Banjaw K, Rottmann O (1998) Reproductive performance of dairy cattle at Asella livestock farm, Arsi, Ethiopia. I: Indigenous cows versus their F1 crosses. *J Anim Breed Genet* 115: 267-280.
12. Shiferaw Y, Tenhagen BA, Bekana M, Kassa T (2003) Reproductive Disorders of Crossbred Dairy Cows in the Central Highlands of Ethiopia and Their Effect on Reproductive Performance. *Trop Anim Health Prod* 37: 427-441.
13. Asseged B, Birhanu M (2004) Survival analysis of calves and reproductive performance of cows in commercial dairy farms in and around Addis Ababa, Ethiopia. *Trop Anim Health Prod* 36: 663-672.
14. Ibrahim N, Abraha A, Mulugeta S (2011) Assessment of reproductive performances of crossbred dairy cattle (Holstein Friesian× Zebu) in Gondar town. *Global Vet* 6: 561-566.
15. Dinka H (2012) Reproductive performance of crossbred dairy cows under smallholder condition in Ethiopia. *Int J Livestock Prod* 3: 25-28.
16. Thomas C (1989) A preliminary survey on infertility aspects of selected dairy farms around Addis Ababa. A seminar presented to the orientation course in Tropical Animal and Health Production, Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia.
17. Barnouin J, Chacronac J (1992) A nutritional risk factor for early metritis in dairy farms in France. *Pre Vet Med* 13: 23-27.
18. Angesom H, Adem A, Fikru R (2013) Major prepartum and postpartum reproductive problems of dairy cattle in Central Ethiopia. *J Vet Med Anim Health* 5: 118-123.
19. Birhanu T (2012) Assessments of the risk factors and financial impacts of LSD in selected districts of Tigray and Afar Regional States, Northeastern Ethiopia. M.Sc. Thesis.
20. Moges N, Hailemariam T, Fentahun T, Chanie M, Melaku A (2012) Bovine Mastitis and Associated Risk Factors in Small Holder Lactating Dairy Farms in Hawassa, Southern Ethiopia. *Global Vet* 9: 441-446.
21. Wudu T (1999) Study on Bovine mastitis in and around Mekelle. DVM thesis, Faculty of Veterinary Medicine, Addis Ababa University, Ethiopia.
22. Mungube EO, Tenhagen BA, Regassa F, Kyule MN, Shiferaw Y, et al. (2005) Reduced milk production in udder quarters with subclinical mastitis and associated economic losses in crossbred dairy cows in Ethiopia. *Trop Anim Health Prod* 37: 503-512.
23. Sori H, Zerihun A, Abdicho S (2005) Dairy cattle mastitis in and around Sebeta, Ethiopia. *Intern J Appl Res Vet Med* 3: 332-338.

24. Birhanu S (1997) Bovine Mastitis in Dairy Farm in Dire Dawa Administrative Council and Western Harareghe Zone. Prevalence, isolation and *in vitro* antimicrobial susceptibility study. DVM thesis. Addis Ababa University, Faculty of Veterinary medicine, Ethiopia.
25. Workneh S, Bayleyegn M, Mekonnen H, Potgieter LN (2002) Prevalence and aetiology of mastitis in cows from two major Ethiopian dairies. Trop Anim Health Prod 34: 19-25.
26. Dego OK, Tareke F (2003) Bovine mastitis in selected areas of southern Ethiopia. Trop Anim Health Prod 35: 197-205.
27. Anteneh S, Guadu T, Fentahun T, Chanie M (2012) Incidence of Milk Fever on Dairy Cows and its Risk Factors in Gondar Town, Northwest Ethiopia. Int J Nat Sci 9: 659-662.
28. Horst RL (1986) Regulation of calcium and phosphorus homeostasis in the dairy cow. J Dairy Sci 69: 604-616.
29. Brandly CA, Cornelius CE (1969) Parturient Hypocalcaemia in Dairy Cows. Advances in Veterinary Science and Comparative Medicine. Academic press, USA, 15: 143-150.
30. Micheal K (2003) Major clinical reproductive problem of small holder dairy cows in and around Awassa. DVM Thesis, Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia.
31. Adane H, Yisehak T, Niguse T (2014) Assessment of major reproductive disorders of dairy cattle in urban and per urban area of Hosanna, Southern Ethiopia. Animal Vet Sci 2: 135-141.
32. Hadush A, Abdella A, Regassa F (2013) Major prepartum and postpartum reproductive problems of dairy cattle in Central Ethiopia. J Vet Med Anim Health 5: 118-123.
33. Arthur G, Noakes D, Pearson H (1989) Veterinary Reproduction and Obstetrics. Theriogenology 6th edition. Baillier Tindall UK, Pp: 83-85.
34. Molalegne B, Shiv P (2011) Study on Major Reproductive Health Problems in Indigenous and Cross Breed Cows in and Around Bedelle, South West Ethiopia. J Anim Vet Adv 10: 723-727.
35. Mamo T (2004) Study on major postpartum reproductive problems of smallholder dairy cows in and around Debre Zeit. DVM Thesis, Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia.
36. Gashaw A, Worku F, Mulugeta S (2011) Assessment of small holder dairy production system and their reproductive health problems in Jimma town Southwestern Ethiopia. Intern J Appl Res Vet Med 9: 80-86.
37. Tesfaye D, Shamble A (2013) Reproductive health problems of cows under different management systems in kombolcha, North east Ethiopia. Adv Biol Res 7: 104-108.
38. Noakes D (1986) Fertility and Obstetrics in cattle. Blackwell Science Publishers, Oxford, UK, Pp: 28-30.
39. Befekadu U (2007) A field investigation of delayed resumption of postpartum ovarian activity and prolonged postpartum anestrus in cross bred dairy cows in Debre Zeit dairy herds (Doctoral dissertation, MSc Thesis, Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia).
40. Dinka H (2013) Major reproductive disorders of dairy cows in and around Asella town, Central Ethiopia. J Vet Med Anim Health 5: 113-117.
41. Kidusan K (2009) Study on major reproductive disorders of dairy cattle in and around Wukro. DVM Thesis, College of Veterinary Medicine, Mekelle University, Tigray, Northern Ethiopia.
42. Asrat M, Tadesse GH, Gounder RV, Nagappan R (2013) Prevalence and Treatment of Ketosis in Dairy Cows in and Around Addis Ababa, Ethiopia. Br J Dairy Sci 3: 26-30.
43. Blood D, Radostists O, Henderson J (1983) Veterinary Medicine. 6th edition. Bailliere Tindall, London and New York.
44. Mugerwa ME (1989) A review of reproductive performance of female Bos indicus (Zebu) cattle. ILCA monograph No. 6. ILCA, Addis Ababa.
45. Plan LM (2008) Livestock development master plan study-Phase 1 report: Data collection and analysis. Volume H-Animal Nutrition (November 2008).
46. Stevenson JS, Call EP (1988) Reproductive disorders in the periparturient dairy cow. J Dairy Sci 71: 2572-2583.

Citation: Fasil N, Juta TS (2016) Major Health Challenges of Dairy Cattle in Hawassa Town SNNPRS, Ethiopia. J Vet Sci Technol 7: 367. doi:[10.4172/2157-7579.1000367](https://doi.org/10.4172/2157-7579.1000367)

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.omicsonline.org/submission/