

Medicinal Plants: Safety Must Be First and Foremost

Yann A Meunier*

Former Stanford University School of Medicine, Stanford Hospital and Clinics, USA

Introduction

In the general population the sentiment that what is natural is also wholesome prevails. While it may be generally true, unfortunately there are deleterious exceptions in the flora especially regarding plants used for real or perceived medical benefits.

More than 1,000 plants and associations of different plants can be found on the United States market. There are more than 7,000 species of medicinal plants. Several of them contain toxic substances which, if ingested, at worst can lead to death. Heavy metals such as mercury, lead, cadmium, copper, iron, manganese, nickel, zinc, and arsenic have been found in their analysis. In a study performed in the United States on 251 preparations coming from Asia, arsenic was found in 36 samples, mercury in 35 samples, and lead in 24 others. Lead encephalopathy and mercury poisoning have been described as detrimental effects of herbal decoctions or raw plant absorption. Anecdotally, a Tibetan child receiving "vitamin herbs" had absorbed 63 g of lead over 4 yrs. Another example is an Indian child who developed eye cancer from an herbal remedy containing arsenic and given to him by his parents [1,2].

Plant potential harmful effects of plant consumption include but are not limited to:

Some *idaturas* have an anticholinergic action while others contain atropine, which can lead to dry mouth, blurred vision, sensitivity to light, lack of sweating, dizziness, nausea, loss of balance, hypersensitivity reactions (such as skin rash), and tachycardia, for example.

- *Ginkgo biloba* has been shown to produce spontaneous cerebral or ocular hemorrhage, fainting, and hypoglycemia.
- *Chan su*, a traditional Chinese medicine used for sore throat and cardiac palpitations contains toxic secretions collected from toad glands and which have an action comparable to digoxin. They stimulate the cardiovascular system and can cause death by arrhythmia.

Herbs from the *aristolochia*, or *teucrium chamaedrys* families have triggered toxic hepatitis.

Kava (piper methylisticum), *pennyroyal* (mentha pulegium, sometimes used to induce miscarriage), *Jin bu huan* and others like a Chinese remedy dubbed "eternal life" have induced hepatitis fulminans.

Aristolochia fang chi, an herb containing aristocholic acid, can generate urinary tract cancer.

Garlic can increase the risk of hemorrhage.

Echinacea can raise the risk of allergy and immune suppression.

Valerian can enhance the risk of sedation.

Other adverse effects of medicinal plants include but are not limited to:

- Cardiovascular system: Arrhythmia, myocardial infarction, heart failure, pericarditis, hypertension.

- Neurological system: Stroke, fainting.
- Renal system: Tubulo-interstitial nephritis.
- Gastro-intestinal system: Diarrhea, constipation, vomiting, rectal bleeding.

Furthermore, medicinal plants can interact with allopathic medications, for example:

Hypericum perforatum reduces the action of anticoagulants used to prevent cardiac and cerebral complications of some diseases. It also decreases the efficacy of digoxin, theophylline (prescribed to treat asthma) as well as some anti-HIV medications.

Garlic enhances the action of antidiabetic drugs and the hepatic toxicity of paracetamol.

To ensure good traceability of all products, each plant and each group of plants must be specifically tagged. They must be analyzed thoroughly with state of the art equipment providing, for example:

- Organoleptic characteristics (aspect, color, smell, texture, taste).
- Identification (showing the matter identity or that of the expected product).
- Physical and chemical test results [3].
- Dosages of active ingredients: vitamins, minerals, preservatives (quality of the product).
- Microbiological analyses (compliance with healthcare criteria).
- Total number of germs, fungi, yeasts and research of specific germs results.

Using the following techniques, for example:

- Thin layer chromatography (particularly for the identification of plants).
- High performance liquid chromatography (in particular for the dosage of active ingredients and the purity of components).
- Gas chromatography (for the dosage of fatty acids, chemo types, essential oils).

*Corresponding author: Yann A Meunier, Senior Healthcare Consultant, Former Stanford University School of Medicine, Stanford Hospital and Clinics, USA, E-mail: ymeuniermd@gmail.com

Received August 22, 2017; Accepted September 06, 2017; Published September 13, 2017

Citation: Meunier YA (2017) Medicinal Plants: Safety Must Be First and Foremost. J Tradit Med Clin Natur 6: 241.

Copyright: © 2017 Meunier YA. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

- Infra-red and UV spectrophotometry (for the identification and purity of raw materials).
- Atomic absorption (for the dosage of minerals and heavy metals).

Systematic and systemic approaches must be adopted to warrant the safety of all medicinal plant consumers [4-29]. It is a public health necessity, a moral duty, and a professional obligation. This is why quality control is of paramount importance. Patients must be protected as much as possible: "Primum non nocere" should be the mantra of everyone involved in healthcare.

References

1. Talas ZS, Gogebakan A, Orun I (2013) Effects of propolis on blood biochemical and hematological parameters in nitric oxide synthase inhibited rats by N^ω-Nitro-L-arginine methyl ester. *Pak J Pharm Sci* 26: 915-919.
2. Ribeiro B, Lopes R, Andrade PB, Seabra RM, Gonçalves RF, et al. (2008) Comparative study of phytochemicals and antioxidant potential of wild edible mushroom caps and stipes. *Food Chem* 110: 47-56.
3. Wong JY, Chye FY (2009) Antioxidant properties of selected tropical wild edible mushrooms. *J Food Comp Anal* 22: 269-277.
4. Soares AA, de Sá-Nakanishi AB, Bracht A, da Costa SMG, Koehnlein EA, et al. (2013) Hepatoprotective effects of mushrooms. *Molecules* 18: 7609-7630.
5. Dias DA, Urban S, Roessner U (2012) A historical overview of natural products in drug discovery. *Metabolites* 2: 303-336.
6. Tsai SY, Huang SJ, Lo SH, Wu TP, Lian PY (2009) Flavour components and antioxidant properties of several cultivated mushrooms. *Food Chem* 113: 578-584.
7. Palacios I, Lozano M, Moro C, D'arrigo M, Rostagno MA, et al. (2011) Antioxidant properties of phenolic compounds occurring in edible mushrooms. *Food Chem* 128: 674-678.
8. Obodai M, Ferreira IC, Fernandes Â, Barros L, Mensah DLN, et al. (2014) Evaluation of the chemical and antioxidant properties of wild and cultivated mushrooms of Ghana. *Molecules* 19: 19532-19548.
9. Doğan HH, Şanda MA, Akata I (2012) Mn, Fe, K, Na, and P contents in some *Tricholoma* (Fr.) staude (Tricholomataceae) taxa from central Anatolia, Turkey. *Fresenius Environ Bull.* 21: 3389-3393.
10. Sarıkurucu C, Copur M, Yıldız D, Akata I (2011) Metal concentration of wild edible mushrooms in Soguksu National Park in Turkey. *Food Chem* 128: 731-734.
11. Sevcikova L, Pechova A, Pavlata L, Antos D, Mala E, et al. (2011) The effect of various forms of selenium supplied to pregnant goats on the levels of selenium in the body of their kids at the time of weaning. *Biol Trace Elem Res* 143: 882-892.
12. Talas ZS, Gok Y, Ozdemir I, Ates B, Gunal S, et al. (2015) Synthesis, antioxidant and anti-microbial properties of two organoselenium compounds. *Pak J Pharm Sci* 28: 611-616.
13. İlahi I, Samar S, Khan I, Ahmad I (2013) *In vitro* antioxidant activities of four medicinal plants on the basis of DPPH free radical scavenging. *Pak J Pharm Sci* 26: 949-952.
14. Kozarski M, Klaus A, Jakovljevic D, Todorovic N, Vunduk J, et al. (2015) Antioxidants of edible mushrooms. *Molecules* 20: 19489-19525.
15. Barros L, Falcão S, Baptista P, Freire C, Vilas-Boas M, et al. (2008) Antioxidant activity of *Agaricus* sp. mushrooms by chemical, biochemical and electrochemical assays. *Food Chem* 111: 61-66.
16. <http://www.saujs.sakarya.edu.tr/article/view/5000205272>
17. Shimada K, Fujikawa K, Yahara K, Nakamura T (1992) Antioxidative properties of xanthan on the autooxidation of soybean oil in cyclodextrin emulsion. *J Agric Food Chem* 40: 945-948.
18. Lee JC, Kim HR, Kim J, Jang YS (2002) Antioxidant property of an ethanol extract of the stem of *Opuntia ficus-indica* var. *saboten*. *J Agric Food Chem* 50: 6490-6496.
19. Erel O (2004) A novel automated direct measurement method for total antioxidant capacity using a new generation, more stable ABTS radical cation. *Clin Biochem* 37: 277-285.
20. Acharya K, Samui K, Rai M, Dutta BB, Acharya R, et al. (2004) Antioxidant and nitric oxide synthase activation properties of *Auricularia auricula*. *Indian J Exp Biol* 42: 538-540.
21. Cai M, Lin Y, Luo YL, Liang HH, Sun P, et al. (2015) Extraction, antimicrobial, and antioxidant activities of crude polysaccharides from the wood ear medicinal mushroom *Auricularia auricula-judae* (Higher Basidiomycetes). *Int J Med Mushrooms* 17: 591-600.
22. Xu S, Zhang Y, Jiang K (2016) Antioxidant activity *in vitro* and *in vivo* of the polysaccharides from different varieties of *Auricularia auricula*. *Food & function* 7: 3868-3879.
23. Mallikarjuna SE, Ranjini A, Haware DJ, Vijayalakshmi MR (2013) Mineral composition of four edible mushrooms. *Journal of Chemistry*.
24. Cheung YC, Siu KC, Wu JY (2013) Kinetic models for ultrasound-assisted extraction of water-soluble components and polysaccharides from medicinal fungi. *Food Bioproc Tech* 79: 214-220.
25. Akgül H, Nur AD, Sevindik M, Doğan M (2016) *Tricholoma terreum* ve *Coprinus micaceus*' un bazı biyolojik aktivitelerinin belirlenmesi. *Artvin Çoruh Üniversitesi Orman Fakültesi Dergisi* 17: 158-162.
26. Erel O (2005) A new automated colorimetric method for measuring total oxidant status. *Clin Biochem* 38: 1103-1111.
27. Gadd GM, Mowll JL, White C, Newby PJ (1986) Methods for assessment of heavy metal toxicity towards fungi and yeasts. *Environ Toxicol* 1: 169-185.
28. Hosseinihashemi SK, Salem MZ, HosseinAshrafi SK, Latibari AJ (2016) Chemical composition and antioxidant activity of extract from the wood of *Fagus orientalis*: Water resistance and decay resistance against *Trametes versicolor*. *BioResources* 11: 3890-3903.
29. Lee IK, Kim YS, Jang YW, Jung JY, Yun BS, et al. (2007) New antioxidant polyphenols from the medicinal mushroom *Inonotus obliquus*. *Bioorg Med Chem Lett* 17: 6678-6681.

Citation: Meunier YA (2017) Medicinal Plants: Safety Must Be First and Foremost. J Tradit Med Clin Natur 6: 241.

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.omicsonline.org/submit/