

# GDP<sub>βS</sub> Activates Excitatory Synapses in CA1 Pyramidal Cells by Disinhibiting the PKA Activating Pathway

Mohamed Ouardouz\*

NRAC, Gatineau, Quebec, Canada

## Abstract

**Objective:** During the early postnatal brain developmental stages, excitatory synapses mediated by AMPA receptors are weak or silent. Activity-dependent insertion of AMPA receptors into synapses depends on the activation of protein kinase A. In this work, we investigated the effect of Guanosine 5'-[βthiol] diphosphate (GDP<sub>βS</sub>) on excitatory and inhibitory synaptic currents in CA1 pyramidal cells at postnatal days 9-12.

**Methods:** Whole-cell patch-clamp recordings from identified hippocampal CA1 pyramidal cells were used. GDP<sub>βS</sub> was applied through the recording electrode.

**Results:** GDP<sub>βS</sub> induces an increase in excitatory synaptic current amplitude, but not in the inhibitory synaptic current amplitude. An analysis of the change in excitatory synaptic current amplitude in the presence of GDP<sub>βS</sub> revealed a progressive increase, which is blocked by the protein kinase A inhibitor Rp-3',5'-cyclic monophosphothioate triethylamine (Rp-cAMP), suggesting that GDP<sub>βS</sub> inhibits G-protein with a tonic negative control on a protein kinase A activating pathway. In addition, GDP<sub>βS</sub> has no effect on paired-pulse facilitation, suggesting that the glutamate release machinery is not affected. Moreover, as GDP<sub>βS</sub> was applied to postsynaptic neurons, the increase in excitatory postsynaptic current amplitude is related to changes at the postsynaptic side.

**Conclusion:** Those results suggest that in developing hippocampal CA1 pyramidal cells, the tonic inhibition of a protein kinase A activating pathway by a G-protein prevents the activation of excitatory synapses.

**Keywords:** Hippocampus; EPSCs; Rat; Development; G-protein; Protein kinase A

## Introduction

Excitatory synaptic transmission in the central nervous system is mainly mediated through the activation of glutamate receptors. In the early development of the rodent hippocampus, excitatory synapses contain mostly N-methyl-D-aspartate (NMDA) receptors, lack α-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) receptors, and are considered silent synapses as they are inactive at the resting membrane potential [1-4]. These synapses can be turned on by the insertion of new AMPA receptors [5-11]. The cascade of events that regulates AMPA receptor insertion into synapses requires phosphorylation by protein kinase A (PKA) [8]. In addition, PKA activation drives AMPA receptors into synapses [6-8,10,12,13].

The mechanisms by which AMPA receptors are prevented from inserting into synapses during the early developmental stage remain unclear. Since the insertion of AMPA receptors depends on phosphorylation by PKA, PKA is possibly inactive at silent synapse sites, thereby preventing the insertion of new receptors at these sites. In the present work, the effect of GDP<sub>βS</sub> G-proteins inhibitor on evoked excitatory postsynaptic currents (EPSCs) in CA1 hippocampal pyramidal cells is investigated. The results suggest the involvement of G-proteins in inhibiting the PKA activating pathway involved in the activation of silent or weak synapses.

## Materials and Methods

Male Wistar rat pups (from different litters, n = 18) were obtained from the animal care unit at the University of British Columbia on postnatal day 9-12. Pups were kept with their mother (standard cages) in a 12-hour light/dark cycle with freely available food and water. Animal care and use conformed to the guidelines and the policies of the Animal Care Committee at The University of British Columbia.

Hippocampal slices (400 μm) were prepared from male Wistar rat (9-12 days old) as previously described [14]. The rat pups were anesthetized using isoflurane (Sigma aldrich) then decapitated, and the brain was quickly removed and placed in cold artificial cerebrospinal fluid containing the following: 120 mM NaCl, 3 mM KCl, 1.8 mM NaH<sub>2</sub>PO<sub>4</sub>, 2 mM MgSO<sub>4</sub>, 2 mM CaCl<sub>2</sub>, 26 mM NaHCO<sub>3</sub> and 10 mM dextrose; the fluid was also saturated with 95% O<sub>2</sub> and 5% CO<sub>2</sub>. Hippocampal slices were cut in cold ACSF with a vibrating blade microtome Leica (Leica Microsystems) and transferred into a container filled with oxygenated ACSF at room temperature. After an incubation period of 1 h, a slice was placed into a recording chamber mounted in an upright microscope (Zeiss), and continuously perfused with oxygenated ACSF.

Electrophysiological recordings were made in a whole-cell patch-clamp using an Axopatch 200A (Molecular Devices, Foster City, CA, USA). The recording pipette contained the following: 145 mM K-gluconate, 10 mM HEPES, 10 mM KCl, 1 mM K<sub>4</sub>-bis-(2-aminophenoxy)-N,N,N',N'-tetra acetic acid (BAPTA, Sigma Aldrich), 5 mM Mg-ATP, 0.1 mM CaCl<sub>2</sub>, 0.4 mM Na<sub>3</sub>-GTP. The pH was adjusted to 7.2-7.3 with KOH. Na<sub>3</sub>-GTP was replaced by GDP<sub>βS</sub> (Sigma Aldrich) to assess the role of the G-protein on synaptic responses. Recordings were acquired at 5 KHz and filtered

\*Corresponding author: Mohamed Ouardouz, NRAC, 39 du météore Gatineau, Quebec, J9A 3B5, Canada, E-mail: [ouardouz@bell.net](mailto:ouardouz@bell.net)

Received June 28, 2014; Accepted September 29, 2014; Published October 06, 2014

**Citation:** Ouardouz M (2014) GDP<sub>βS</sub> Activates Excitatory Synapses in CA1 Pyramidal Cells by Disinhibiting the PKA Activating Pathway. J Neurol Neurophysiol 5: 235. doi:10.4172/2155-9562.1000235

**Copyright:** © 2014 Ouardouz M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

at 2 KHz and digitized using digidata 1300 and windows based programme Pclamp 9 (molecular devices). Series resistance and holding currents were monitored during all recordings.

Inhibitory postsynaptic currents were evoked in the presence of 6,7-dinitroquinoxaline-2,3-dione(DNQX, 10  $\mu$ M, Tocris) and (+2)-amino-5-phosphonovaleric acid (APV, 50  $\mu$ M, Tocris). Excitatory synaptic currents were evoked by electrical stimulation using a bipolar electrode placed in stratum radiatum. The connection between CA3 and CA1 was cut.

GABA $_B$  receptor-mediated postsynaptic currents were evoked by a train of 4 pulses (0.1-0.2ms, 200-400  $\mu$ A) at 200 Hz applied in stratum radiatum in the presence of ionotropic glutamate receptor antagonists, DNQX, (10  $\mu$ M), APV (50  $\mu$ M) and GABA $_A$ -receptor antagonist: bicuculline methiodide (30  $\mu$ M, Tocris).

Data are presented as mean + sem and were analysed using Student's t-test and paired t-test (Microsoft Excel); n represents the number of cells. The difference between groups is considered significant if  $p < 0.05$ .


## Results

Experiments were performed in visually identified pyramidal cells in the CA1 area of the hippocampus using an upright Zeiss microscope. To verify that GDP $\beta$ S applied through the patch pipette, blocks G-proteins, we tested its effect on GABA $_B$  receptor, a member of G-protein activated receptors. Electrical stimulation of stratum radiatum in the presence of DNQX (10  $\mu$ M), AP-5 (50  $\mu$ M) and bicuculline (30  $\mu$ M) evoked GABA $_B$  inhibitory synaptic potentials in older animals (3 weeks). In the presence of GDP $\beta$ S in the patch solution, the amplitude of GABA $_B$  IPSPs significantly decreased with time (to  $37.2 \pm 6.4$  % after 10 min of recording;  $n = 5$ ,  $P = 0.004$ ; Figure 1). In contrast, GDP $\beta$ S had no effect on GABA $_A$  inhibitory synaptic currents (IPSC) (Figure. 2B). GABA $_A$  IPSCs were evoked via electrical stimulation in the stratum radiatum in the presence DNQX, 10  $\mu$ M) and APV, 50  $\mu$ M) at -30 mV. The mean amplitude of the IPSCs is not significantly affected by GDP $\beta$ S ( $53.8 \pm 3.5$  pA after breaking into the cell and  $46.7 \pm 3.4$  pA after 10 minutes of recording (Figure. 2B,  $n = 3$ ,  $p = 0.12$ , paired t-test).


Excitatory synaptic currents were evoked in the absence of DNQX and APV at -70 mV (Figure 2). The mean amplitude of EPSCs significantly increased from  $98.2 \pm 9.6$  pA after breaking into the cell to  $230.5 \pm 30.2$  pA after 10 minutes of recordings (Figure 2C-D,  $n = 17$ ,  $p = 0.00003$ , paired t-test). The mean amplitude of EPSCs significantly increased with time with GDP $\beta$ S in the recording solution as compared to the control solution (Figure 2). Under the control conditions, the increase in mean EPSC amplitude is not statistically significant (i.e.,  $114.3 \pm 14.4$  pA after breaking into the cell and  $154.4 \pm 30.7$  pA after 10 minutes of recording;  $n = 14$ ,  $p = 0.5$ , paired t-test). This increase in EPSC amplitude induced by GDP $\beta$ S is not accompanied by changes in EPSC kinetics. The EPSC rise time and decay time are not affected by GDP $\beta$ S (i.e., the rise time after breaking into the cell was  $4.9 \pm 0.5$  ms, and the rise time after 10 minutes of recording in the presence of GDP $\beta$ S was  $4.2 \pm 0.3$  ms,  $p = 0.1$ , paired t-test; the EPSC decay time after breaking into the cell was  $57.0 \pm 6.5$  ms, and the decay time after 10 minutes of recording in the presence of GDP $\beta$ S was  $51.9 \pm 3.2$  ms,  $p = 0.5$ , paired t-test).

GDP $\beta$ S is known to block all G-proteins, including the ones that inhibit the production of second messengers, like cyclic adenosine mono-phosphate (cAMP) (e.g., Gi-proteins), which activate PKA.



PKA is known to modulate AMPA receptors [15,16]. To test the involvement of PKA in EPSC potentiation by GDP $\beta$ S, we used Rp-3',5'-cyclic monophosphothioetriethyl-amine (Rp-cAMP), an analogue of cAMP with a selective and competitive antagonist effect on PKA [17]. The application of Rp-cAMP (100  $\mu$ M, in the recording solution) completely blocks changes in EPSC amplitude induced by GDP $\beta$ S (i.e.,  $103.8 \pm 41.5$  pA after breaking into the cell and  $99.3 \pm 38.1$  pA after 10 minutes of recording;  $n = 5$ ,  $p = 0.65$ , paired t-test)



**Figure 1:** GDP $\beta$ S blocks GABA $_B$  receptor-mediated inhibitory postsynaptic potentials. GABA $_B$  receptor mediated synaptic potentials (IPSPs) were evoked by a train of electrical stimulations (4 pulses at 200Hz). GDP $\beta$ S Applied through the recording electrode induces a progressive reduction of GABA $_B$  IPSPs.



**Figure 2:** GDP $\beta$ S induces an increase in excitatory postsynaptic current (EPSC) amplitude but does not affect inhibitory synaptic currents (IPSC) amplitude. A: Representative traces of IPSCs recorded at -30 mV just after breaking into the cell (i.e., the control) and after 10 minutes of recording in the presence of Guanosine 5'-[ $\beta$ thio]diphosphate (GDP $\beta$ S, 10 minutes GDP $\beta$ S). B: Data summary histogram showing no significant effect of GDP $\beta$ S on IPSC amplitude. C: Representative traces of EPSCs recorded at -70 mV just after breaking into the cell (i.e., the control) and after 10 minutes of recording in the presence of GDP $\beta$ S (10 minutes GDP $\beta$ S). D: Data summary histogram showing a significant increase in EPSC amplitude by GDP $\beta$ S. Error bar represent the sem, \*\*\* statically significant  $P < 0.001$ .



(Figure 3). These results suggest that GDP $\beta$ S blocks a Gi-protein that inhibits adenylatecyclase and cAMP production. The disinhibition of adenylatecyclase increases the production of cAMP and the activation of PKA, which potentiates EPSCs.

To further characterize the mechanisms by which GDP $\beta$ S induces an increase in EPSC amplitude, we analysed the changes in spontaneous EPSC amplitude and frequency between the first minute after breaking into the cell and the eighth to tenth minute of recordings. As reflected by the inter-event interval, the frequency of spontaneous EPSCs significantly increased with time in the presence of GDP $\beta$ S (i.e., inter-event interval:  $3.98 \pm 0.45$  s during the first min after breaking into the cell and  $1.91 \pm 0.31$  s during the eighth to tenth minute of recordings;  $n = 4$ ;  $p = 0.0057$  in a paired t-test). Additionally, the spontaneous EPSC amplitude did not significantly change (i.e.,  $4.7 \pm 0.3$  pA during the first minute of recording, and  $4.3 \pm 0.5$  pA during the eighth to tenth minute of recordings;  $n = 4$ ,  $p = 0.5$  in a paired t-test) (Figure 4). The increase in spontaneous EPSC frequency may be due to changes in the probability of glutamate release from the presynaptic terminal.

To assess the effect of GDP $\beta$ S even if it was applied via the recording pipette on glutamate release, we used paired pulse stimulation at 100 and 200 ms intervals. Under the control conditions, the amplitude of the second EPSC increases by  $39.2 \pm 4.8$  % ( $n = 12$ , at 200 ms interval) and as compared to the amplitude of the first EPSC. After 10 minutes of recording in the presence of GDP $\beta$ S, the percentage increase in the amplitude of the second EPSC as compared to the first is not statistically different from that of the control ( $27.2 \pm 3.5$  %;  $n = 13$ ,  $p > 0.05$  at 200 ms interval), thereby suggesting that GDP $\beta$ S does not affect the probability of glutamate release.

## Discussion

Silent excitatory glutamatergic synapses lacking AMPA receptors are present in CA1 pyramidal cells during the first postnatal days [1-4]. In this study, we report that GDP $\beta$ S, which is known to block G-protein activity, potentiates the evoked EPSCs and the effect is blocked by the PKA inhibitor Rp-cAMP.

The increase in evoked EPSC amplitude by GDP $\beta$ S depends on

PKA activation as the application of its inhibitor Rp-cAMP reverses the effect of GDP $\beta$ S. The G-proteins involved in the activation of adenylatecyclase are named Gs, and the one involved in its inhibition are named Gi. GDP $\beta$ S inhibits both Gi and Gs-proteins [18-21]. The inhibition by GDP $\beta$ S of Gi-proteins will remove the negative control over adenylatecyclase and increase the production of cAMP, hence increasing the activity of PKA. Evoked EPSCs were recorded at a holding membrane potential of -70 mV and are mainly mediated by AMPA receptors since most NMDA receptors are inactive at this membrane potential. The increase in evoked EPSC amplitude by GDP $\beta$ S is mostly due to the modulation of AMPA receptors. In fact, previous studies have shown that an increase in the activity of PKA modulates AMPA receptors mediated current [15,16] and plays an important role in AMPA receptor insertion [6-8,13,22-24].

The increase in EPSC amplitude by GDP $\beta$ S is accompanied by an increase in spontaneous EPSC frequency without a change in the spontaneous EPSC amplitude. The increase in spontaneous EPSC frequency may be due to a change in the probability of glutamate release at the presynaptic terminals. Under our experimental conditions, the probability of glutamate release is not affected by GDP $\beta$ S in the recording solution as revealed by the lack of a GDP $\beta$ S effect in the paired pulse facilitation of EPSCs. In addition, GDP $\beta$ S was applied into the postsynaptic cell. Therefore, the change in evoked EPSC amplitude and spontaneous EPSC frequency is likely due to a change on the postsynaptic side. The most probable scenario is the insertion of AMPA receptors and the activation of silent synapse as described for long-term potentiation in CA1 pyramidal cells [5,9,22-24]. GDP $\beta$ S induces an increase in evoked EPSC amplitude via the activation of silent synapses through the insertion of AMPA receptors after phosphorylation by PKA.

In conclusion, our data highlighted the presence of a Gi-protein-dependent system that prevents the activation of excitatory synapses though the modulation of PKA activity. This may be done through the inhibition of adenylatecyclase, which produces cAMP; by the activation of Gi-protein dependent receptors. It will be interesting to determine which of those systems is involved in the sustained inhibition of the PKA activation pathway and hindering the insertion of AMPA receptors into synapses.

#### Acknowledgments

I wish to thank an American Manuscript Editor for proof reading an earlier version of the manuscript. This work was conducted in neuroscience research laboratory in the Department of Pharmacology and Therapeutics at the University of British Columbia.

#### References

1. Isaac JT, Nicoll RA, Malenka RC (1995) Evidence for silent synapses: implications for the expression of LTP. See comment in PubMed Commons below *Neuron* 15: 427-434.
2. Malenka RC, Nicoll RA (1997) Silent synapses speak up. See comment in PubMed Commons below *Neuron* 19: 473-476.
3. Liao D, Hessler NA, Malinow R (1995) Activation of postsynaptically silent synapses during pairing-induced LTP in CA1 region of hippocampal slice. See comment in PubMed Commons below *Nature* 375: 400-404.
4. Durand GM, Kovalchuk Y, Konnerth A (1996) Long-term potentiation and functional synapse induction in developing hippocampus. See comment in PubMed Commons below *Nature* 381: 71-75.
5. Liao D, Scannevin RH, Huganir R (2001) Activation of silent synapses by rapid activity-dependent synaptic recruitment of AMPA receptors. See comment in PubMed Commons below *J Neurosci* 21: 6008-6017.
6. Dias RB, Ribeiro JA, Sebastião AM (2012) Enhancement of AMPA currents and GluR1 membrane expression through PKA-coupled adenosine A(2A)

receptors. See comment in PubMed Commons below *Hippocampus* 22: 276-291.

7. Zheng Z, Keifer J (2009) PKA has a critical role in synaptic delivery of GluR1- and GluR4-containing AMPARs during initial stages of acquisition of in vitro classical conditioning. See comment in PubMed Commons below *J Neurophysiol* 101: 2539-2549.
8. Esteban JA, Shi SH, Wilson C, Nuriya M, Huganir RL, et al. (2003) PKA phosphorylation of AMPA receptor subunits controls synaptic trafficking underlying plasticity. See comment in PubMed Commons below *Nat Neurosci* 6: 136-143.
9. Lu W, Man H, Ju W, Trimble WS, MacDonald JF, et al. (2001) Activation of synaptic NMDA receptors induces membrane insertion of new AMPA receptors and LTP in cultured hippocampal neurons. See comment in PubMed Commons below *Neuron* 29: 243-254.
10. Kollerker A, Zhu JJ, Schupp BJ, Qin Y, Mack V, et al. (2003) Glutamatergic plasticity by synaptic delivery of GluR-B(long)-containing AMPA receptors. See comment in PubMed Commons below *Neuron* 40: 1199-1212.
11. Man HY, Sekine-Aizawa Y, Huganir RL (2007) Regulation of {alpha}-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor trafficking through PKA phosphorylation of the Glu receptor 1 subunit. See comment in PubMed Commons below *Proc Natl Acad Sci U S A* 104: 3579-3584.
12. Ehlers MD (2000) Reinsertion or degradation of AMPA receptors determined by activity-dependent endocytic sorting. See comment in PubMed Commons below *Neuron* 28: 511-525.
13. Cousin MA, Evans GJ (2011) Activation of silent and weak synapses by cAMP-dependent protein kinase in cultured cerebellar granule neurons. See comment in PubMed Commons below *J Physiol* 589: 1943-1955.
14. Ouardouz M, Xu JY, Sastry BR (2006) Theta bursts set up glutamatergic as well as GABA-ergic plasticity in neonatal rat hippocampal CA1 neurons. See comment in PubMed Commons below *Brain Res* 1068: 65-69.
15. Banke TG, Bowie D, Lee H, Huganir RL, Schousboe A, et al. (2000) Control of GluR1 AMPA receptor function by cAMP-dependent protein kinase. See comment in PubMed Commons below *J Neurosci* 20: 89-102.
16. Roche KW, O'Brien RJ, Mammen AL, Bernhardt J, Huganir RL (1996) Characterization of multiple phosphorylation sites on the AMPA receptor GluR1 subunit. See comment in PubMed Commons below *Neuron* 16: 1179-1188.
17. Rothermel JD, Stec WJ, Baraniak J, Jastorff B, Botelho LH (1983) Inhibition of glycogenolysis in isolated rat hepatocytes by the Rp diastereomer of adenosine cyclic 3',5'-phosphorothioate. See comment in PubMed Commons below *J Biol Chem* 258: 12125-12128.
18. Pieroni JP, Jacobowitz O, Chen J, Iyengar R (1993) Signal recognition and integration by Gs-stimulated adenylyl cyclases. See comment in PubMed Commons below *Curr Opin Neurobiol* 3: 345-351.
19. Rens-Domiano S, Hamm HE (1995) Structural and functional relationships of heterotrimeric G-proteins. See comment in PubMed Commons below *FASEB J* 9: 1059-1066.
20. Oldham WM, Hamm HE (2006) Structural basis of function in heterotrimeric G proteins. See comment in PubMed Commons below *Q Rev Biophys* 39: 117-166.
21. Holinstat M, Oldham WM, Hamm HE (2006) G-protein-coupled receptors: evolving views on physiological signalling: symposium on G-protein-coupled receptors: evolving concepts and new techniques. See comment in PubMed Commons below *EMBO Rep* 7: 866-869.
22. Bredt DS, Nicoll RA (2003) AMPA receptor trafficking at excitatory synapses. See comment in PubMed Commons below *Neuron* 40: 361-379.
23. Barry MF, Ziff EB (2002) Receptor trafficking and the plasticity of excitatory synapses. See comment in PubMed Commons below *Curr Opin Neurobiol* 12: 279-286.
24. Santos SD, Carvalho AL, Caldeira MV, Duarte CB (2009) Regulation of AMPA receptors and synaptic plasticity. See comment in PubMed Commons below *Neuroscience* 158: 105-125.

Citation: Ouardouz M (2014) GDP $\beta$ S Activates Excitatory Synapses in CA1 Pyramidal Cells by Disinhibiting the PKA Activating Pathway. *J Neurol Neurophysiol* 5: 235. doi:[10.4172/2155-9562.1000235](https://doi.org/10.4172/2155-9562.1000235)