

# Microbiological Study of Turkey Meat Marketed in Kenitra (North-oust of Morocco)

Jaber H\*, Ijoub R, Zaher A, Chakit M, Rhaïem N, Bourkhiss B and Ouhssine M

Department of Biology, Ibn Tofail University, Kenitra, Morocco

## Abstract

Turkey meat is considered among meat of poultry most consumed by the Moroccan population, and this due to its richness in vitamins, proteins and essential elements for growth, which gives him an excellent nutritional value. In addition, it is characterized by a low content of fatty acids. But this highly perishable commodity could constitute a real danger of food-borne illness. The aim of present study is to assess the hygienic quality of 168 samples collected from twenty four points of sales located in six neighborhoods in Kenitra. The analysis of the microbiological quality of the samples shows that the percentage of non-compliance for the parameter Total Aerobic Mesophilic Flora is 35%, with the mean concentration 7.21 ( $\log_{10}$  cfu/g). The respective percentages of non-compliance of the total and fecal coliforms are 68 and 75% with bacterial concentrations 6.69 ( $\log_{10}$  cfu/g) and 6.42 ( $\log_{10}$  cfu/g). This non-compliance is due to the presence of the germs of fecal contamination among the identified isolates. The four identified species showed following prevalence: *Escherichia coli* (67.9%), *Klebsiella pneumonia* (17, 9%), *Pseudomonas* sp. (14.1%) and *Salmonella* (0%).

**Keywords:** Turkey meat; Kenitra; Non-compliance; Bacterial concentration; Contamination

## Introduction

In Morocco, the production of turkey meat has experienced a pace of significant development, it is increased from 70,000 tones in 2010 to 80,000 tones in 2015 and its consumption is becoming more and more important given its price suitable for the citizen [1].

Turkey meat is considered among the most consumed poultry meats in the city of Kenitra knowing that the most requested body part by the citizen is the muscle. It is regarded as a potential source of food-borne illness caused by lack of hygiene [2,3]. Turkey meat is a highly perishable food whose hygienic quality depends on the quality of the different operations including the slaughter, cutting, storage and distribution [4]. In Morocco, between 2000 and 2004, 7,118 cases of food-borne infections have been reported in which more than 86% are of bacterial origin [5]. The contamination of foods of animal origin and mainly of meat products is responsible for 28% of cases of food-borne diseases [6]. Many factors have contributed to the increase of foodborne diseases. Mass production, environmental factors, and inadequate knowledge on food handling have contributed to the increase of contaminated foodstuffs [5].

This problem of contamination, therefore, proves the importance of the food hygiene. This work constitutes a contribution to the evaluation of the hygienic quality of turkey meat marketed in Kenitra city.

## Materials and Methods

### Sample collection

A total of 168 samples of turkey meat have been collected, in aseptic conditions, from 24 points of sale in the city of Kenitra to reason of seven samples by the point of wind, during the period of June-September 2015. Each sample was placed in sterile stomacher bag and transported in a cooler to the laboratory. In the arrival, the samples are analyzed immediately in three repetitions.

### Bacteriological analysis

A portion of 25 g of each sample is cut aseptically and then crushed carefully using a chopper ultra-turax and then, added

in an Erlenmeyer flask containing 225 ml of sterile physiological water. From this last, we have prepared a series of tenfold dilutions ranging up to  $10^{-6}$  [7]. The bacteriological analysis has focused on the following germs: the Total Aerobic Mesophilic Flora (TAMF), the Total and Fecal coliforms, *Escherichia coli*, *Klebsiella* sp., *Pseudomonas* sp. and *Salmonella* sp.

### Enumeration of the Total Aerobic Mesophilic Flora (TAMF):

The enumeration of the TAMF was carried out by using the method of seeding in depth on the Plate Count Agar (PCA, Biokar). The series of tenfold dilution were deposited aseptically in different boxes of sterile Petri dishes containing 10-15 ml of the medium PCA maintained at  $45 \pm 1^\circ\text{C}$ . The incubation was driving at  $30^\circ\text{C}$  for 72 h. Only the boxes of Petri dishes with a number of colonies between 30 and 300 have been retained for the enumeration [8].

### Enumeration of total and fecal coliforms:

The Enumeration of Total and Fecal coliforms was carried out on Violet Red Bile Lactose Agar (VRBL). Total coliforms are counted after 24 h of incubation at  $37^\circ\text{C}$ . For Fecal coliforms, the boxes of Petri dishes are incubated at  $44^\circ\text{C}$  for 24 h [9]. Only colonies with a yellow aspect orange and a diameter of more than 0.5 mm were counted.

**Escherichia coli:** Colonies of Fecal coliforms were isolated in the Middle VRBL and subcultured on another selective medium Eosin Methylene Blue (EMB). Then, the boxes of Petri dishes have incubated at  $37^\circ\text{C}$  for 24 h. The presence of *Escherichia coli* (*E. coli*) is indicated by the metallic reflection and the green color of the Colonies [10].

**\*Corresponding author:** Jaber H, Department of Biology, Faculty of Science, University Ibn Tofail, University Campus, Kenitra, 14000, Morocco, Tel: (00212) 6 01 21 72 15; E-mail: [jaberhassna@yahoo.com](mailto:jaberhassna@yahoo.com)

**Received** July 20, 2017; **Accepted** July 24, 2017; **Published** July 31, 2017

**Citation:** Jaber H, Ijoub R, Zaher A, Chakit M, Rhaïem N, et al. (2017) Microbiological Study of Turkey Meat Marketed in Kenitra (North-oust of Morocco). J Nutr Food Sci 7: 620. doi: [10.4172/2155-9600.1000620](https://doi.org/10.4172/2155-9600.1000620)

**Copyright:** © 2017 Jaber H, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Klebsiella sp.:** Similarly, colonies of Fecal coliforms were isolated from the mid-VRBL and sown on the mid-Eosin Methylene Blue (EMB), after their incubation at 37°C, for 48 h [10]. The *Klebsiella* colonies have shown large, rounded and very mucous membranes [11].

**Pseudomonas sp.:** The presence of colonies of *Pseudomonas* has been carried out on the mid-Pseudomonas Agar Base, after incubation boxes of Petri dishes to 44°C for 48 h [12]. The *Pseudomonas* is characterized by the yellow color with a clear halo.

**Salmonella:** The research of *Salmonella* requires the achievement of several of the systematic steps. The pre-enrichment was carried out in 225 ml of buffered peptone water (Difco) incubated at 37°C for 24 h, followed by an enrichment in the selective liquid medium: Rappaport Vassiliadis (RV). The high concentration of magnesium chloride and the presence of the green malachite are slowing down the growth of microorganisms except salmonella. A volume of 0.1 ml of the medium pre-enriched was added to 10 ml of the medium RV and incubated at +41°C ± 2°C, for 24 h. The isolation was carried out on the mid-Hektoen (Difco). The selective enriched liquid medium RV was stocked on the selective agar medium Hecktoen, previously sunk in boxes of Petri dishes, and were incubated at +36°C ± 2°C, for 24 h [13,14].

### Biochemical identification

All above-cited bacteria have been tested by Gram staining and by oxidase activity. The identification of the isolates was carried out by the determination of the biochemical profile using the API gallery 20 (Bio Mérieux, Marcy l'Etoile, France).

### Results and Discussion

The non-conformity of samples was assessed according to the joint order of the Ministry of Agriculture and Rural Development, the Ministry of Health and the Ministry of Industry, Trade and Telecommunications No. 624-04 Safar 1425 (8 April 2004) published in the Official Bulletin n: 5214-30 Rabbi 1-1425 (20-5-2004) relating to microbiological standards and which must meet the animal foodstuffs. The acceptable upper limit of the Total Aerobic Mesophilic Flora (TAMF) in the meat of poultry is of the order of 6.7 (log<sub>10</sub> cfu/g), 2 (log<sub>10</sub> cfu/g) for Total coliforms and 4 (log<sub>10</sub> cfu/g) for Fecal coliforms. However, the Moroccan standards are devoid of security criteria concerning *Escherichia coli*, *Klebsiella pneumonia* and *Pseudomonas* sp. Therefore, the comparison of our results was made in reference to other studies. For *Salmonella* and according to the Moroccan standard, it must be absent in 25 g of the sample of the meat [13].

### Percentage of the non-compliance of the studied samples

**Total aerobic mesophilic flora:** The results of the microbiological analysis of turkey meat samples marketed in the city of Kenitra (Table 1) show that the average value of the Total Aerobic mesophilic flora (FMAT) 6.44 (log<sub>10</sub> cfu/g) is less than the value declared by the Moroccan standards of the microbiological safety criteria of foods (6.7 (log<sub>10</sub> cfu/g)). These results are lower than those of Amara et al. [14] and Cohen et al. [15], the found values are respectively in the order of 7.15 and 7.4 (log<sub>10</sub> cfu/g). The values of our results are lower also to those found in the meat of poultry by Chaiba and Rhazi Filali [16] (7.45 (log<sub>10</sub> cfu/g)). The non-compliance of the studied samples has presented a rate of 35% (Figure 1).

**Total coliforms:** The degree of contamination of the samples by the total germs has been assessed by the calculation of the average of the loads found in the analyzed samples, the load of Total coliforms is 3.87 (log<sub>10</sub> cfu/g). This value is lower than that found in the carcasses of

chickens of the traditional slaughter of Meknes city; the abundance of Total coliforms was 4.4 (log<sub>10</sub> cfu/g) [16]. In our study, it was found that 68% of the analyzed samples are non-complaint (Figure 2).

**Fecal coliforms:** The average rate of contamination of turkey meat samples is 5.98 (log<sub>10</sub> cfu/g). In this study, the load is less than these mentioned by El-Allaoui et al. [13] (8.15 (log<sub>10</sub> cfu/g)), and higher than these reported by Cohen in 2006 (2.5 ± 1.2 (log<sub>10</sub> cfu/g)) [5]. Our results revealed a rate of non-compliance of 75% of the totality of the analyzed samples (Figure 3). This contamination would seem to be obvious during the evisceration although the majority of these germs can be caused the gastro-enteritis disorders for consumer.

### Origin of the non-compliance





On 168 samples, 118 bacterial strains were isolated (Table 2). The results of the microscopic study and the biochemical identification of bacteria responsible for the non-compliance of turkey meat show that coliform bacteria are responsible for the non-compliance of the majority of the analyzed samples.

**Escherichia coli:** The prevalence of *Escherichia coli* is 67.8%. This value is compatible with that found by Patyal et al. (68%) [17], it is lower than that found by El-Allaoui et al. (83%) [18], and largely superior to that found by Gupta (22.5%) [19] and Iroha (2%) [20]. *E. coli* is a hygienic indicator, its presence in turkey meat is a sign of enteric pathogenicity and which may cause a risk of contamination to the consumer. It was found in abundance. This can be explained by the

Points of sale	TAMF (log <sub>10</sub> cfu/g)	CT (log <sub>10</sub> cfu/g)	CF (log <sub>10</sub> cfu/g)
1	6.25	2.65	6.33
2	6.48	3.57	6.33
3	6.46	3.14	6.01
4	6.52	3.15	5.92
5	6.45	3.20	5.62
6	6.33	3.02	5.40
7	6.28	3.29	5.34
8	6.28	3.65	5.38
9	6.40	3.85	5.34
10	6.41	4.02	5.53
11	6.51	3.98	5.71
12	6.41	4.11	5.78
13	6.51	4.53	6.13
14	6.53	4.69	6.19
15	6.47	5.30	6.40
16	6.57	5.40	6.49
17	6.46	4.97	6.52
18	6.48	4.92	6.49
19	6.50	4.21	6.36
20	6.40	3.57	6.03
21	6.48	3.45	5.98
22	6.49	3.22	5.89
23	6.45	3.40	5.97
24	6.50	3.67	6.27
Average	6.44	3.87	5.98
Standard*	6.7	2	4

C.T: Total Coliforms; C.F: Fecal Coliforms; TAMF: Total Aerobic Mesophilic Flora. \*According to the joint order of the Ministry of Agriculture and Rural Development, the Ministry of Health and the Ministry of Industry, Trade and Télécommunications No. 624-04 Safar 1425 (8 April 2004) published in the Official Bulletin n: 5214-30 Rabbi 1-1425 (20-5-2004) relating to microbiological criteria of use for food.

**Table 1:** Result of microbiological analyzes of samples collected from 24 points of sale.



lack of good hygienic practices among the personnel responsible for the sale or in their premises during the process of slaughter and cutting of turkey meat (Figure 4).

***Klebsiella pneumoniae*:** The prevalence of *Klebsiella pneumoniae* was 17.8%. This rate is far lower than that found in the turkey meat by Davis (38%) [21], and by Neslihan in the chicken meat (47%) [22]. The presence of *Klebsiella pneumoniae* in meat is an indicator of faecal contamination and a sign of poor conditions of hygiene [23,24].

***Pseudomonas sp.*:** For *Pseudomonas sp.* the prevalence was 14.4%. This value is very lower than that found by Arnaut-Rollier in chicken meat (19.2%) [25]. The presence of this bacterium is probably due to its prototrophic properties and to these needs in mineral nitrogen, carbohydrates, mineral salts but it does not require the presence of the growth factors as well as its psychrotroph characteristic which the growth temperature varies between 0 and 37°C [26,27]. Therefore, its presence in the analyzed samples denotes a secondary contamination which could be associated to a bad conservation in the time [28] and the modification of the organoleptic characteristics of the meat.

***Salmonella sp.*:** The *Salmonella* presents an importance in the veterinarian areas and on the medical plan, by the economic losses caused by animal diseases and by the high incidence in the consumer. Therefore, the typhoid fever and food-borne diseases due to *Salmonella* [29] provoke most often the gastro-enteritis. *Salmonella* is involved in 76.2% of food-borne illness. Given the broad spectrum of animals that can be carriers of *Salmonella*, a great variety of food products may be at the origin of a human contamination: meat and particularly poultry, meat products, eggs and dairy products. Salmonellosis can give rise to important spots which can reach a national scale if a food marketed to wide dissemination is contaminated.

In this study, the rate contamination of turkey meat by *Salmonella* is zero (0%), although the contamination of faecal origin has been relatively important. Indeed, a high load in fecal coliforms can predict the presence of salmonella.

The rate of contamination by *Salmonella* is variable according to studies. In Morocco, Bennani et al. [30] have reported a rate of 13% of the positive samples to the presence of *Salmonella sp* on samples of poultry meat in Fez [30]. In Casablanca, Karraouan et al. have found a contamination rate of 20.3% in Turkey meat [31]. Chaiba [32] reported a rate of 20.83% in chicken meat in Meknes. However, Beli et al. have revealed a low prevalence of *Salmonella* in turkey meat in Albania (8.2%) [33]. In Ireland, Jordan and al have found a rate of 3.1% [34] and in the United Kingdom, Zhao et al. have found a rate of 5.6% [35].

The rate of Isolation of *Salmonella* is higher after evisceration, which confirms that the operations of slaughter and preparation in the slaughterhouse and cutting plants can be considered as a major source of contamination of the meat by *Salmonella*.

The presence of the pathogenic germs in the analyzed samples shows that it is interesting to determine the origin of contamination. This can be achieved only by the adoption and implementation of the quality approach in applying the guidelines of the Hazard Analysis Critical Control Point (HACCP) plan, as well as the mastery of the guide to good hygiene practices.

## Conclusion

The results of analyses of 168 turkey meat samples, collected from 24 sale points in the city of Kenitra, show the presence of four bacterial strains: *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas sp.* and

Species	F	G	Ox	Cat	Glu	Lac	Cit	Ur	TDA	ONPG	Vp	H <sub>2</sub> S	Ind	N.S.I
<i>Escherichia coli</i>	B	-	-	+	+	+	-	-	-	+	-	-	+	80
<i>Klebsiella pneumonia</i>	B	-	-	+	+	+	+	+	-	+	+	-	-	21
<i>Pseudomonas sp.</i>	B	-	+	+	-	-	+	-	-	-	-	-	-	17
<i>Salmonella sp.</i>	B	-	-	+	+	-	+	-	-	-	-	+	-	0
<b>Total</b>														118

F: Form, G: Gram, Ox: Oxidase; Cat: Catalase; Glu: Fermentation of Glucose; Lac: Fermentation Lactose; Cit: Citrate, Reduction of Nitrates to Nitrites; Ur: Urease; TDA: Tryptophan Deaminase; ONPG: Ortho-Nitro-Phényl-Galactopyranoside; Vp: Voges-Proskauer; H<sub>2</sub>S: Production of the Gas; Ind: Training of Indole; N.S.I: Number of Isolated Strains.

**Table 2:** Biochemical characteristics of the Strains *Escherichia coli*, *Klebsiella pneumonia*, *Pseudomonas sp.*, and *Salmonella sp.* isolated of turkey meat costs.

*Salmonella*. The *E. coli* strain was found in all points of sale with a total dominance (67.8%), followed by *K. pneumonia* with a prevalence of 17.8%, then *Pseudomonas sp.* with a rate of 14.4%. While the *Salmonella* has been absent.

These results suggest that there may be a lack of strategies of adequate control of turkey meat. Such inadequate controls can include the absence of good hygiene practices during the slaughter, transport, storage and distribution.

## References

- Foreign Intelligence Surveillance Act (FISA) (2015) Interprofessional Federation in the poultry sector-poultry sector in figures.
- Dennai N, Kharrattib B, El Yachoui M (2001) Appreciation of the microbiological quality of the carcasses of cattle freshly slaughtered. *Ann Vet Med* 145: 270-274.
- Pit J, Cappelier JM, Laroche M, Fradin N, Giraudet K, et al. (2006) Beef and veal: an analysis of biological hazards for the consumer applied to the slaughterhouse. *Meeting Ruminant Research* 13: 411-414.
- El-Hadef El, Okki S, ElGroud R, Kenana H, Quessy S (2005) Evaluation of the superficial contamination of carcasses beef/veal and sheep meat sectors from the municipal slaughterhouse of Constantine in Algeria. *Can Vet J* 46: 638-640.
- Cohen N, Ennaji H, Hassar M, Karib H (2006) The bacterial quality of red meat and offal in Casablanca (Morocco). *Mol Nutr Food Res* 50: 557-562.
- Anonymous (2005) Morocco foodborne disease outbreaks, searchable data 2000-2005. *Yearly Reports* 2000, 2001, 2002, 2003, 2004.
- NM 08.0.117 (2007) Food microbiology: Preparation of samples, the suspension mother and decimal dilutions in view of the Microbiological Examination-Part 2: Specific rules for the preparation of meat and meat products.
- NM 08.0.184 (2012) Enumeration of micro-organisms by counting the colonies obtained at 30°C after seeding by the method spiral.
- NM 08.0.124 (2012) Enumeration of thermotolerant coliforms by counting colonies obtained at 44°C.
- NM 08.0.127 (2012) Horizontal method for the research and the enumeration of *Escherichia coli* presumed-TECHNICAL The number the more likely.
- Brisse S, Grimont F, Grimont PAD (2006) The genus *Klebsiella* prokaryotes. *The Prokaryotes* 6: 159-196.
- NM 08.6.104 (2006) Enumeration of *Pseudomonas sp.*
- No. 624-04 (2004) Joint Decree of the Minister of Agriculture and Rural Development, the Minister of Health and the Minister of Industry, Trade and Telecommunications, of 17 Safar 1425 (8 April 2004) relating to microbiological standards which must be met by the animal foodstuffs or animal origin.
- Amara A, Badoum M, Faid M, Bouzoubaa K (1994) Microbial contamination of poultry slaughtered in traditional shops in Morocco. *Sci Int* 12: 323-327.
- Cohen N, Ennaji H, Bouchrif B, Hassan M, Karib H (2007) Comparative study of microbiological quality of raw poultry meat at various seasons and for different slaughtering processes in Casablanca (Morocco). *I Appl Poult Res* 16: 502-508.
- Chaiba A, Rhazi Filali F (2011) Impact of slaughter operations in the traditional killings on the bacteriological quality of poultry meat in Meknès (Morocco). *Tropicultura* pp: 161-167.
- Patyal A, Gangil R., Singh PK, Mathur KN, Sudan V (2012) Bacteriological quality of market chicken meat in Jaipur city. Department of Veterinary Public Health and Epidemiology Apollo College of Veterinary Medicine, Agra Road, Jaipur. *J Vet Public Health* 10: 45-48.
- El Allaoui A, Rhazi Filali F, Oumokhtar B (2013) Prevalence and Antibiogram Study of *Escherichia coli* and *Staphylococcus aureus* in turkey meat in Morocco. *Anal Pharmaceut Acta* 4: 270.
- Gupta N, Gupta S (2009) Microbial Quality of chicken meat samples collected from Durg (CG). *J Vet Public Health* 7: 79-81.
- Iroha IR, Ugbo EC, Ilang DC, Oji AE, Ayogu TE (2011) Bacteria Contamination of raw meat sold in Abakaliki, Ebonyi State Nigeria. *J Pub Health Epidemiol* 3: 49-53.
- Davis GS, Waits K, Nordstrom L, Weaver B, Aziz M, et al. (2015) Intermingled *Klebsiella pneumonia* populations between retail meats and human urinary tract infections. *Clin Infect Dis Adv*.
- Neslihan G, Sumru C, Emel Y (2011) Virulence properties of extended spectrum  $\beta$ -lactamase-producing *Klebsiella* species in meat samples. *J Food Prot* 74: 559-564.
- Cossi MVC, DeAlmeida MV, Dias MR, Pinto PSD, Nero LA (2012) Comparison of destructive and nondestructive sampling techniques of retail chicken carcasses for enumeration of hygiene indicator microorganisms. *Food Protect* 75: 29-33.
- Ould Abeid A, Mennane Z, Oudda H, Ouhsine M (2015) Microbiological study and identification of the strains isolated from the fish (*Mugil cephalus*) dried-pounding "Lekhli" (Microbiological study and identification of strains isolated from the fish (*Mugil cephalus*) dried-pounded Lekhli. *J Mater Approx Sci* 6: 1142-1146.
- Arnaut-Rollier I, De Zutter L, Hoof JV (1999) Identities of the *Pseudomonas* spp. in flora from chilled chicken. *Int J Food Microbiol* 48: 87-96.
- Asian Family Support Services of Austin (AFSSA) (2008) Opinion of the French Food Safety Agency relating to the transport of the carcasses of pork not having reached the regulatory temperature at the outlet of the Slaughterhouse.
- Salifou CFA, Boko KC, Attakpa YE, Agossa R, Ogbankota I, et al. (2013) Evaluation of the bacteriological quality of fresh meat of bovine animals slaughtered in the slaughterhouses of Cotonou Porto-Novo during the distribution chain. *J Anim Plant Sci* 17: 2567-2579.
- Gill CO, Landers C (2003) Effects of spray-cooling processes on the microbiological conditions of contaminated beef carcasses. *J Food Protect* 66: 1247-1252.
- Bornert G (2000) Chicken without *Salmonella*: Myth or Reality? *Med Review Vet* 151: 1083-1094.
- Bennani S, Berrada BS, Aabouch M, Ouali E, Lalami A (2016) Evaluation of the hygienic quality of meat and some meat products collected from the city of Fez, Morocco. *Int J Innov Appl Stud* 15: 547-554.
- Karraouan B, Fassouane A, EL Ossmani H, Cohen N, Charafeddine O, et al. (2010) Prevalence and genes for virulence of *Salmonella* isolated from raw ground meats of Turkey in Casablanca (Morocco). *Med Review Vet* 3: 127-132.
- Chaiba A (2011) Impact of practices of production of chicken flesh to Meknes on the bacteriological quality, the antibiorésistances and antibiotic residues in the avian products finished, Doctoral thesis national, University Moulay Ismail, Faculty of Sciences of Meknes, Morocco.
- Beli E, Telo A, Duraku E (2001) *Salmonella* serotypes isolated from turkey meat in Albania. *Int J Food Microbiol* 63: 165-167.

34. Jordan E, Egan J, Dullea C, Ward I, Mc-Gillicudy K, et al. (2006). Salmonella Surveillance in Raw and cooked meat and meat products in the 98 Republic of Ireland from 2002 to 2004. Int J Food Microbiol 112: 66-70.
35. Zhao C, GE B, Villena JD, Sulder R, Yeh E, et al. (2001) Prevalence of *Campylobacter* spp., *Escherichia coli*, and *Salmonella* serovars in retail chicken, turkey, pork and beef from Grater Washington, D.C, Area. Approx Appl Microbiol 67: 5431-5436.

**Citation:** Jaber H, Ijoub R, Zaher A, Chakit M, Rhaiem N, et al. (2017) Microbiological Study of Turkey Meat Marketed in Kenitra (North-oust of Morocco). J Nutr Food Sci 7: 620. doi: [10.4172/2155-9600.1000620](https://doi.org/10.4172/2155-9600.1000620)

### OMICS International: Open Access Publication Benefits & Features

#### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

#### Special features:

- 700+ Open Access Journals
- 50,000+ Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>