

Molecular Basis for Mercury-Induced Alteration in Endothelial Function: NO and its Modulators

Omanwar S¹, Saidullah B¹ and Fahim M²¹School of Sciences, Indira Gandhi National Open University, New Delhi, India²Hamdard Institute of Medical Science and Research, Hamdard University, New Delhi, India

Abstract

Epidemiological and animal studies have suggested a strong association between the environmental and occupational exposure to mercury and risk of cardiovascular diseases (CVD). One of the triggering factors of CVD is endothelial dysfunction. The endothelium can evoke relaxations and contractions of the underlying smooth muscle, by releasing vasoactive agents. Nitric oxide (NO), formed by endothelial NO synthase (eNOS), is the best characterized endothelium derived relaxing factor (EDRF). The release of NO is down regulated/upregulated by factors like oxidative stress, estrogen and diseases like diabetes and hypercholesterolemia, etc. The inhibition/activation of eNOS by mercury, affecting the NO release is one of the proposed mechanisms for mercury-induced vascular diseases. In addition, during exposure to mercury, overproduction of reactive oxygen species (ROS) can occur, resulting in oxidative stress, which is another major risk factor for endothelial dysfunction. In this article, molecular basis for mercury-mediated alteration in endothelium derived vasodilator (NO) and factors modulating the release of NO are being reviewed.

Background

For many years, the vascular endothelium, a monolayer of cells covering the vascular lumen was thought to be relatively inert. It is now recognized, as metabolically active with important paracrine, endocrine and autocrine functions, requisite for the maintenance of vascular homeostasis under physiological conditions [1,2]. The endothelium plays a vital role in the regulation of vascular tone, controlling tissue blood flow and inflammatory responses and maintaining blood fluidity [3-5]. Endothelium releases various vasoactive substances: vasodilators (NO, prostacyclin, endothelium-derived hyperpolarizing factor (EDHF), bradykinin, adrenomedullin, C-natriuretic peptide) and vasoconstrictors (endothelin-1, angiotensin-II, thromboxane A₂, prostaglandins, hydrogen peroxide (H₂O₂) and free radicals) which help in regulating the vascular tone [6]. As a major regulator of vascular homeostasis, the endothelium maintains the balance between vasodilation and vasoconstriction and disturbing this tightly regulated equilibrium leads to endothelial dysfunction [7].

There is a growing data (epidemiological and animal studies) associating mercury exposure with higher risk of CVD which has been thoroughly documented [8-18]. One of the events that triggers development of cardiovascular disease is functional disruption of the vascular endothelium. Man has been exposed to mercury for centuries as it occurs naturally in the environment, and anthropogenic activities causes the release of this element into the environment, leading to pollution of air, water, and soil [19,20]. Its exposure is the second-most common cause of heavy metal poisoning. Toxicity from mercury is associated with *in-vivo* oxidative stress. The loss of endothelial function due to oxidative stress is one of the most commonly observed vascular effects of mercury exposure [21-24].

This review investigates and appraises the impact of mercury exposure on endothelium derived vasodilator, NO, highlighting the molecular basis for mercury-mediated alteration on endothelial function and modulators of NO.

Vascular Function, NO and Mercury

Nitric oxide

The L-arginine-NO pathway is thought to be the most important enzymatic vasodilator source. In addition to its function as a vasodilator, NO released from endothelial cells is also a potential inhibitor of the aggregation and adhesion of platelets to the vascular wall.

The endothelial cells synthesize NO using substrate L-arginine with help of an enzyme eNOS (endothelial nitric oxide synthase). The by-product of the reaction, l-citrulline, inhibits arginase II, which cause hydrolysis of arginine [7]. eNOS is an NADPH-dependent oxygenase that requires tetrahydrobiopterin, flavin adenine dinucleotide (FAD) and flavin mononucleotide (FMN) as cofactors [25,26]. In endothelial cells, the enzyme is localized preferentially in caveolae following post-translational acylation [27,28]. NO is formed when eNOS is stimulated by agonists such as bradykinin, acetylcholine and shear stress. Stimulation of endothelial cells causes the dissociation of the caveolin/NOS complex. This dissociation of caveolin/NOS causes binding and activation of Ca²⁺/calmodulin and NO synthesis by eNOS [27,28]. During, shear stress activation of NOS is Ca²⁺-independent and protein tyrosine kinase-dependent [29]. NO relaxes the vascular smooth muscle by stimulating of soluble guanylate cyclase which results in an increased formation of cyclic GMP (cGMP) [30]. The cGMP activates cGMP-dependent protein kinase which results in an increased extrusion of Ca²⁺ from the cytosol in vascular smooth

***Corresponding author:** Omanwar S, School of Sciences, Indira Gandhi National Open University, New Delhi, India, Tel: 91-011-29572818; E-mail: swatiomanwar@gmail.com

Received December 03, 2015; **Accepted** December 10, 2015; **Published** December 15, 2015

Citation: Omanwar S, Saidullah B, Fahim M (2015) Molecular Basis for Mercury-Induced Alteration in Endothelial Function: NO and its Modulators. Cardiovasc Pharm Open Access 4: 167. doi:10.4172/2329-6607.1000167

Copyright: © 2015 Omanwar S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

muscle, and the inhibition of the contractile machinery [31]. Cyclic GMP-dependent protein kinase also phosphorylates K^+_{ATP} channels which culminates into hyperpolarization and thereby inhibition of vasoconstriction [32]. In certain arteries, NO directly activates K^+_{ATP} channels independently of cGMP [33] (Figure 1).

Mercury exposure and NO signaling

NO release from vascular endothelium is affected by mercury exposure (Figure 2) was reported for the first time by Golpon and co-workers [34]. They observed that in the isolated aortic tissue, low concentrations of mercury produced an endothelium dependent vasorelaxation which was blocked by the NOS inhibitor L-NAME. However at higher concentrations, mercury altered the structure and function of vascular endothelium and vasoconstriction was observed [34]. Further studies from our group on isolated aortic rings validated that mercury produces a dual response: vasoconstriction at high concentration and vasorelaxation at lower concentrations. L-NAME and glybenclamide block the vasorelaxation produced at low concentrations of mercury suggesting that mercury acts through activation of NOS and K^+_{ATP} channel [35]. It may be hypothesized that mercury stimulates NOS, which forms NO and consequently the synthesis of cGMP. cGMP increases extrusion of Ca^{2+} from the cytosol in vascular smooth muscle, causing vasorelaxation [35]. The cGMP and NO also directly phosphorylate K^+_{ATP} channels. K^+_{ATP} channels induce hyperpolarization, which results also in vasorelaxation [31,32]. Decreased acetylcholine (ACh) induced vasorelaxation has been observed when rat aortic rings were exposed to mercury chloride (10^{-5} M) and methyl mercury chloride (10^{-5} M) separately *in-vitro* suggesting endothelial dysfunction. However, an increased ACh induced vasodilation was observed in rat aortic rings exposed to low concentrations of mercury chloride (10^{-9} M), and methyl mercury chloride (10^{-9} M), separately indicating modulation of the endothelial function. The NOS inhibitor, L-NAME significantly reduced the ACh induced vasorelaxation in mercury chloride (10^{-9} M) and methyl mercury chloride (10^{-9} M) exposed aortic rings, in comparison to mercury chloride (10^{-5} M) and methyl mercury chloride (10^{-5} M) exposed aortic rings. Mercury induces increased/decreased production of NO depending on the dose of exposure [36]. The above experiments by Golpon and our group were performed on isolated vascular tissue where other physiological mechanisms do not play any role. In an animal model many factors get involved, the effects of mercury on vascular function especially those on NO and factors modulating the release of NO are interesting.

Mercury and modulation of protective role of Nitric oxide

The ability of the endothelium to release NO can be upregulated/downregulated by several factors. NO is upregulated by shear stress, estrogen, insulin, adiponectin, aldosterone, arginine and ω unsaturated fatty acid, etc. NO is downregulated by oxidative stress, aging, obesity, hypercholesterolemia etc [7]. Several studies have shown a connection between mercury exposure and factors which may upregulate and down regulate the release of endothelium from NO.

Reactive oxygen species

Several enzymes in the endothelial cells namely NADPH oxidase, xanthine oxidase, cyclooxygenase and eNOS itself generate super-oxide anions [7]. eNOS, generates super-oxide anions when it is uncoupled by lack of substrate (L-arginine) or shortage of the essential co-factor tetrahydrobiopterin (BH4) [37]. Super-oxide anions are dismutated to hydrogen peroxide (H_2O_2) by superoxide dismutase (SOD). H_2O_2 contribute to endothelium-dependent relaxations or is broken down

by catalase [3,38]. However, superoxide anions also react avidly with NO resulting in the formation of a potent vasoconstrictor peroxynitrite [37,39,40]. Superoxide anion also reduces the bioavailability of NO resulting in reduced endothelium-dependent relaxations [39] (Figure 1). Antioxidants have shown to acutely improve endothelial responses *in-vitro* and *in-vivo* both in animals and humans [41-44].




Figure 1: Schematic representation of possible mechanisms (only those mechanism affected by mercury exposure) by which production of nitric oxide is regulated in endothelial cells. Nitric oxide is produced through enzymatic conversion of L-arginine by nitric oxide synthase (eNOS). Insulin and estrogen cause upregulation of eNOS by Akt phosphorylation. Oxidative stress causes downregulation of eNOS due to low tetrahydrobiopterin (BH₄). The by-product of the eNOS reaction, L-citrulline (L-Cit), inhibits arginase which hydrolyses arginine.




Figure 2: Schematic representation of effect of low/high mercury exposure on endothelium-derived vasoactive factors. In response to low mercury exposure, an increase in the eNOS activity/expression causes an increase in NO. In response to high mercury exposure decrease in the eNOS activity/expression causes decrease in NO with persistence or upregulation of EDHF. Mercury exposure also causes oxidative stress, hypercholesterolemia, decrease in insulin and affects estrogen.

There are a number of studies revealing that mercury exposure generates oxygen radicals, with subsequent oxidative damage in several organs and systems as well as alters the antioxidant defense system in the cells [45-49]. Cell toxicity caused by heavy metal ions is attributed to oxidative and nitrosative stress, defined as an excess of oxidants over antioxidants. Macromolecules in cells are damaged by mercury-induced production of oxygen and nitrogen-containing free radicals (oxidants) and/or metal-induced depletion of the antioxidant defences. The vascular endothelium is very sensitive to oxidative stress [50-52]. Although multiple processes may lead to endothelial damage, the generation of oxygen-derived free radicals and subsequent lipid peroxidation may be one of the key components in the cascade of events.

On exposure to mercury, vascular Endothelial cells (ECs) in culture, generates superoxide anions (oxidative stress) in ECs. Mercury induced oxidative stress in ECs is caused by the activation of phospholipase D (PLD) which results in generation of, 2-diacylglycerol (DAG), a second messenger for vasoconstriction [53]. Similarly in Bovine Pulmonary Artery Endothelial Cells (BPAECs), mercury-induced PLD activity is attenuated by L-type calcium channel blockers demonstrating the importance of calcium and calmodulin in the regulation of mercury-induced PLD activation and the protective action of L-type calcium channel blockers against mercury cytotoxicity in vascular ECs [54]. In bovine pulmonary artery endothelial cell monolayers, it has been reported that mercury ions induce oxidative stress through depletion of GSH and inactivation of thiol enzymes [55].

Studies on isolated aortic rings indicate that the dual response (vasoconstriction at high concentration and vasorelaxation at lower concentrations) produced by mercury is ameliorated by antioxidants SOD and catalase and L-type Ca channel [35].

In another *in-vitro* mercury exposure (HgCl_2 , 6 nM) study, mercury induced modulation of vascular reactivity was observed due to increased release of ROS derived from NADPH oxidase which culminated into reduced bioavailability of NO [56]. Mercury causes contraction of the vascular smooth muscle in isolated rat vascular bed which is mediated by the formation of superoxide anions and by reduction in the endothelial vasodilator activity [57].

Acute and chronic exposure to mercury in rats causes an increase in vascular O_2^- production, plasma malondialdehyde levels and total antioxidant [35,55,57-61]. Low dose of mercury exposure causes endothelial dysfunction and alters the coronary vascular reactivity. This alteration is partially, due to increase in both NOX-1 and NOX-4 subunits suggesting the involvement of NADPH oxidase resulting in increased ROS production. A decrease in antioxidant defenses could also be a contributory factor to the increased superoxide production observed after mercury treatment [60]. In contrast several authors have reported augmented antioxidant defenses to protect cells against the increased oxidative stress after acute and chronic mercury exposure [55,56,62,63].

The interplay between NO and oxidative stress occurs on mercury exposure as free radical scavengers along with L-NAME blocks the mercury induced vasorelaxant or vasoconstrictor response in isolated aortic rings. SOD β catalase blocks the ROS mediated vasoconstrictor response and L-NAME blocks the NO-mediated vasorelaxant response. Mercury evokes the production of both NO and ROS, and it is their percentage that produces the response. If ROS produced is more than NO, peroxynitrite is formed and a vasoconstriction is produced [35].

This interaction of NO and oxidative stress was further validated by

an acute exposure study of methyl mercury chloride (5 mg/kg; po.) in rats. Oxidative stress was produced along with an increase in serum NO levels. A significant increase in the acetylcholine vasodilator response in isolated aortic rings from mercury exposed rats was observed. This effect was mediated by increased production of NO, because of stimulation of eNOS. It is interesting to note that this increase occurred even when there was oxidative stress, suggesting a state of inclination towards NO [63] (Figure 2).

Contrary, to above an opposite interaction between NO and oxidative stress was observed in a chronic study on healthy wistar rats exposed to inorganic mercuric chloride in drinking water for 30 days. Oxidative stress accompanied with increased NO levels and endothelial dysfunction, was observed. Even though there was an increase in the serum NO level, endothelial dysfunction was detected. As there was a significant increase in free radical production, the free radicals must have interacted with NO and reduced the bioavailability of NO resulting in endothelial dysfunction. The EDHF pathway was relatively resistant to mercury exposure and oxidative stress, suggesting that there may be an up-regulation of K_{ATP}^+ channels in order to maintain circulation to compensate the attenuated NO-mediated vasodilatation (Figure 2) [58].

The evidence available suggests that antioxidants may play an important role in abating some health hazards (endothelial dysfunction) of mercury exposure.

Insulin

In-vitro studies on insulin have recorded an enhancement in the expression of eNOS in native endothelial cells *in-vitro* [64] and facilitation of NO-dependent vasorelaxations *in-vivo* [65-67] suggesting that NO is upregulated in presence of insulin (Figure 1). Decrease in insulin may result in attenuation in endothelium release of NO. There is a significant correlation between diabetes (impaired/decreased insulin secretion) and total hair mercury levels [68-71]. Follow up studies of patients with Minamata disease (organic mercury poisoning) in Japan, have also shown an increased incidence of diabetes 1 due to pancreatic β -cells dysfunction [72]. Similarly, *in-vitro* studies have furthermore shown a mercury dependent decrease in the function and viability of pancreatic β -cells due to increased oxidative stress, causing impaired insulin secretion [73]. One of the reasons of endothelial dysfunction in mercury exposure worth characterizing may be decrease in insulin levels leading to downregulation of eNOS and endothelial dysfunction (Figure 2).

Estrogen

The potentiating effect of estrogens on endothelium-dependent relaxations involves both genomic and non-genomic effects [74-76]. It depends presumably both on a reduction in oxidative stress leading to an increased bioavailability of NO and an increased responsiveness of the vascular smooth muscle cells to vasodilator stimuli [76-79] (Figure 1). Rowland et al. (1994) reported that 418 women with high exposure to mercury (i.e., female dental assistants) were less fertile than unexposed controls [80]. In a study by Baranski and Szymczyk (1973), female rats exposed via inhalation to metallic mercury (at an average of 2.5 mg/m³, 6 hours a day, 5 days a week for 21 days) experienced longer estrous cycles than unexposed animals [81]. Decrease in estrogen level after mercury exposure renders testis more susceptible to oxidative damage leading to its functional inactivation [82]. Mercuric chloride also exhibits estrogen-like effect through binding and activating estrogen receptor (ER) [83]. It is likely that the potentiating effect of mercury on NO release presumably may be due to its estrogen like

effect and further studies are needed to test the hypothesis (Figure 2).

Omega-3 Unsaturated Fatty Acid

Consumption of fish has always been considered beneficial for the health of the cardiovascular system. The benefits of fish are mainly due to its content of omega-3 long chain polyunsaturated fatty acids (n-3 PUFA), including eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). Fish consumption is associated with decreased circulating biomarkers of endothelial dysfunction and inflammation [84], and flow mediated dilation values seem to improve after n-3 PUFA intake [85]. The chronic intake of Omega-3-unsaturated fatty acids potentiates the endothelium-dependent relaxations of coronary arteries [86-88]. Omega-3 fatty acid might attenuate Mercury chloride-induced toxicity by improving antioxidant status and acute phase response in mice [89]. There are a contrary number of studies, that say the consumption of fish contaminated with mercury, causes high mercury levels in blood, hair, urine and toe nail, which diminishes/does not diminish its cardio protective effect [90-94]. Omega 3-fatty acids attenuates vascular endothelial dysfunction, by activating eNOS through opening of K^+_{ATP} channels in sodium arsenite exposed rats [95]. As mercury is known to modulate eNOS and also K^+_{ATP} channels the beneficial effect of habitual consumption of high amounts of fish is associated with better endothelial function despite higher serum concentrations of mercury [96].

Hypercholesterolemia

Hypercholesterolemia reduces endothelium-dependent relaxations due to increased oxidative stress leading to a reduced bioavailability of NO [97-100]. Long-term MeHg treatment induces dyslipidemia, characterized by increased serum cholesterol levels in mice [101]. In a study done on 274 school children, an increased level of mercury in urine was associated with elevated cholesterol level which is a known risk factor of myocardial infarction, coronary disease, and cardiovascular disease [102]. Hypercholesterolemia reduces eNOS activity and may act as a triggering factor of mercury induced reduction in endothelium dependent relaxation (Figure 2).

Conclusion

Vascular endothelial cells respond to low concentration of mercury by releasing NO, which relaxes the vascular smooth muscle that surrounds them. High concentration of mercury causes decrease in endothelial NO and vasoconstriction. Decrease/increase in release of NO on mercury exposure is majorly regulated by superoxide anions and minorly may be by insulin, estrogen, omega-3 unsaturated fatty acid and hypercholesterolemia. It emerges that a fragile equilibrium exists between NO released and superoxide anions generated by mercury exposure. When this balance leans towards NO, enhanced endothelial function is detected. When the equilibrium inclines in favour of oxidative stress, endothelial dysfunction is observed. The endothelial dysfunction caused by the imbalance between the production of NO and oxidative stress on mercury exposure can be compensated by EDHF-mediated vasorelaxation. Mercury exposure affects insulin (decrease/impaired) and causes hypercholesterolemia, downregulating the release of NO. Mercury has estrogen like affect and may be increasing the release of NO. The beneficial effects of omega-3 unsaturated fatty acid on mercury exposure and endothelial vascular release of NO are a point of contention. NO signalling mechanism and oxidative stress play a vital function in the mercury-induced cardiovascular diseases in the populations exposed to mercury.

References

1. Vallance P (2001) Importance of asymmetrical dimethyl arginine in cardiovascular risk. *Lancet* 358: 2096-2097.
2. Bonetti PO, Lerman LO, Lerman A (2003) Endothelial dysfunction: a marker of atherosclerotic risk. *Arterioscler Thromb Vasc Biol* 23: 168-175.
3. Félétou M, Vanhoutte PM (2006) Endothelial dysfunction: a multifaceted disorder (The Wiggers Award Lecture). *Am J Physiol Heart Circ Physiol* 291: H985-1002.
4. Moncada S, Higgs EA (2006) Nitric oxide and the vascular endothelium. *Handb Exp Pharmacol* 176 Pt 1: 213-254.
5. Félétou M (2011) The endothelium: Part 1: Multiple functions of the endothelial cells—focus on endothelium-derived vasoactive mediators, San Rafael (CA): Morgan and Claypool Life Sciences.
6. McGuire JJ, Ding H, Triggle CR (2001) Endothelium-derived relaxing factors: a focus on endothelium-derived hyperpolarizing factor(s). *Can J Physiol Pharmacol* 79: 443-470.
7. Vanhoutte PM, Shimokawa H, Tang EH, Feletou M (2009) Endothelial dysfunction and vascular disease. *Acta Physiol (Oxf)* 196: 193-222.
8. Omanwar S, Fahim M (2015) Mercury Exposure and Endothelial Dysfunction: An Interplay Between Nitric Oxide and Oxidative Stress. *Int J Toxicol* 34: 300-307.
9. Salonen JT, Seppänen K, Nyyssönen K, Korpela H, Kahvanen J, et al. (1995) Intake of mercury from fish, lipid peroxidation, and the risk of myocardial infarction and coronary, cardiovascular, and any death in eastern Finnish men. *Circulation* 9: 645-655.
10. Virtanen JK, Voutilainen S, Rissanen TH, Mursu J, Tuomainen TP, et al. (2004) Mercury, fish oils, and risk of acute coronary events and cardiovascular disease, coronary heart disease, and all-cause mortality in men in eastern Finland. *Arterioscler Thromb Vasc Biol* 25: 228-233.
11. Houston MC (2007) The role of mercury and cadmium heavy metals in vascular disease, hypertension, coronary heart disease, and myocardial infarction. *Altern Ther Health Med* 13: S128-133.
12. García Gómez M, Boffetta P, Caballero Klink JD, Español S, Gómez Quintana J (2007) [Cardiovascular mortality in mercury miners]. *Med Clin (Barc)* 128:766-771.
13. Choi AL, Weihe P, Budtz-Jørgensen E, Jørgensen PJ, Salonen JT, et al. (2009) Methylmercury exposure and adverse cardiovascular effects in Faroese whaling men. *Environ Health Perspect* 117: 367-372.
14. Trakhtenberg IM (1974) Chronic effects of mercury on organisms: the micromercurialism phenomenon on mercury handlers. Chap VI: 109-134.
15. Snodgrass W, Sullivan JB Jr, Rumack BH, Hashimoto C (1981) Mercury poisoning from home gold ore processing. Use of penicillamine and dimercaprol. *JAMA* 246: 1929-1931.
16. Barregård L, Sällsten G, Järholm B (1990) Mortality and cancer incidence in chloralkali workers exposed to inorganic mercury. *Br J Ind Med* 47: 99-104.
17. Boffetta P, Sällsten G, Garcia-Gómez M, Pompe-Kirn V, Zaridze D, et al. (2001) Mortality from cardiovascular diseases and exposure to inorganic mercury. *Occup Environ Med* 58: 461-466.
18. Sibirerud RL (1990) The relationship between mercury from dental amalgam and the cardiovascular system. *Sci Total Environ* 99: 23-35.
19. Clarkson TW, Magos L, Myers GJ (2003) The toxicology of mercury—current exposures and clinical manifestations. *N Engl J Med* 349: 1731-1737.
20. Bhan A, Sarkar NN (2005) Mercury in the environment: effect on health and reproduction. *Rev Environ Health* 20: 39-56.
21. Huang YL, Cheng SL, Lin TH (1996) Lipid peroxidation in rats administered with mercuric chloride. *Biol Trace Elem Res* 52: 193-206.
22. Mahboob M, Shireen KF, Atkinson A, Khan AT (2001) Lipid peroxidation and antioxidant enzyme activity in different organs of mice exposed to low level of mercury. *J Environ Sci Health B* 36: 687-697.
23. Reus IS, Bando I, Andrés D, Cascales M (2003) Relationship between expression of HSP70 and metallothionein and oxidative stress during mercury chloride induced acute liver injury in rats. *J Biochem Mol Toxicol* 17: 161-168.

24. Chen C, Qu L, Li B, Xing L, Jia G, et al. (2005) Increased oxidative DNA damage, as assessed by urinary 8-hydroxy-2'-deoxyguanosine concentrations, and serum redox status in persons exposed to mercury. *Clin Chem* 51: 759-767.
25. Moncada S, Rees DD, Schulz R, Palmer RM (1991) Development and mechanism of a specific supersensitivity to nitrovasodilators after inhibition of vascular nitric oxide synthesis in vivo. *Proc Natl Acad Sci U S A* 88: 2166-2170.
26. Förstermann U, Closs EI, Pollock JS, Nakane M, Schwarz P, et al (1994) Nitric oxide synthase isozymes. Characterization, purification, molecular cloning, and functions. *Hypertension* 23: 1121-1131.
27. Michel JB, Feron O, Sacks D, Michel T (1997) Reciprocal regulation of endothelial nitric-oxide synthase by Ca^{2+} -calmodulin and caveolin. *J Biol Chem* 272: 15583-15586.
28. Feron O, Saldana F, Michel JB, Michel T (1998) The endothelial nitric-oxide synthase-caveolin regulatory cycle. *J Biol Chem* 273: 3125-3128.
29. Ayajiki K, Kindermann M, Hecker M, Fleming I, Busse R (1996) Intracellular pH and tyrosine phosphorylation but not calcium determine shear stress-induced nitric oxide production in native endothelial cells. *Circ Res* 78: 750-758.
30. Rapoport RM, Murad F (1983) Agonist-induced endothelium-dependent relaxation in rat thoracic aorta may be mediated through cGMP. *Circ Res* 52: 352-357.
31. Lincoln TM, Komalavilas P, Cornwell TL (1994) Pleiotropic regulation of vascular smooth muscle tone by cyclic GMP-dependent protein kinase. *Hypertension* 23: 1141-1147.
32. Bolotina VM, Najibi S, Palacino JJ, Pagano PJ, Cohen RA (1994) Nitric oxide directly activates calcium-dependent potassium channels in vascular smooth muscle. *Nature* 368: 850-853.
33. Ignarro LJ, Napoli C, Loscalzo J (2002) Nitric oxide donors and cardiovascular agents modulating the bioactivity of nitric oxide: an overview. *Circ Res* 90: 21-28.
34. Golpon HA, Püchner A, Barth P, Welte T, Wichert P, et al (2003) Nitric oxide-dependent vasorelaxation and endothelial cell damage caused by mercury chloride. *Toxicology* 192: 179-188.
35. Omanwar S, Saidullah B, Ravi K, Fahim M (2014) Vasorelaxant effects of mercury on rat thoracic aorta: the nitric oxide signaling mechanism. *Hum Exp Toxicol* 33: 904-910.
36. Omanwar S, Fahim M (2015) Endothelial dysfunction of rat aorta after exposure to mercury is dependent on reactive oxygen species and nitric oxide. *Proceedings of The Physiological Society*.
37. Kojda G, Harrison D (1999) Interactions between NO and reactive oxygen species: pathophysiological importance in atherosclerosis, hypertension, diabetes and heart failure. *Cardiovasc Res* 43: 562-571.
38. Matoba T, Shimokawa H, Nakashima M, Hirakawa Y, Mukai Y, et al. (2000) Hydrogen peroxide is an endothelium-derived hyperpolarizing factor in mice. *J Clin Invest* 106: 1521-1530.
39. Miyagawa K, Ohashi M, Yamashita S, Kojima M, Sato K, et al. (2007) Increased oxidative stress impairs endothelial modulation of contractions in arteries from spontaneously hypertensive rats. *J Hypertens* 25: 415-421.
40. Macarthur H, Westfall TC, Wilken GH (2008) Oxidative stress attenuates NO-induced modulation of sympathetic neurotransmission in the mesenteric arterial bed of spontaneously hypertensive rats. *Am J Physiol Heart Circ Physiol* 294: H183-189.
41. Aubin MC, Carrier M, Shi YF, Tardif JC, Perrault LP (2006) Role of probucol on endothelial dysfunction of epicardial coronary arteries associated with left ventricular hypertrophy. *J Cardiovasc Pharmacol* 47: 702-710.
42. Liu YH, You Y, Song T, Wu SJ, Liu LY (2007) Impairment of endothelium-dependent relaxation of rat aortas by homocysteine thiolactone and attenuation by captopril. *J Cardiovasc Pharmacol* 50: 155-161.
43. Kanani PM, Sinkey CA, Browning RL, Allaman M, Knapp HR, et al. (1999) Role of oxidant stress in endothelial dysfunction produced by experimental hyperhomocyst(e)inemia in humans. *Circulation* 100: 1161-1168.
44. Taddei S, Virdis A, Ghiadoni L, Salvetti G, Bernini G, et al. (2001) Age-related reduction of NO availability and oxidative stress in humans. *Hypertension* 38: 274-279.
45. Miller DM, Woods JS (1993) Urinary porphyrins as biological indicators of oxidative stress in the kidney. Interaction of mercury and cephaloridine. *Biochem Pharmacol* 46: 2235-2241.
46. Huang YL, Cheng SL, Lin TH (1996) Lipid peroxidation in rats administered with mercuric chloride. *Biol Trace Elem Res* 52: 193-206.
47. Mahboob M, Shireen KF, Atkinson A, Khan AT (2001) Lipid peroxidation and antioxidant enzyme activity in different organs of mice exposed to low level of mercury. *J Environ Sci Health B* 36: 687-697.
48. Wildemann TM, Siciliano SD, Weber LP (2015) The mechanisms associated with the development of hypertension after exposure to lead, mercury species or their mixtures differs with the metal and the mixture ratio. *Toxicology* 339: 1-8.
49. Rangel-Méndez JA, Arcega-Cabrera FE, Fargher LF, Moo-Puc RE (2015) Mercury levels assessment and its relationship with oxidative stress biomarkers in children from three localities in Yucatan, Mexico. *Sci Total Environ* 543(PtA): 187-196.
50. Stohs SJ, Bagchi D (1995) Oxidative mechanisms in the toxicity of metal ions. *Free Radic Biol Med* 18: 321-336.
51. Ercal N, Gurer-Orhan H, Aykin-Burns N (2001) Toxic metals and oxidative stress part I: mechanisms involved in metal-induced oxidative damage. *Curr Top Med Chem* 1: 529-539.
52. Pompella A, Visvikis A, Paolicchi A, De Tata V, Casini AF (2003) The changing faces of glutathione, a cellular protagonist. *Biochem Pharmacol* 66: 1499-503.
53. Peltz A, Sherwani SI, Kotha SR, Mazerik JN, O'Connor Butler ES, et al. (2009) Calcium and calmodulin regulate mercury-induced phospholipase D activation in vascular endothelial cells. *Int J Toxicol* 28: 190-206.
54. Mazerik JN, Mikkilineni H, Kuppusamy VA, Steinhour E, Peltz A, et al. (2007) Mercury activates phospholipase a(2) and induces formation of arachidonic Acid metabolites in vascular endothelial cells. *Toxicol Mech Methods* 17: 541-557.
55. Wolf MB, Baynes JW (2007) Cadmium and mercury cause an oxidative stress-induced endothelial dysfunction. *Biometals* 20: 73-81.
56. Lemos NB, Angeli JK, Faria Tde O, Ribeiro Junior RF, Vassallo DV, et al. (2012) Low mercury concentration produces vasoconstriction, decreases nitric oxide bioavailability and increases oxidative stress in rat conductance artery. *PLoS One* 7: e49005.
57. Da Cunha V, Souza HP, Rossoni LV, França AS, Vassallo DV (2000) Effects of mercury on the isolated perfused rat tail vascular bed are endothelium-dependent. *Arch Environ Contam Toxicol* 39: 124-130.
58. Omanwar S, Ravi K, Fahim M (2011) Persistence of EDHF pathway and impairment of the nitric oxide pathway after chronic mercury chloride exposure in rats: mechanisms of endothelial dysfunction. *Hum Exp Toxicol* 30: 1777-1784.
59. Wiggers GA, Peçanha FM, Briones AM, Pérez-Girón JV, Miguel M, et al. (2008) Low mercury concentrations cause oxidative stress and endothelial dysfunction in conductance and resistance arteries. *Am J Physiol Heart Circ Physiol* 295: H1033-H1043.
60. Jindal M, Garg GR, Mediratta PK, Fahim M (2011) Protective role of melatonin in myocardial oxidative damage induced by mercury in murine model. *Hum Exp Toxicol* 30: 1489-1500.
61. Rizzetti DA, Torres JG, Escobar AG, Peçanha FM, Santos FW et al. (2013) Apocynin prevents vascular effects caused by chronic exposure to low concentrations of mercury. *PLoS One* 8: e55806.
62. Harisa GI, Alanazi FK, El-Bassat RA, Malik A, Abdallah GM et al. (2012) Protective effect of pravastatin against mercury induced vascular cells damage: erythrocytes as surrogate markers. *Environ Toxicol Pharmacol* 34: 428-435.
63. Omanwar S, Saidullah B, Ravi K, Fahim M (2013) Modulation of vasodilator response via the nitric oxide pathway after acute methyl mercury chloride exposure in rats. *Biomed Res Int* 2013: 530603.
64. Fisslthaler B, Benzing T, Busse R, Fleming I (2003) Insulin enhances the expression of the endothelial nitric oxide synthase in native endothelial cells: a dual role for Akt and AP-1. *Nitric Oxide* 8: 253-261.
65. Lembo G, Iaccarino G, Vecchione C, Barbato E, Izzo R, et al. (1997) Insulin modulation of an endothelial nitric oxide component present in the alpha2- and beta-adrenergic responses in human forearm. *J Clin Invest* 100: 2007-2014.

66. Steinberg HO, Brechtel G, Johnson A, Fineberg N, Baron AD (1994) Insulin-mediated skeletal muscle vasodilation is nitric oxide dependent. A novel action of insulin to increase nitric oxide release. *J Clin Invest* 94: 1172-1179.
67. Taddei S, Virdis A, Mattei P, Natali A, Ferrannini E, et al. (1995) Effect of insulin on acetylcholine-induced vasodilation in normotensive subjects and patients with essential hypertension. *Circulation* 92: 2911-2918.
68. Nakagawa R (1995) Concentration of mercury in hair of diseased people in Japan. *Chemosphere* 30: 135-140.
69. Chang JW, Chen HL, Su HJ, Liao PC, Guo HR, et al. (2011) Simultaneous exposure of non-diabetics to high levels of dioxins and mercury increases their risk of insulin resistance. *J Hazard Mater* 185: 749-755.
70. Chen YW, Huang CF, Tsai KS, Yang RS, Yen CC, et al. (2006) Methylmercury induces pancreatic beta-cell apoptosis and dysfunction. *Chem Res Toxicol* 19: 1080-1085.
71. Uchino M, Tanaka Y, Ando Y, Yonehara T, Hara A, et al. (1995) Neurologic features of chronic minamata disease (organic mercury poisoning) and incidence of complications with aging. *J Environ Sci Health B* 30: 699-715.
72. Chen YW, Huang CF, Tsai KS, Yang RS, Yen CC, et al. (2006) Methylmercury induces pancreatic beta-cell apoptosis and dysfunction. *Chem Res Toxicol* 19: 1080-1085.
73. Tostes RC, Nigro D, Fortes ZB, Carvalho MH (2003) Effects of estrogen on the vascular system. *Braz J Med Biol Res* 36: 1143-1158.
74. Keung W, Vanhoutte PM, Man RY (2005) Nongenomic responses to 17beta-estradiol in male rat mesenteric arteries abolishes intrinsic gender differences in vascular responses. *Br J Pharmacol* 146:1148-55.
75. Miller VM, Duckles SP (2008) Vascular actions of estrogens: functional implications. *Pharmacol Rev* 60: 210-241.
76. Wassmann S, Bäumer AT, Strehlow K, van Eickels M, Grohé C et al. (2001) Endothelial dysfunction and oxidative stress during estrogen deficiency in spontaneously hypertensive rats. *Circulation* 103: 435-441.
77. Scott PA, Tremblay A, Brochu M, St-Louis J (2007) Vasorelaxant action of 17-estradiol in rat uterine arteries: role of nitric oxide synthases and estrogen receptors. *Am J Physiol Heart Circ Physiol* 293: H3713-3719.
78. Lee MY, Man RY (2003) The phytoestrogen genistein enhances endothelium-independent relaxation in the porcine coronary artery. *Eur J Pharmacol* 481: 227-232.
79. Han G, Ma H, Chintala R, Miyake K, Fulton DJ et al. (2007) Nongenomic, endothelium-independent effects of estrogen on human coronary smooth muscle are mediated by type I (neuronal) NOS and PI3-kinase-Akt signaling. *Am J Physiol Heart Circ Physiol* 293: H314-321.
80. Rowland AS, Baird DD, Weinberg CR, Shore DL, Shy CM et al. (1994) The effect of occupational exposure to mercury vapour on the fertility of female dental assistants. *Occup Environ Med* 51: 28-34.
81. Baranski B, Szymczyk I. (1973) Effects of mercury vapor upon reproductive functions of female white rats. *Med Pr* 24: 248.
82. Boujbiha MA, Hamden K, Guermazi F, Bouslama A, Omezzine A et al. (2010) Impairment of spermatogenesis in rats by mercuric chloride: involvement of low 17β-estradiol level in induction of acute oxidative stress. *Biol Trace Elem Res* 142: 598-610.
83. Zhang X, Wang Y, Zhao Y, Chen X (2008) Experimental study on the estrogen-like effect of mercuric chloride. *Biometals* 21: 143-150.
84. Van Bussel BC, Henry RM, Schalkwijk CG, Ferreira I, Feskens EJ, et al. (2011) Fish consumption in healthy adults is associated with decreased circulating biomarkers of endothelial dysfunction and inflammation during a 6-year follow-up. *J Nutr* 141: 1719-1725.
85. Xin W, Wei W, Li X (2012) Effect of fish oil supplementation on fasting vascular endothelial function in humans: a meta-analysis of randomized controlled trials. *PLoS One* 7: e46028.
86. Shimokawa H, Vanhoutte PM (1989) Dietary omega 3 fatty acids and endothelium-dependent relaxations in porcine coronary arteries. *Am J Physiol* 256: H968-973.
87. Shimokawa H, Lam JY, Chesebro JH, Bowie EJ, Vanhoutte PM (1987) Effects of dietary supplementation with cod-liver oil on endothelium-dependent responses in porcine coronary arteries. *Circulation* 76: 898-905.
88. Sekikawa A, Curb JD, Ueshima H, El-Saed A, Kadowaki T et al. (2008) ERA JUMP (Electron-Beam Tomography, Risk Factor Assessment Among Japanese and U.S. Men in the Post-World War II Birth Cohort) Study Group. Marine-derived n-3 fatty acids and atherosclerosis in Japanese, Japanese-American, and white men: a cross-sectional study. *J Am Coll Cardiol* 52:417-424.
89. Karapehliyan M, Ogun M, Kaya I, Ozen H, Deveci HA, et al. (2014) Protective effect of omega-3 fatty acid against mercury chloride intoxication in mice. *J Trace Elem Med Biol* 28: 94-99.
90. Virtanen JK, Laukkanen JA, Mursu J, Voutilainen S, Tuomainen TP (2012) Serum long-chain n-3 polyunsaturated fatty acids, mercury, and risk of sudden cardiac death in men: a prospective population-based study. *PLoS One* 7: e41046.
91. Guallar E, Sanz-Gallardo MI, van't Veer P, Bode P, Aro A, et al. (2002) Mercury, fish oils, and the risk of myocardial infarction. *N Engl J Med* 347: 1747-1754.
92. Guallar E, Sanz-Gallardo MI, van't Veer P, Bode P, Aro A, et al. (2002) Mercury, fish oils, and the risk of myocardial infarction. *N Engl J Med* 347: 1747-1754.
93. Chan HM, Egeland GM (2004) Fish consumption, mercury exposure, and heart diseases. *Nutr Rev* 62: 68-72.
94. Landmark K, Aursnes I (2004) [Mercury, fish, fish oil and the risk of cardiovascular disease]. *Tidsskr Nor Laegeforen* 124: 198-200.
95. Buscemi S, Vasto S, Di Gaudio F, Grosso G, Bergante S, et al. (2014) Endothelial function and serum concentration of toxic metals in frequent consumers of fish. *PLoS One* 9: e112478.
96. Nirwane A, Pawar V, Majumdar A (2015) Therapeutic interventions using a combination of Telmisartan and omega 3-fatty acids in sodium arsenite-induced vascular endothelial dysfunction in rats: modulation through ATP-sensitive K+ channels and eNOS. *J Complement Integr Med* 12: 143-151.
97. Vanhoutte PM (1991) Hypercholesterolaemia, atherosclerosis and release of endothelium-derived relaxing factor by aggregating platelets. *Eur Heart J* 12 Suppl E: 25-32.
98. Sena CM, Nunes E, Louro T, Proença T, Fernandes R, et al. (2008) Effects of alpha-lipoic acid on endothelial function in aged diabetic and high-fat fed rats. *Br J Pharmacol* 153: 894-906.
99. Landmesser U, Bahlmann F, Mueller M, Spiekermann S, Kirchhoff N, et al. (2005) Simvastatin versus ezetimibe: pleiotropic and lipid-lowering effects on endothelial function in humans. *Circulation* 111: 2356-2363.
100. August M, Wingerter O, Oelze M, Wenzel P, Kleschyov AL, et al. (2006) Mechanisms underlying dysfunction of carotid arteries in genetically hyperlipidemic rabbits. *Nitric Oxide* 15: 241-251.
101. Moreira EL, de Oliveira J, Dutra MF, Santos DB, Gonçalves CA, et al. (2012) Does methylmercury-induced hypercholesterolemia play a causal role in its neurotoxicity and cardiovascular disease? *Toxicol Sci* 130: 373-382.
102. Kim DS, Lee EH, Yu SD, Cha JH, Ahn SC, et al. (2005) [Heavy metal as risk factor of cardiovascular disease—an analysis of blood lead and urinary mercury]. *J Prev Med Public Health* 38: 401-407.

Citation: Omanwar S, Saidullah B, Fahim M (2015) Molecular Basis for Mercury-Induced Alteration in Endothelial Function: NO and its Modulators. *Cardiovasc Pharm Open Access* 4: 167. doi:[10.4172/2329-6607.1000167](https://doi.org/10.4172/2329-6607.1000167)