

# NAP-Related Protein 1 (Atnrp1) Overexpression Increases the Heat Tolerance of Arabidopsis Cells/Plantlets

Bíró J, Domoki M and Fehér A\*

Institute of Plant Biology, Biological Research Center, Hungarian Academy of Sciences, Temesvári krt 62, H-6726 Szeged, Hungary

## Abstract

Nucleosome assembly protein-related proteins, NRPs, were overexpressed in transgenic Arabidopsis plants. This resulted in increased survival rate of seedlings and cells exposed to 45°C for one hour. Heat shock resulted in the accumulation of the proteins, detected in the cytosol of untreated plants, in the nuclear protein fraction. DNA repair and genotoxic stress related-gene expression were unaltered in the transgenic plants as compared to controls.

**Keywords:** Nucleosome assembly protein-related protein; Elevated temperature; Survival; Transgenic; Cellular localization

## Introduction

Nucleosome-assembly protein-related proteins (NRPs) are multifunctional proteins having histone chaperone [1] and histone phosphatase inhibitor [2] properties and being implicated in root meristem maintenance [1], mitotic regulation [1], chromatin-mediated gene silencing [1], genotoxic sensitivity [1] and homologous recombination [3]. The drosophila homologue of plant NRPs, the SET protein, has been shown to accumulate at heat-shock gene loci in response to increased temperature where it co-localized with Ser10-phosphorylated histoneH3 [4]. It was therefore hypothesized that the drosophila SET protein prevents phosphatase2A (PP2A)-mediated histoneH3 dephosphorylation and contributes to chromatin modifications facilitating heat shock protein gene transcription [4].

Based on the structural and functional homology of SET and NRP proteins [2], we investigated whether the altered expression of the plant At NRP1 protein interferes with the heat sensitivity of Arabidopsis cells/seedlings.

## Experimental

### Plant cultures and heat treatment

Experiments were carried out with the wild-type Columbia ecotype of Arabidopsis thaliana (L.) Heynh and its At NRP1 overexpressor genetic transformant (NRP OX; kindly provided by Valerie Frankard, CropDesign N.V., Ghent, Belgium). In these plants, the Arabidopsis *nrp1-1* cDNA clone was cloned after the *Helianthus annuus* Gubb1 promoter for constitutive expression. T4 generation homozygous transgenic (OX) and wild type (WT) segregants of the same transformation event were used in the experiments.

Seeds of Arabidopsis thaliana were surface-sterilized by soaking in 70% ethanol (v/v) for 1 minute and in 30% bleach (v/v) for 10 minutes and then rinsed five times with sterile distilled water, and plated on solid Murashige and Skoog (MS) medium (Duchefa Biochemie, B.V., Haarlem, The Netherlands) containing 3% (w/v) sucrose, 0.6% (w/v) agar, with pH adjusted to 5.8. The seeds were germinated at 24°C using an 8 hours light/16 hours dark light regime. Experiments were carried out at least in triplicates with 7-8-days-old seedlings grown in glass petri dishes on MS medium [2]. Heat shock treatment (45°C for 45-60 minutes as indicated) was applied in a water bath (50 seedlings/dish; 4 dishes/line). Survival rates of plants (%) were visually determined one week after the treatment as the ratio of seedlings staying green and

growing six weeks after the treatment. In the case of leaf protoplast-derived cells (for isolation and culture, [5]) dead cells were stained by Evans blue (0, 5% w/v). Altogether app. 1000 cells were investigated per line in three repetitions.

## Western blotting


The antibodies used and the immunoblotting protocol is described elsewhere in details [2].

## Gene expression analysis by real-time quantitative PCR (RT-QPCR)

RNA isolation, cDNA synthesis and RT-QPCR analysis by an ABI Prism 7700 sequence detection system have been described earlier [2]. DNA damage and repair-associated (PARP2, At4g02390; RAD51, At5g20850; AtMYB, At5g03780; RNR2, At3g27060) as well as the control unresponsive (UBC18, At5g42990) genes were selected based on the work of Chen et al. [6] (Table 1).

## Results and Discussion

Using a specific antibody recognizing Arabidopsis NRP proteins



**Figure 1:** NRP protein level in wild type (WT) and overexpressing (OX) Arabidopsis plants as detected by a specific antibody ( $\alpha$ -NRP1). Proteins were stained by Ponceau reagent as loading control.

\*Corresponding author: Fehér A, Institute of Plant Biology, Biological Research Center, Hungarian Academy of Sciences, Temesvári krt. 62, H-6726 Szeged, Hungary, Tel: +36 62599701; E-mail: feher.attila@brc.mta.hu

Received April 26, 2013; Accepted July 10, 2013; Published July 17, 2013

**Citation:** Bíró J, Domoki M, Fehér A (2013) NAP-Related Protein 1 (Atnrp1) Overexpression Increases the Heat Tolerance of Arabidopsis Cells/Plantlets. J Plant Biochem Physiol 1: 115. doi:10.4172/2329-9029.1000115


**Copyright:** © 2013 Bíró J, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

[2] it was verified that NRP proteins accumulated in the AtNRP1 overproducing transgenic plants (Figure 1). 8-day-old seedlings of the wild type (WT) and NRP overexpressing (OX) plants were subjected to heat shock at 45°C. In a series of experiments the NRP overexpressing seedlings exhibited increased survival rate one week after the treatment in comparison to controls (Figure 2A and 2B).


In order to test whether this difference in heat sensitivity of seedlings is also exhibited at the cellular level, leaf protoplasts have been isolated from WT and OX plants and were similarly subjected to a one hour 45°C heat shock. Cell viability was determined using Evans blue staining dead cells. Similarly, as was observed in the case of seedlings, an app. 20% difference in cell viability could be detected in favor of NRP overexpressing cells (Figure 2C).

Basal thermo tolerance of plants depends on several factors, including hormonal regulation (ethylene, salicylic acid), production and scavenge of reactive oxygen species and the expression of various genes including heat shock transcription factors and heat shock proteins [7,8], several of which play also role during acquired thermotolerance. However, we have previously shown that NRPs are dispensable for the heat-induced expression of heat shock protein genes in Arabidopsis; despite the fact that they act as potent histoneH3 phosphatase inhibitors *in vitro* [2].


Strong heat causes DNA strand breaks what if not repaired can finally result in cell death. NRPs has recently been implicated in the maintenance of genome stability [3]. Therefore we investigated whether NRP overexpression may contribute to increased DNA repair and in this way can improve cellular survival. However, we could not detect any difference in the expression of selected DNA repair-associated (PARP2, RAD51) or genotoxic stress-upregulated (AtMYB and RNR2) genes (Figure 3). Therefore at present it is not known via what mechanism NRPs contribute to cellular heat tolerance in Arabidopsis. However, based on the presently known biochemical functions of plant NRPs,



**Figure 3.** Results of a gene expression analysis by real-time quantitative PCR. 8-days-old seedlings of non-transgenic (WT) and NRP1-overexpressing transgenic (OX) Arabidopsis lines were heat shocked (45°C for 1 hour) or were grown in the presence of 10 µg/ml bleomycin. Relative expression (treated/control) is shown for the DNA repair-associated (PARP2, RAD51) and genotoxic stress-upregulated (AtMYB and RNR2) genes [6]. Values were normalized to the genotoxic stress un-responsive UBC18 gene expression [6].



**Figure 4.** The NRP proteins are detected in the cytoplasmic fraction in seedlings grown at normal culture conditions (24°C) but in the nuclear fraction of heat shocked (45°C for 1 hour) cells as detected by a specific antibody (α-NRP1). Proteins were stained by Ponceau reagent as a loading control.



**Figure 2.** Increased survival of NRP overexpressing Arabidopsis plants (A,B) and cells (C) subjected to heat shock at 45°C.

(A) Survival rate of one-week-old seedlings treated at 45°C for 45 minutes (Exp I and II) or for 60 minutes (Exp III and IV) as determined 7-days after the treatment.

(B) Example of seedling growth one week after or without the treatment. Arrows indicate dead plants.

(C) Survival rate of leaf protoplast derived cells treated at 45°C for 60 minutes and scored for survival by the help of Evans blue staining at the indicated time points. WT-wild type; OX-overexpressor.

it might be related to their capability to influence chromatin structure and/or phosphatase activity [1,2].

Several functions, such as gene silencing, mitosis regulation and homologous recombination [1,3] of NRPs are associated with nuclear functions, and especially with the organization of the chromatin. Interestingly, when NRP proteins were detected by Western blotting in control cells and in cells heat shocked for one hour at 45°C, it was observed that while in untreated cells the proteins were in the cytosolic, in heat shocked cells they were in the nuclear protein fractions (Figure 4). Therefore one can suppose that these proteins shuttle between the cytosol and the nucleus in response to heat shock. However, the detection of NRP proteins in the cytosol of WT plants is contradictory to a previous report describing the nuclear localization of green fluorescent protein-NRP fusions [1].

The animal homologues of NRPs are also nuclear proteins having, however, cytoplasmic functions as well [9]. Therefore, further studies are required to determine the dynamic cellular location of plant NRPs and its role in the high temperature response of plant cells.

#### Acknowledgements

The presented work was supported by the NKTH-OTKA grant K68051.

Primer name	UPL	Sequence (5'-3')
PARP2 FW	68	GCGAACTATTGCTATGCCAAC
PARP2 REV		ATGTCTCCCAAAGCAACCTC
RAD51 FW	134	GATCACGGGAGCTCGATAAA
RAD51 REV		GCGGAACTCACCATATAACTCTG
UBC18 FW	67	ACAGCAATGGACATATTTGTTAGA
UBC18 REV		TGATGCAGACTGAACTCACTGTC
RNR2 FW	143	GAATCATCGCTTTCGCTTG
RNR2 REV		TTCAGCCAGAAGATTGAACAAA
AtMyb FW	134	TGTGGCTAAGTCAAAGACAGTGA
AtMyb REV		TCCACTCCCACCTTTAGCAT

(UPL = Roche "Universal probe library" number of TaqMan probes used together with the primers to ensure specificity.)

**Table 1:** Sequence of primers used in the study.

## References

- Zhu Y, Dong A, Meyer D, Pichon O, Renou JP (2006) *Arabidopsis NRP1* and *NRP2* Encode Histone Chaperones and Are Required for Maintaining Postembryonic Root Growth. *Plant Cell* 18: 2879-2892.
- Bíró J, Farkas I, Domoki M, Otvös K, Bottka S (2012) The histone phosphatase inhibitory property of plant nucleosome assembly protein-related proteins (NRPs). *Plant Physiol Biochem* 52: 162-168.
- Gao J, Zhu Y, Zhou W, Molinier J, Dong A, et al. (2012) NAP1 family histone chaperones are required for somatic homologous recombination in Arabidopsis. *Plant Cell* 24: 1437-1447.
- Nowak SJ, Pai CY, Corces VG (2003) Protein phosphatase 2A activity affects histone H3 phosphorylation and transcription in *Drosophila melanogaster*. *Mol Cell Biol* 23: 6129-6138.
- Abel S, Theologis A (1994) Transient transformation of Arabidopsis leaf protoplasts: a versatile experimental system to study gene expression. *Plant J* 5: 421-427.
- Chen IP, Haehnel U, Altschmied L, Schubert I, Puchta H (2003) The transcriptional response of Arabidopsis to genotoxic stress - a high-density colony array study (HDCA). *Plant J* 35: 771-786.
- Larkindale J, Knight MR Protection against heat stress-induced oxidative damage in Arabidopsis involves calcium, abscisic acid, ethylene, and salicylic acid. *Plant Physiol* 128: 682-695.
- Larkindale J, Hall JD, Knight MR, Vierling E (2005) Heat stress phenotypes of Arabidopsis mutants implicate multiple signaling pathways in the acquisition of thermotolerance. *Plant Physiol* 138: 882-897.
- Park YJ, Luger K (2006) Structure and function of nucleosome assembly proteins. *Biochem Cell Biol* 84: 549-558.

Citation: Bíró J, Domoki M, Fehér A (2013) NAP-Related Protein 1 (Atnrp1) Overexpression Increases the Heat Tolerance of Arabidopsis Cells/Plantlets. J Plant Biochem Physiol 1: 115. doi:10.4172/2329-9029.1000115

## Submit your next manuscript and get advantages of OMICS Group submissions

### Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

### Special features:

- 250 Open Access Journals
- 20,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>