

## Novel Diagnostic Tool CISH to Demonstrate CMV DNA by Light Microscopy-A Case Report of CMV Colitis

Barodawala SM<sup>1</sup>, Majethia NK<sup>2\*</sup> and Chadha K<sup>3</sup>

<sup>1</sup>Department of Histopathology, Grant Medical College and Sir JJ Group of Hospitals, Maharashtra, India

<sup>2</sup>Department of Pathology, Grant Medical College and Sir JJ Group of Hospitals, Maharashtra, India

<sup>3</sup>Department of Oncopathology, Grant Medical College and Sir JJ Group of Hospitals, Maharashtra, India

\*Corresponding author: Majethia NK, Grant Medical College and Sir JJ Group of Hospitals, Maharashtra, India, Tel: 02223735555; Email id: [nikhilmajethia@gmail.com](mailto:nikhilmajethia@gmail.com)

Received date: August 03, 2016; Accepted date: September 19, 2016; Publication date: September 27, 2016

Copyright: © 2016 Majethia NK, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License; which permits unrestricted use; distribution; and reproduction in any medium; provided the original author and source are credited.

### Abstract

Cytomegalovirus (CMV) infections are generally thought to be opportunistic especially in the setting of an immunosuppressed host. We present an unusual case of CMV colitis with liver abscess observed in an immunocompetent patient. The right diagnosis was made after implementation of Chromogenic *In Situ* Hybridization (CISH) on colonic biopsy tissue. Our purpose is therefore to familiarize fellow colleagues and clinicians with possible role of CISH in the diagnosis of this entity.


**Keywords:** Chromogenic *in situ* hybridization; Cytomegalovirus; Colitis

### Introduction


The prevalence of CMV infection is high, ranging from 30%-100%, depending on age and race of population under evaluation. Most cases of cytomegalovirus (CMV) colitis occur in immunocompromised patients, including those with congenital or acquired immunodeficiency diseases, those receiving immunosuppressive drugs and transplant recipients. Only a small number of immunocompetent patients with CMV colitis have been reported worldwide [1-3]. CMV has a propensity to involve rapidly growing tissue such as granulation tissue in an ulcer bed [4]. We present an unusual case of CMV colitis with liver abscess observed in an immunocompetent patient. The right diagnosis was made after implementation of Chromogenic *In Situ* Hybridization (CISH) on colonic biopsy tissue. Our purpose is therefore to familiarize fellow colleagues and clinicians with possible role of CISH in the diagnosis of this entity.

### Case Report

A 69 year old male presented with diarrhea, gastrointestinal bleed, abdominal pain and fever. Ultrasonography of the abdomen examination revealed a solitary liver abscess and caecal thickening. Endoscopic evaluation showed swollen and thickened walls of colon with multiple erythematous patches and ulcerative changes in caecum and right colon. The patient also had antral gastritis. The clinical diagnosis was amoebic colitis. A caecal biopsy was submitted for histopathologic examination. It revealed moderate colitis with focal ulceration and submucosal granulation tissue. The latter showed proliferating capillaries and fibroblasts along with dense neutrophilic and moderate lymphoplasmacytic infiltrate. Periodic acid-Schiff (PAS) stain did not reveal trophozoites of *Entamoeba histolytica* ruling out the clinical diagnosis. The noteworthy feature was the presence of enlarged endothelial cells with basophilic intra nuclear inclusions (owl eye) consistent with CMV infected cells (Figure 1 and 2).



**Figure 1:** Photomicrograph of enlarged endothelial cells with basophilic intra nuclear inclusions consistent with CMV infected cells.



**Figure 2:** Photomicrograph of CMV inclusion on CISH.

The tissue was subjected to Chromogenic *in situ* hybridization (CISH) for CMV using Zytostat Digoxigenin-labeled oligonucleotide probe for the detection of Cytomegalovirus (CMV) DNA. The CMV DNA in the cells or tissue links to the secondarily polymerized enzyme-conjugated antibody to this DNA. The chromogenic substrate is then linked to this antibody which leads to the formation of a color precipitate that is visualized by light microscopy. Following the report of CMV infection, the patient received antiviral therapy and he is doing well.

## Discussion

Case reports for CMV infection in colon have been well documented in immunocompetent patients, the first case reported in 1992 [5,6], sigmoid colon being the most common site [1]. In immunocompetent patients, diabetes mellitus, renal failure, severe trauma, sepsis, shock, burns, cirrhosis, are predisposing like in our case. Our patient presented with multiple ulcers in caecum along with a liver abscess. Imitating other more common conditions, like amoebic colitis. Other differential diagnosis include ischemic colitis, pseudomembranous colitis and Inflammatory bowel disease [7]. However, these were ruled out on histopathological examination and histochemical stains i.e. PAS stain. Due to presence of large cells with basophilic intranuclear inclusions surrounded by a clear halo (owl eye) in endothelial cells a diagnosis of cytomegalovirus infection was suspected and confirmed by CISH for CMV. Until recently, the diagnosis of CMV infection had been limited to serological conversion, culture and detection of the virus by cytopathic effects on body fluids or tissue or the identification of typical CMV inclusion bodies in biopsy specimens. The application of mRNA or DNA probes specific for CMV has been shown to be sensitive and specific in adjunct to histological criteria. Similarly, detection of CMV antigens like pp65 with IgM monoclonal antibodies using immunohistochemistry can be carried out [8], CISH is a type of Bright field *in situ* hybridization (BRISH) which can be carried out on the slide prepared from the paraffin block. The BRISH includes CISH and SISH where in a chromogen (e.g. DAB) and silver are used to highlight the DNA /RNA of infectious agents like HPV DNA, EBER RNA of EBV (Epstein Barr virus in cases of lymphoma) [9] and CMV DNA. The latter is what we report in our study. It is to be noted that the specificity of demonstrating CMV DNA by CISH is 100%. The use of *in situ* hybridization is not new as it is more popularly used as Fluorescence *in situ* hybridization (FISH) for demonstrating HER-2/neu in breast carcinoma [10] cases and prognosticating hemato-lymphoid malignancies. It offers a real advantage by providing a cytomorphologic link where in slides can be analyzed by an anatomical pathologist using a light microscope [11].

## Conclusion

The purpose of documenting this case is to highlight the fact that although *in situ* hybridization technique is already being utilized in oncology practice but its role in confirming infectious etiologies on tissues is less explored. Our 3 years experience in recognizing HPV, EBV, CMV has optimized many patients management. This is one such case where CMV illustration by CISH could help treat the patient. CISH is a novel the tool for diagnosing such cases where other methods cannot be used and can be an integral part of armamentarium available to the surgical pathologist.

## References

1. Lin MW, Liang JT, Chang KJ, Lee PH, Lin BR (2008) Cytomegalovirus Colitis in an Immunocompetent Patient: Report of a Case and Review of the Literature. J Soc Colon Rectal Surgeon 19: 27-32.
2. Karakozis S, Gongora E, Caceres M, Brun E, Cook JW, et al. (2001) Life-threatening cytomegalovirus colitis in the immunocompetent patient: report of a case and review of the literature. Dis Colon Rectum 44: 1716-1720.
3. Lee CS, Low AH, Ender PT, Bodenheimer HC (2004) Cytomegalovirus colitis in an immunocompetent patient with amebiasis: case report and review of the literature. Mt Sinai J Med 71: 347-350.
4. Christidou A, Zambeli E, Mantzaris G (2007) Cytomegalovirus and Inflammatory Bowel Disease: pathogenicity, diagnosis and treatment. Ann Gastroenterol 20: 110-115.
5. Blair SD, Forbes A, Parkins RA (1992) CMV colitis in an immunocompetent adult. J R Soc Med 85: 238-239.
6. Paparoupa M, Schmidt V, Weckauf H, Ho H, Schuppert F (2016) CMV Colitis in Immunocompetent Patients: 2 Cases of a Diagnostic Challenge. Case Rep Gastrointest Med.
7. Subbalaxmi MVS, Kumar VG, Chandra N, Raju YSN (2014) Cytomegalovirus colitis in a human immunodeficiency virus seropositive individual with moderately severe immunodeficiency. The J Clin Sci Res 3: 42-45.
8. Weiss LM, Chen YY (2013) EBER *in situ* hybridization for Epstein-Barr virus. Methods in Molecular Biology 999: 223-230.
9. Garrido E, Carrera E, Manzano R, Lopez-Sanroman A (2013) Clinical significance of cytomegalovirus infection in patients with inflammatory bowel disease. World Journal of Gastroenterology 19: 17-25.
10. Gupta D, Middleton LP, Whitaker MJ, Abrams J (2003) Comparison of fluorescence and chromogenic *in situ* hybridization for detection of HER-2/neu oncogene in breast cancer. American journal of clinical pathology 119: 381-387.
11. Nitha H, Grogan TM (2013) Bright Field *in situ* hybridization methods to discover gene amplifications and rearrangements in clinical samples. Methods in Molecular Biology 986: 341-352.