



This is triangle shape.



This is dot shape.

Figure 1: Clinical findings of erythema on the base of the head.




Figure 2: Shapes of typical rashes.

The reason for this high incidence remains unknown, and the shape of the patch may be associated with its likelihood to fade or disappear. To the best of our knowledge, there are no available statistical global data on nuchal salmon patches, possibly because most people, including physicians, are not interested in investigating it. The nuchal salmon patch should be investigated in studies that include more cases.

Conclusion

Nuchal salmon patches tend to persist and our study found an incidence of persistent nuchal salmon patch that was both higher than that in previous report and higher than we expected.

References

1. Leung AK (2009) Salmon patches in *The Encyclopedia of Molecular Mechanisms of Disease*, F. Lang, Ed., Berlin, Germany. Springer: 1880-1881.
2. Leung AKC, Telmesani AMA (1989) Salmon patches in Caucasian children. *Pediatr Dermatol* 6: 185-187.
3. Rivers JK, Frederiksen PC, Dibdin C (1990) A prevalence survey of dermatoses in the Australian neonate. *J Am Acad Dermatol*. 23: 77-81.
4. Sachdeva M, Kaur S, Nagpal M, Dewan SP (2002) Cutaneous lesions in new born. *Indian J Dermatol Venereol Leprol* 68: 334-337.
5. Ferahbas A, Utas S, Akcakus M, Gunes T, Mistik S, et al. (2009) Prevalence of cutaneous findings in hospitalized neonates: a prospective observational study. *Pediatr Dermatol*. 26: 139-142.
6. Monteagudo B, Labandeira J, Acevedo A, Cabanillas M, León-Muiños E, et al. (2011) Salmon patch: a descriptive study. *Actas Dermosifiliogr* 102: 24-27.
7. Corson EF (1934) Nevus flammeus nuchae: its occurrence and abnormalities. *The Am J Medical Sci* 187:121-124.
8. Verbov J, Steinberg R (1974) The persistent nuchal or occipital salmon patch. *Br J Dermatol* 90: 586-587.
9. Oster J, Nielsen A (1970) Nuchal naevi and interscapular telangiectases. *Acta Paediatrica Scand* 59: 416-423.