

Nutrient Profile of Native Woody Species and Medicinal Plants in Northeastern Mexico: A Synthesis

Ratikanta Maiti¹, Humberto Gonzalez Rodriguez^{1*} and Aruna Kumari²

¹Facultad de Ciencias Forestales, Universidad Autónoma de Nuevo León, Carr. Nac. No. 85 Km. 45, Linares, Nuevo León 67700, México

²Plant Physiology, Agricultural College, Professor Jaya Shankar Telangana State Agricultural University, Polasa, Jagtial, Karimnagar, Telangana, India

Abstract

Leaf nutrients have an important role in governing the trees growth and development of trees and are sources of nutrients for ruminants in forest. The present paper makes a synthesis of two earlier studies undertaken on macro and micronutrients of 1) 37 woody species and 2) 44 medicinal plants used traditionally of Northeastern Mexico. In the both the studies we estimated the contents of six nutrients in the leaves, three macronutrients (P, Mg, K, protein, C, N, C/N and three micronutrients (Cu, Fe, ZN) both for woody species and medicinal plants. A large variability exists in nutrient contents in woody species and medicinal plants. The species selected for the highest macronutrients and micronutrients may be utilized for confirming their physiological efficiency and probable better growth and productivity as well as good sources of macro and micronutrients for grazing wild animals. Similar to the results of woody trees, medicinal plants also exhibited a wider variability in macro and micronutrients. Based on the analysis of macro and micro nutrients the following species *Phalaris canariensis*, *Eruca sativa*, *Ocimum basilicum*, *Tragia ramosa*, *Mentha piperita*, *Mimosa malacophylla*, *Acalypha monostachya*, *Salvia hispánica*, *Tillandsia usenoides*, *Letsea glauscens*, *Tagetes lucida*, *Erygium heterophylla*, *Dyssodia setifolia*, *Nicotiana glauca*, *Ruta graveolens*, *Olea europea*, *Equisetum hyemale*, *Rosmarinus officinalis* are selected containing a majority of each of these nutrients.

Keywords: Macro-Micronutrients; Native woody plants; Medicinal plants; Mexico; Variability; High nutritional values

Introduction

Woody plants and medicinal plants possess various macro nutrients and micronutrients. These are essential to bring about the growth and development of the plants. These because of their medicinal values are used traditionally to cure various diseases. We made a short synthesis of researches undertaken on macro and micronutrients of woody species and medicinal plants.

Nutrients of woody plants

Woody plants of Tamaulipan Thornscrubs in the semiarid regions are of great economic importance for various uses such as timber for furniture, fences, firewood and serve as sources of forage for wild animal grazing [1]. Few studies are undertaken on nutrient contents of woody plants. The specific leaf area (SLA, leaf area per unit mass) appears to be correlated with leaf nitrogen per unit dry mass, photosynthesis and dark respiration sites [2].

A large variation exists among species in nutrient conservation. Species having nutrient conservation have long life span, high leaf mass per area, low nutrient concentrations and low photosynthetic capacity [3]. In a review a synthesis of crop responses to nutrient stress was made with special emphasis on nutritional values of nitrogen and phosphorus because these elements limit plant growth [4]. Leaf nutrient contents depend on the availability of macro and micronutrients that are present in the soil habitat. Nutrient-poor habitats are generally dominated by species that are nutrient conserving. The fertile habitats are generally dominated by species which have short term productivity per unit leaf mass [5]. Within a given habitat, the coexistence of the species is seen that exhibit variability in a large range of leaf traits [3].

With the age of leaves, nutrients are resorbed from older leaves prior to leaf abscission and reemployed in the developing tissues of leaves, fruits, and seeds. Resorption in general occurs throughout a leaf's life, more particularly when the leaves are shaded [6,7]. A major

phase of resorption occurs shortly before leaf abscission [8] nutrient is recycled via resorption process [9]. It is reported that the presence of nutrient sinks that are active in particular have control over resorption [10]. Wright et al. [2] reported that most plants withdraw nutrients from leaves with advancement in age.

The nutrient availability and management were studied by [11] in the rhizosphere showing genotypic differences. Understanding the role of plant-microbe-soil interaction governing the nutrient availability will enhance environmental sustainability. Green- leaf and senesced-leaf nitrogen and phosphorous contents quantified revealed that these in green and senesced leaves were positively correlated with leaf length in all species.

A study was conducted on chemical composition and potential value of subtropical tree species of Combretum in southern Africa for ruminants. It was concluded that the foliage tested would not be a suitable resource of N to supplement protein deficiencies in low quality herbage [12].

Nutrients in medicinal plants

The role of traditional medicinal plants to alleviate various diseases in different countries since remote times is well known. In Mexico, many medicinal plants are commonly used traditionally to cure various

***Corresponding author:** Humberto Gonzalez Rodriguez, Facultad de Ciencias Forestales, Universidad Autónoma de Nuevo León, Carr. Nac. No. 85 Km. 45, Linares, Nuevo León 67700, México, Tel: 8212124895 extn. 136 y 236; E-mail: humberto.gonzalez@uanl.mx

Received May 04, 2016; Accepted May 26, 2016; Published May 31, 2016

Citation: Maiti R, Rodriguez HG, Kumari A (2016) Nutrient Profile of Native Woody Species and Medicinal Plants in Northeastern Mexico: A Synthesis. J Bioprocess Biotech 6: 283. doi:10.4172/2155-9821.1000283

Copyright: © 2016 Maiti R, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

diseases in rural and urban areas [13-15]. Although various studies on ethanobotany of medicinal plants have been undertaken [16], only few cases were documented and reported in the northeastern area of Mexico [17-22]. Besides, very little information about these species is available to examine the efficacy of each species based on of chemical analysis.

Several studies in medicinal plants have been undertaken on the use of the use of the medicinal plants to cure human and animal diseases such as in eye problem [23] and diabetes [24,25]. The micronutrients are essential for promoting good health of mankind, plants and animals. Their requirement is though in minor quantities the deficiency of these elements causes abnormalities that lead to infection of diseases [23].

Several medicinal plants possessing micronutrients are reported to be effective to control blood sugar such as Mulberry leaf, soybean, black bean, Persian shallot, *Zizyphus lotus*, etc. It is reported that Carbon, Nitrogen and C/N ratios contribute to the production of a number of secondary metabolites, antioxidants and flavonoids which are useful for the maintenance of health of human beings [26,27]. Antioxidants have the ability to reduce inflammations and promote normal cardiovascular functions [28] inhibits cancerous tumour [29] reduce the ageing process in the brain and nervous system [30] and delay or inhibit oxidation [31]. Several antioxidant compounds are present in some fruits, vegetables viz., *Occimum sanctum*, *Terminalia belerica*, *Zingiber officinales* and several Chinese and Indian spices such as *Poper cababa*, *Allium sativa*. The flavones, isoflavones, flavonoids, anthocyanins, coumarins ligans, catechin and isocatechin and several secondary metabolites that are commonly found in plants exhibit the antioxidant activities [32,33]. High peroxidation of the secondary metabolites was reported in plants that have high C/N ratio and low nitrogen fertilization especially when exposed to elevated CO₂ levels. Under low nitrogen conditions, the growth and photosynthesis in plant increases the C/N ratio which leads to an increase in the production of secondary metabolites [30,34].

Very recently the consumption of nutraceuticals from plant has become popular to improve health and to prevent or treat various diseases. Some popular phyto-nutraceuticals are glucosamine from ginseng, Omega 3 fatty acids from linseed. Many of the nutraceuticals have high potentials for multiple therapeutic uses but research is lacking [35].

Methodology

These two studies were carried out at the experimental station of Facultad de Ciencias Forestales, Universidad Autonoma de Nuevo Leon, located in the municipality of Linares (24 47N 99 32 W), at elevation of 350 m. The climate is subtropical or semiarid with warm summer, monthly mean air temperature vary from 14.7°C in January to 23°C in August, although during summer the temperature goes up to 45°C. Average annual precipitation is around 805 mm with a bimodal distribution. The dominant type of vegetation is the Tamaulipan Thorn scrub or subtropical Thorn scrub wood land. The dominant soil is deep, dark grey, lime-grey, vertisol with montmorrillonite, which shrink and swell remarkably in response to change in moisture content.

Chemical analysis

The leaf samples were collected from each species and were placed to dry on a newspaper for a week. The leaves were separated from the rest of the plant and were passed twice through a mesh of 1 × 1 mm in diameter using a mill Thomas Wiley and subsequently dried for more than three days at 65°C in an oven (Precision model 16EG) to remove

moisture from the sample and later these were placed in desiccators. A 2.0 mg of the sample was weighed in an AD6000 Perkin balance elmer in a vial of tin, bent perfectly. This was placed in Chons analyzer Perkin Elmer Model 2400 for determining carbon, and nitrogen. For estimating the mineral contents, the samples were incinerated in a muffle oven at 550°C for 5 hours. The ashed sample is then digested in a solution containing HCL and HNO₃, using the technique of wet digestion [36]. The estimations of Carbon and nitrogen foliar contents (% dry mass basis) were carried out in 0.020 g of milled dried leaf tissue by using a CHN analyser (PerkinElmer, model 2400). Protein is calculated following $N \times 6.25$.

Mature leaf samples (1.0 g dry weight) obtained from each plant and shrub species was used for determining the contents of minerals (Cu, Fe, Zn, Mg, K and P). Mineral content was estimated by incinerating samples in a muffle oven at 550°C during 5 hours. Ashes were digested in a solution containing HCl and HNO₃, using the wet digestion technique [36]. Cu, Fe, Zn, K and Mg (air/acetylene flame) were determined by atomic absorption spectrophotometry (Varian, model SpectrAA-200), whereas P was quantified spectrophotometrically using a PerkinElmer spectrophotometer (Model Lamda 1A) at 880 nm [37].

Results

In the present paper we presented the brief results of two studies, 1) leaf macro and micronutrients of 37 woody species and 2) of 44 medicinal plant species that were used traditionally in Northeastern Mexico during summer 2015.

Study 1: Macro and micronutrients of 37 woody species

Results: The data regarding the macro- and microminera; contents of 37 native shrubs and trees are presented in Table 1. It is observed that there exists a large variability in the contents of macro (P, Mg, K, protein, C, N ($\mu\text{g g}^{-1}$ dw), C/N ratio) and micronutrients (Cu, Fe, Zn ($\mu\text{g g}^{-1}$ dw), thereby giving large opportunity for the selection of the species with high nutrient contents. In the following we selected species with high macro and micronutrients which could serve as good source of nutrients.

Macronutrients (mg g^{-1} dw):

P: The species containing high P are *Croton suaveleons* (2.43), *Eysenhardtia polystachya* (1.84), *Prosopis laevigata* (1.65), *Parkinsonia aculeate* (1.56), *Acacia farnesiana* (1.54), *Salix lasiolepis* (1.15).

Mg (mg g^{-1} dw): The species containing high Mg are *Ehretia anacua* (9.45), *Condalia hookeri* (6.50), *Parkinsonia aculeata* (5.29), *Helietta parviflora* (3.17), *Guaiacum angustifolia* (4.1).

K (mg g^{-1} dw): The species containing high K are *Croton suaveolens* (75.62), *Cordia boissieri* (45.58), *Celtis pallida* (42.6), *Acacia rigidula* (38.75), *Diospyros texana* (36.55), *Acacia farnesiana* (34.72), *Prosopis laevigata* (34.04), *Bumelia celastrina* (33.02).

C%: The species containing high C are *Leucophyllum frutescens* (49.97), *Foresteria angustifolia* (49.47), *Bumelia celastrina* (49.25), *Acacia berlandieri* 49.18, *Acacia rigidula* (48.23), *Gymnosperma glutinosum* (46.19), *Acacia farnesiana* (46.17), *Croton suaveolens* (45.17), *Sargentia gregii* (44.07).

N%: The species containing high N% are *Gymnosperma glutinosum* (5.89), *Lantana macropoda* (4.43), *Acacia shaffneri* (4.32), *Bernardia myricifolia* (2.21), *Celtis pallida* (4.12), *Eysenhardtia polystachya* (4.06), *Cercidium macrum* (4.01).

Species	Leaf macro-nutrient contents						Leaf micronutrient contents		
	(mg g ⁻¹ dw)			(%)			(µg g ⁻¹ dw)		
	K	Mg	P	C	N	C:N	Cu	Fe	Zn
<i>Acacia berlandieri</i>	0.78 ± 0.08	2.69 ± 0.41	6.80 ± 2.10	23.88	49.18 ± 1.25	3.82 ± 0.14	3.52 ± 0.62	73.46 ± 8.01	15.08 ± 3.61
<i>Acacia farnesiana</i>	1.54 ± 0.11	0.22 ± 0.17	34.72 ± 2.20	21.31	46.17 ± 2.63	3.41 ± 0.18	24.62 ± 1.11	259.76 ± 2.66	15.47 ± 0.83
<i>Acacia rigidula</i>	1.25 ± 0.14	0.43 ± 0.09	38.75 ± 0.55	16.25	48.23 ± 1.56	2.60 ± 0.22	7.09 ± 0.36	252.33 ± 3.04	10.23 ± 1.16
<i>Acacia schaffneri</i>	1.44 ± 0.22	1.72 ± 1.17	19.86 ± 1.77	27	39.52 ± 0.99	4.32 ± 0.16	3.18 ± 0.94	138.93 ± 32.25	44.6 ± 5.71
<i>Acacia wrightii</i>	1.22 ± 0.19	3.03 ± 1.23	20.5 ± 3.41	24.75	36.59 ± 1.11	3.96 ± 0.18	8.11 ± 2.97	99.04 ± 23.21	28.14 ± 2.29
<i>Amyris texana</i>	1.09 ± 0.08	2.67 ± 0.36	19.56 ± 4.79	23.25	38.06 ± 1.89	3.72 ± 0.33	9.18 ± 1.17	99.88 ± 31.75	17.40 ± 1.24
<i>Berberis chococo</i>	0.90 ± 0.05	2.35 ± 0.86	12.42 ± 2.06	15.19	36.91 ± 1.25	2.43 ± 0.19	5.12 ± 0.38	58.79 ± 13.95	50.68 ± 9.41
<i>Bernardia myricifolia</i>	1.09 ± 0.10	3.61 ± 0.38	11.54 ± 1.18	26.31	42.69 ± 1.13	4.21 ± 0.49	8.03 ± 0.85	139.73 ± 24.69	16.17 ± 0.93
<i>Bumelia celastrina</i>	0.09 ± 0.78	0.68 ± 0.13	33.02 ± 1.30	15.13	49.25 ± 1.56	2.42 ± 0.36	25.24 ± 1.50	249.00 ± 15.03	14.10 ± 7.38
<i>Caesalpinia mexicana</i>	1.29 ± 0.23	1.20 ± 0.12	13.32 ± 1.88	18.19	41.12 ± 1.96	2.91 ± 0.38	4.55 ± 0.96	48.47 ± 19.26	17.36 ± 5.74
<i>Celtis laevigata</i>	1.57 ± 0.23	2.19 ± 0.27	20.67 ± 3.28	18.81	39.45 ± 0.51	3.01 ± 0.18	8.88 ± 2.09	254.09 ± 40.44	42.28 ± 5.49
<i>Celtis pallida</i>	1.24 ± 0.18	3.20 ± 0.25	42.60 ± 0.90	25.75	38.66 ± 0.88	4.12 ± 0.67	25.98 ± 1.04	276.89 ± 5.70	12.42 ± 0.29
<i>Cercidium macrum</i>	1.10 ± 0.11	2.95 ± 0.87	14.5 ± 6.07	25.06	43.41 ± 3.44	4.01 ± 0.30	5.97 ± 1.50	96.08 ± 24.65	25.29 ± 5.86
<i>Condalia hoockeri</i>	0.89 ± 0.06	6.50 ± 0.84	25.76 ± 3.33	19.13	30.07 ± 2.81	3.06 ± 0.41	5.02 ± 0.39	73.79 ± 18.44	11.54 ± 3.41
<i>Cordia boissieri</i>	1.42 ± 0.12	2.72 ± 0.31	45.58 ± 1.65	20.5	43.43 ± 1.20	3.28 ± 0.09	30.71 ± 0.55	280.55 ± 8.46	51.87 ± 1.80
<i>Croton suaveolens</i>	2.43 ± 0.14	0.22 ± 0.09	75.62 ± 3.67	14.5625	45.17 ± 0.35	2.33 ± 0.53	26.87 ± 1.66	229.13 ± 24.25	34.55 ± 4.11
<i>Diospyros palmeri</i>	0.96 ± 0.06	2.84 ± 0.92	18.13 ± 1.60	13.56	37.59 ± 1.72	2.17 ± 0.12	5.36 ± 1.13	92.96 ± 14.59	18.58 ± 5.90
<i>Diospyros texana</i>	0.98 ± 0.08	2.59 ± 0.55	36.55 ± 2.26	11.81	40.79 ± 1.46	1.89 ± 0.06	2.80 ± 0.16	72.47 ± 22.66	41.45 ± 4.03
<i>Ebenopsis ebano</i>	0.90 ± 0.03	2.88 ± 0.25	14.06 ± 1.27	24.13	37.57 ± 1.21	3.86 ± 0.20	8.85 ± 2.14	130.60 ± 94.49	17.21 ± 3.81
<i>Ehretia anacua</i>	1.12 ± 0.06	9.45 ± 0.37	16.72 ± 1.18	15.25	34.09 ± 2.51	2.44 ± 0.10	12.62 ± 0.79	68.90 ± 5.82	40.07 ± 5.31
<i>Eysenhardtia polystachya</i>	1.84 ± 0.22	2.22 ± 0.24	16.43 ± 2.54	25.375	36.26 ± 0.58	4.06 ± 0.27	16.16 ± 2.92	82.86 ± 10.47	51.39 ± 10.66
<i>Forestiera angustifolia</i>	0.89 ± 0.06	1.27 ± 0.35	28.32 ± 3.61	18.75	49.47 ± 0.43	3.00 ± 0.41	4.03 ± 0.39	70.10 ± 11.86	48.56 ± 6.13
<i>Fraxinus greggii</i>	0.88 ± 0.05	1.55 ± 0.54	23.07 ± 1.63	13.44	38.06 ± 1.89	2.15 ± 0.14	8.16 ± 1.21	125.13 ± 43.75	30.94 ± 3.97
<i>Guaiaecum angustifolium</i>	0.80 ± 0.18	4.10 ± 0.87	12.69 ± 1.32	18.13	41.89 ± 3.56	2.90 ± 0.42	3.99 ± 2.07	83.30 ± 6.70	20.59 ± 4.08
<i>Gymnosperma glutinosum</i>	1.40 ± 0.04	1.90 ± 0.13	21.49 ± 3.31	36.8125	46.19 ± 1.04	5.89 ± 0.29	8.93 ± 2.40	167.40 ± 13.81	12.16 ± 1.64
<i>Harvardia pallens</i>	1.11 ± 0.11	3.15 ± 0.68	22.86 ± 2.52	18.5625	43.49 ± 1.24	2.97 ± 0.15	3.51 ± 0.46	109.87 ± 10.53	29.57 ± 6.38
<i>Helietta parvifolia</i>	1.01 ± 0.12	5.17 ± 0.45	21.48 ± 1.83	15.19	31.13 ± 1.03	2.43 ± 0.25	9.11 ± 2.21	72.94 ± 4.97	37.55 ± 9.07
<i>Karwinskia humboldtiana</i>	1.05 ± 0.12	1.31 ± 0.37	16.41 ± 2.28	17.75	31.35 ± 0.70	2.84 ± 0.10	5.86 ± 0.75	70.41 ± 12.17	13.19 ± 1.61
<i>Lantana macropoda</i>	1.37 ± 0.06	3.71 ± 0.31	26.04 ± 2.04	27.69	42.91 ± 3.74	4.43 ± 0.39	13.00 ± 0.22	145.81 ± 48.14	28.29 ± 4.72
<i>Leucaena leucocephala</i>	0.98 ± 0.05	2.60 ± 0.38	14.21 ± 2.03	23.63	43.16 ± 1.98	3.78 ± 0.50	6.03 ± 0.41	77.59 ± 2.20	11.06 ± 1.12
<i>Leucophyllum frutescens</i>	0.8 ± 0.10	2.69 ± 0.28	13.69 ± 3.10	14.06	49.97 ± 0.94	2.25 ± 0.27	6.45 ± 0.79	118.12 ± 15.44	27.23 ± 3.79
<i>Parkinsonia aculeata</i>	1.56 ± 0.35	5.29 ± 1.82	24.93 ± 2.81	19	36.63 ± 3.25	3.04 ± 0.41	7.44 ± 2.20	165.63 ± 69.17	51.66 ± 8.09
<i>Prosopis laevigata</i>	1.65 ± 0.29	2.88 ± 1.12	34.04 ± 2.03	24.06	41.64 ± 0.71	3.85 ± 0.21	5.17 ± 1.53	128.92 ± 41.18	48.47 ± 11.71
<i>Quercus virginiana</i>	0.91 ± 0.07	2.60 ± 1.34	15.04 ± 1.37	12.25	43.02 ± 2.38	1.96 ± 0.18	3.63 ± 0.50	66.32 ± 13.19	39.25 ± 3.88
<i>Salix lasiolepis</i>	1.51 ± 0.09	2.34 ± 1.18	23.57 ± 1.54	12.875	33.37 ± 4.58	2.06 ± 0.50	8.49 ± 0.52	95.49 ± 13.15	144.86 ± 12.77
<i>Sargentia greggii</i>	0.78 ± 0.04	2.45 ± 0.19	13.19 ± 3.84	11.94	44.07 ± 1.22	1.91 ± 0.45	4.79 ± 0.47	87.80 ± 18.23	14.48 ± 0.94
<i>Zanthoxylum fagara</i>	0.99 ± 0.13	2.80 ± 0.46	14.77 ± 2.51	18.63	40.35 ± 3.15	2.98 ± 0.90	15.66 ± 3.17	112.80 ± 22.32	18.92 ± 2.80

Data are means and standard deviation (n=5).

Table 1: Leaf nutrient content in different plant species.

C/N ratio: The species containing high C/N are *Sargentia gregii* (23.13), *Leucophyllum frutescens* (22.17), *Quercus virginiana* (21.95), *Diospyros texana* (21.58), *Bumelia celastrina* (20.35), *Croton suaveolens* (20.16).

Micronutrients: Cu ($\mu\text{g g}^{-1}$ dw): The species containing high Cu are *Cordia boissieri* (30.71), *Croton suaveolens* (26.87), *Celtis pallida* (25.98), *Bumelia celastrina* (23.24), *Acacia farnesiana* (24.62), *Eysenhardtia polystachya* (16.16), *Zanthoxylum fragara* (15.66), *Lantana macropoda* (13.0), *Ehretia anacua* (12.62).

Fe ($\mu\text{g g}^{-1}$ dw): The species containing high Fe are *Cordia boissieri* (280.55), *Celtis pallida* (276.89), *Acacia farnesiana* (259.76), *Celtis laevigata* (234.09), *Acacia rigidula* (252.33), *Bumelia celastrina* (249), *Croton suaveolens* (229.19), *Gymnosperma glutinosum* (167.4), *Parkinsonia aculeata* (165.63), *Lantana macropoda* (145.81), *Bernardia myricifolia* (139.73), *Acacia shaffneri* (138.93).

Zn ($\mu\text{g g}^{-1}$ dw): The species containing high Zn are *Salix lasiolepis* (144.86), *Cordia boissieri* (51.87), *Parkinsonia aculeata* (51.66), *Eysenhardtia polystachya* (51.39), *Foresteria angustifolia* (48.56), *Prosopis laevigata* (48.47), *Acacia shaffneri* (44.60), *Celtis laevigata* (42.28), *Diospyros texana* (41.45), *Ehretia anacua* (40.07).

In the context of the above results it may be stated that the species show large variation in the contents of six macro (P, Mg, K, C, N, C/N, % protein) and three micronutrients Cu, Fe and Zn, thereby offering an opportunity to select species for high macro and micronutrients. The roles of nutrients are emphasized in various studies [4-6] and others.

Discussion: Studies on variability in leaf nutrient contents among woody species are rare but various studies have been demonstrated on nutrient contents of plants with respect to nutrient conservation, photosynthetic capacity (3), availability of leaf nutrients in soil habitat (5), resorption of nutrients from older leaves with age of leaves (6,7); leaf abscission (8); nutrient management in rhizosphere (11); protein deficiencies in low quality herbage (2). In the context of the above results and discussion, there is a research need to study the nutritional values of the selected species for animal nutrition and the productivity of trees and establish a germplasm bank with high nutrient profile for future research.

Study 2: Nutrient profile (Macro and micronutrients) of 44 medicinal plants used traditionally in Northeastern Mexico

Results: We determined macronutrients (K, Mg, P, C, C/N), protein content, and micronutrients (Cu, Fe, Zn) from the 44 medicinal plants (Table 2). The information helps to select plants with high quantity of each nutrient. This will in turn confirm the efficacy of the medicinal activity of the plant.

It is observed from the Table 3 that there exists a large variability in the contents of macro nutrients (K, Mg, P, N, C/N (mg g^{-1} dwt) and micronutrients Fe, Cu, Zn ($\mu\text{g g}^{-1}$ dwt). Therefore, a large variation in the contents of nutrients and protein%, offers an opportunity to scientists working on medicinal plants to select plants with nutritional values. In the following the medicinal plants were grouped based on the maximum content of macro and micro nutrients.

Macronutrients:

• Species containing high Potassium (K) content (mg g^{-1} dwt)

Phalaris canariensis (163.3), *Eruca sativa* (144.2), *Opuntia ficus-indica* (101.5), *Phoradendron villosum* (100.6), *Melia azadirachta* (100), *Moringa oleifer* (96.0), *Marrubium vulgare* (91.3).

• Species containing high Magnesium (Mg) content (mg g^{-1} dwt)

Mimosa malacophylla (8.6), *Acalypha monostachya* (8.1), *Opuntia ficus-indica* (6.4), *Salvia hispanica* (3.5).

• Species containing high Phosphorus (P) content (mg g^{-1} dwt)

Celtis laevigata (4.03), *Carya illinoensis* (2.89), *Croton suaveolens* (2.40), *Litsea glaucescens* (2.40), *Eryngium heterophyllum* (2.27), *Lepidium virginicum* (2.27), *Cnidioscolus aconitifolius* (2.05), *Mentha piperita* (2.0).

• Species containing with high Carbon (C) (Per cent) capacity of Carbon dioxide fixation

Eugenia caryophyllata (51.66), *Rhus virens* (50.38), *Cinnamomum verum* (49.34), *Arbutus xalapensis* (49.1), *Tecoma stans* (48.79), *Eryobotria japonica* (47.98), *Rosamrinus officinalis* (47.77), *Hedeoma palmeri* (46.38), *Moringa oleifer* (45.96), *Buddleja cordata* (45.70), *Croton suaveolens* (45.17), *Mimosa malacophylla* (45.15), *Chrysactinia mexicana* (45.04), *Melia azadirachta* (45.12).

• Species containing high Nitrogen (N) (per cent)

Mimosa malacophylla (8.46), *Moringa oleifer* (6.25), *Tagetes lucida* (5.98), *Tagetes lucida* (5.89), *Marrubium vulgare* (5.58), *Eruca sativa* (5.48), *Salvia hispanica* (5.24).

• Species containing high C/N ratio

Agave macroculmis Todaro (31.04), *Arbutus xalapensis* (26.94), *Eryngium heterophyllum* (24.23), *Rhus virens* (22.92), *Olea europea* (22.08), *Croton suaveolens* (20.16).

• Protein%

Mimosa malacophylla (52.87), *Moringa oleifer* (39.04), *Tagetes lucida* (36.81), *Poliomintha longiflora* (36.54), *Eruca sativa* (34.23), *Lepidium virginicum* (33.73), *Melia azadirachta* (36.55), *Nicotiana glauca* (29.1), *Phoradendron villosum* (30.76).

Micronutrients:

• Species with high Copper (Cu) content ($\mu\text{g g}^{-1}$ dwt)

Celtis laevigata (33.88), *Phalaris canariensis* (33.78), *Tillandsia usenoides* L. (33.70), *Arbutus xalapensis* (33.40), *Lepidium virginicum* (33.06), *Phalaris canariensis* (33.04), *Lepidium virginicum* (31.06), *Buddleja cordata* (29.90), *Cnidioscolus aconitifolius* (29.67).

• Species with high Iron (Fe) content ($\mu\text{g g}^{-1}$ dwt)

Gnaphalium canescens (3973.55), *Tragia ramosa* (1450.25), *Tillandsia usenoides* (936.75), *Nicotiana glauca* (771.51), *Salvia hispanica* (479.51), *Salix lasiolepis* (444.872), *Ruta graveolens* (458.62), *Olea europea* (406.30), *Ocimum basilicum* (405.13).

• Species with high Zinc (Zn) content ($\mu\text{g g}^{-1}$ dwt)

Salix lasiolepis (216.31), *Ocimum basilicum* (118.77), *Rosamrinus officinalis* (86.07), *Salvia hispanica* (62.22), *Phalaris canariensis* (63.97), *Tragia ramosa* (57.87), *Carya illinoensis* (57.69), *Agave macroculmis Todaro* (55.20), *Gnaphalium canescens* (53.73), *Hedeoma palmeri* (53.54), *Melia azadirachta* (52.57), *Ruta graveolens* (50.72), *Opuntia ficus-indica* (50.05), *Phoradendron villosum* (52.02).

Discussion: The medicinal plants used traditionally by Mexicans for various diseases such as for diabetes, few for stomach ache and others in various diseases such as cancer, bronchitis, infertility, inflammation, blood circulation, digestion, fever, kidney problem,

Particular diseases	Common name	Scientific name	Family	Type
Diabetes	Maguey Tódaro	<i>Agave macroculmis Todaro</i>	Agavaceae	Rosetofilius
Diabetes	Madroño	<i>Arbutus xalapensis</i>	Ericaceae	Bush
Diabetes	Pata de vaca	<i>Bauhinia forficata</i>	Fabaceae	Tree
Diabetes	Tepozan	<i>Buddleja cordata</i>	Buddlejaceae	Tree
Diabetes	Nogal	<i>Carya illinoensis</i>	Juglandaceae	Tree
Diabetes	Palo blanco	<i>Celtis laevigata</i>	Ulmaceae	Tree
Diabetes	Canela	<i>Cinnamomum verum</i>	Lauraceae	Tree
Diabetes	Salvia	<i>Croton suaveolens</i>	Euphorbiaceae	Bush
Diabetes	Níspero	<i>Eryobotria japonica</i>	Rosaceae	Bush
Diabetes	Betónica o poleo de hoja ancha	<i>Hedeoma palmeri</i>	Lamiaceae	Bush
Diabetes	Manrubbio	<i>Marrubium vulgare</i>	Lamiaceae	Herb
Diabetes	Neem	<i>Melia azadirachta</i>	Meliaceae	Tree
Diabetes	Moringa	<i>Moringa oleifer</i>	Moriginaceae	Tree
Diabetes	Nopal de t. año	<i>Opuntia ficus-indica</i>	Cactaceae	Bush
Diabetes	Injerto	<i>Phoradendron villosum</i>	Viscaceae	Bush
Diabetes	Lantrisco	<i>Rhus virens</i>	Anacardiaceae	Bush
Diabetes	Sauce	<i>Salix lasiolepis</i>	Salicaceae	Tree
Diabetes	Tronadora	<i>Tecoma stans</i>	Bignoniaceae	Bush
Diabetes	Maguey Tódaro	<i>Agave macroculmis Todaro</i>	Agavaceae	Rosetofilius
Ulcer, tumour, respiratory problems	Uña de gato	<i>Acacia wrightii</i>	Mimosaceae	Tree
Cancer	Hierba del cáncer	<i>Acalypha monostachya</i>	Euphorbiaceae	Herb
Bronchitis, infertility, inflammation	Hierba de San Nicolás	<i>Chrysactinia mexicana</i>	Asteraceae	Bush
Circulation, digestion	Chaya	<i>Cnidoscolus aconitifolius</i>	Euphorbiaceae	Tree
Fever	Parraleña	<i>Dyssodia setifolia</i>	Asteraceae	Herb
Kidney, cancer	Cola de caballo	<i>Equisetum hyemale</i>	Equisetaceae	Stem erect
Stomach ache	Colesia	<i>Eruca sativa</i>	Brassicaceae	Herb
Cholesterol	Hierba del sapo	<i>Eryngium heterophyllum</i>	Apiaceae	Herb
Tonic, stimulant	Clavo de olor	<i>Eugenia caryophyllata</i>	Myrtaceae	Tree
Expectorant	Gordolobo	<i>Gnaphalium canescens</i>	Asteraceae	Herb
Stomachache	Ocotillo	<i>Gochnatia hypoleuca</i>	Asteraceae	Bush
Stomachache	Hierba del pajarito	<i>Lepidium virginicum</i>	Brassicaceae	Herb
Stomachache	Laurel	<i>Litsea glauscesens</i>	Lauraceae	Bush
Stomachache	Yerbabuena	<i>Mentha piperita</i>	Lamiaceae	Herb
Kidney	Charrasquilla	<i>Mimosa malacophylla</i>	Leguminosae	sub Bush
Headache, gum pain, joint pain	Gigante	<i>Nicotiana glauca</i>	Solanaceae	Bush
Earache, taquicardia	Albahaca	<i>Ocimum basilicum</i>	Lamiaceae	herb
Cholesterol	Olivo	<i>Olea europea</i>	Oleaceae	Tree
Circulation, blood pressure	Alpistle	<i>Phalaris canariensis</i>	Poaceae	Bush
Cough, expectorante	Orégano	<i>Poliomintha longiflora</i>	Lamiaceae	Bush
Circulation, hairfall	Romero	<i>Rosamrinus officinalis</i>	Lamiaceae	Bush
Earache, blood pressure	Ruda	<i>Ruta graveolens</i>	Rutaceae	Herb
Cancer, coagulation, antitumour, elimination of fat	Chia	<i>Salvia hispanica</i>	Lamiaceae	Herb
Somachache, antifungal	Tatalencho	<i>Tagetes lucida</i>	Asteraceae	sub Bush
Burning	Paistle	<i>Tillandsia usenoides L.</i>	Bromeliaceae	Caulscent
Bronchitis blood purification	Ortiguilla	<i>Tragia ramosa</i>	Euphorbiaceae	Herb

Table 2: List of medicinal plants used to determine the leaf nutrient content.

Plant Species	Leaf Macro-nutrient Content						Leaf Micro-nutrient Content			
	(mg g ⁻¹ dw)			(%)			% Protein	(µg g ⁻¹ dw)		
	K	Mg	P	C	N	C:N		Cu	Fe	Zn
<i>Agave macroculmis Todaro</i>	78.45 ± 1.07	1.38 ± 0.29	0.73 ± 0.02	41.32 ± 0.74	1.36 ± 0.21	31.05 ± 5.11	8.49	18.76 ± 2.87	210.53 ± 17.37	55.20 ± 3.39
<i>Arbutus xalapensis</i>	30.65 ± 13.9	0.54 ± 0.60	1.78 ± 0.40	49.1 ± 0.42	1.86 ± 0.30	26.94 ± 3.72	11.6	33.40 ± 20.18	347.58 ± 104.2	25.07 ± 2.65
<i>Bauhinia forficata</i>	8.29 ± 1.27	1.45 ± 0.31	1.37 ± 0.11	34.02 ± 2.32	2.33 ± 0.6	14.60 ± 3.86	14.56	12.86 ± 0.73	161.65 ± 5.30	10.73 ± 0.28
<i>Buddleja cordata</i>	39.30 ± 1.82	0.17 ± 0.09	0.56 ± 0.10	45.70 ± 0.56	3.26 ± 0.40	14.16 ± 1.44	20.38	29.90 ± 1.85	148.00 ± 26.25	40.26 ± 3.39
<i>Carya illinoensis</i>	31.16 ± 1.89	0.85 ± 0.34	2.89 ± 0.06	44.27 ± 1.00	3.76 ± 0.71	12.04 ± 1.81	23.5	25.74 ± 1.70	166.49 ± 20.45	57.69 ± 7.74
<i>Celtis laevigata</i>	16.10 ± 4.59	1.51 ± 0.67	4.03 ± 0.29	39.45 ± 0.51	3.01 ± 0.18	13.13 ± 0.70	18.83	33.88 ± 12.60	213.15 ± 49.80	23.53 ± 1.91
<i>Cinnamomum verum</i>	16.14 ± 1.35	0.27 ± 0.03	0.53 ± 0.05	49.34 ± 0.48	2.49 ± 0.20	19.89 ± 1.70	15.59	24.53 ± 2.77	217.14 ± 9.90	9.49 ± 1.47
<i>Croton suaveolens</i>	75.62 ± 3.67	0.22 ± 0.09	2.43 ± 0.14	45.17 ± 0.35	2.33 ± 0.53	20.16 ± 4.52	14.58	26.87 ± 1.66	229.13 ± 24.25	34.55 ± 4.11
<i>Eryobotria japonica</i>	18.77 ± 1.68	1.78 ± 0.36	2.20 ± 0.20	47.98 ± 1.18	3.03 ± 0.35	15.98 ± 1.58	18.94	22.04 ± 3.44	177.91 ± 13.45	17.13 ± 1.56
<i>Hedeoma palmeri</i>	76.50 ± 1.42	0.18 ± 0.14	1.40 ± 0.03	46.38 ± 1.66	2.83 ± 0.78	17.14 ± 3.33	17.69	23.98 ± 1.20	334.23 ± 9.96	53.54 ± 2.88
<i>Marrubium vulgare</i>	91.27 ± 3.70	0.64 ± 0.42	1.85 ± 0.05	40.48 ± 0.32	4.56 ± 0.58	8.99 ± 1.03	28.48	25.14 ± 1.07	374.78 ± 13.18	46.79 ± 3.07
<i>Melia azadirachta</i>	90.99 ± 7.21	3.41 ± 0.80	1.98 ± 0.35	45.12 ± 0.87	5.85 ± 0.32	7.73 ± 0.32	36.55	24.20 ± 5.26	265.59 ± 21.75	52.57 ± 11.28
<i>Moringa oleifer</i>	95.59 ± 7.56	0.81 ± 0.89	1.91 ± 0.15	45.96 ± 0.23	6.25 ± 0.25	7.37 ± 0.31	39.05	10.59 ± 2.22	773.04 ± 198.33	26.74 ± 4.75
<i>Opuntia ficus-indica</i>	101.47 ± 9.1	6.39 ± 0.90	0.84 ± 0.07	25.54 ± 0.99	2.36 ± 0.43	11.1 ± 1.94	14.74	22.76 ± 1.66	135.18 ± 10.44	50.05 ± 4.80
<i>Phoradendron villosum</i>	100.58 ± 7.6	2.29 ± 0.68	2.40 ± 0.04	40.4 ± 0.63	4.92 ± 0.20	8.22 ± 0.44	30.76	25.30 ± 1.28	151.30 ± 10.06	52.02 ± 6.85
<i>Rhus virens</i>	14.77 ± 2.64	0.33 ± 0.15	1.41 ± 0.26	50.34 ± 0.59	2.27 ± 0.45	22.92 ± 4.67	14.19	22.98 ± 6.24	98.28 ± 23.19	12.75 ± 1.54
<i>Salix lasiolepis</i>	15.66 ± 3.34	1.24 ± 0.17	1.19 ± 0.07	33.37 ± 4.58	2.06 ± 0.50	16.24 ± 9.16	12.88	8.73 ± 3.27	444.82 ± 24.76	216.31 ± 10.83
<i>Tecoma stans</i>	57.22 ± 8.90	0.31 ± 0.11	1.36 ± 0.02	48.79 ± 1.21	3.28 ± 0.47	15.17 ± 2.34	20.47	25.67 ± 3.33	263.66 ± 32.88	29.49 ± 1.27
<i>Acacia wrightii</i>	12.23 ± 2.97	0.49 ± 0.28	0.99 ± 0.21	36.59 ± 1.11	3.96 ± 0.18	9.25 ± 6.22	24.75	11.51 ± 12.73	196.38 ± 15.07	11.53 ± 6.33
<i>Acalypha monostachya</i>	25.70 ± 1.86	8.11 ± 0.88	1.58 ± 0.04	29.71 ± 3.28	1.86 ± 0.26	15.95 ± 12.53	11.64	9.80 ± 1.50	275.96 ± 36.79	64.74 ± 3.01
<i>Chrysactinia Mexicana</i>	5.47 ± 0.81	0.57 ± 0.36	0.86 ± 0.05	45.04 ± 0.48	3.39 ± 0.49	13.56 ± 2.46	21.2	12.53 ± 1.69	231.53 ± 21.08	30.30 ± 3.09
<i>Cnidoscopus aconitifolius</i>	39.96 ± 0.98	2.28 ± 0.36	2.05 ± 0.06	39.11 ± 2.20	4.34 ± 0.08	9.02 ± 2.75	27.09	29.67 ± 1.04	354.72 ± 20.59	52.01 ± 4.65
<i>Dyssodia setifolia</i>	11.12 ± 2.24	0.58 ± 0.44	1.06 ± 0.29	39.68 ± 1.35	2.35 ± 0.67	18.59 ± 7.38	14.67	25.25 ± 1.45	3540.19 ± 557.14 ??	27.97 ± 1.25
<i>Equisetum hyemale</i>	12.86 ± 0.86	2.28 ± 0.38	1.51 ± 0.61	26.95 ± 1.09	1.81 ± 0.08	14.89 ± 12.82	11.31	9.97 ± 1.63	229.29 ± 8.67	107.44 ± 17.64
<i>Eruca sativa</i>	144.23 ± 2.3	2.22 ± 0.24	2.22 ± 0.26	41.13 ± 0.72	5.48 ± 0.64	7.59 ± 0.90	34.23	22.54 ± 8.27	312.46 ± 60.02	45.51 ± 1.96
<i>Eryngium heterophyllum</i>	43.07 ± 8.87	0.38 ± 0.29	2.27 ± 0.11	40.9 ± 0.65	1.75 ± 0.40	24.23 ± 4.89	10.94	33.30 ± 12.18	144.26 ± 57.95	15.45 ± 3.49
<i>Eugenia caryophyllata</i>	37.73 ± 1.40	0.97 ± 0.34	0.89 ± 0.08	51.66 ± 1.85	2.9 ± 0.35	18.01 ± 2.30	18.14	27.33 ± 1.25	221.34 ± 156.74	12.27 ± 0.56
<i>Gnaphalium canescens</i>	16.69 ± 0.53	1.40 ± 0.18	1.08 ± 0.26	37.73 ± 1.26	2.56 ± 0.32	14.89 ± 1.37	15.99	25.13 ± 1.70	3973.55 ± 1342.28	53.73 ± 0.79
<i>Gochnatia hypoleuca</i>	22.54 ± 2.48	0.48 ± 0.25	0.82 ± 0.04	49.86 ± 0.87	3.59 ± 0.50	14.11 ± 1.89	22.41	11.37 ± 0.89	292.92 ± 23.68	24.43 ± 0.77
<i>Lepidium virginicum</i>	11.67 ± 0.59	1.89 ± 0.52	2.27 ± 0.05	43.8 ± 1.22	4.46 ± 0.59	9.95 ± 1.11	27.85	5.36 ± 1.80	188.69 ± 7.64	28.26 ± 1.49
<i>Litsea glauscescens</i>	10.94 ± 2.47	0.33 ± 0.28	2.40 ± 0.13	51.34 ± 0.28	3.36 ± 0.45	15.5 ± 2.03	21	22.84 ± 7.79	177.31 ± 27.21	33.09 ± 2.68
<i>Lepidium virginicum</i>	58.22 ± 4.35	1.15 ± 0.24	2.20 ± 0.03	44.14 ± 2.71	5.4 ± 0.15	8.18 ± 0.41	33.73	31.06 ± 0.74	373.38 ± 9.40	49.17 ± 5.46
<i>Mimosa malacophylla</i>	16.39 ± 0.45	8.64 ± 0.99	0.84 ± 0.04	45.15 ± 0.53	8.46 ± 0.18	5.34 ± 0.17	52.87	4.17 ± 1.25	222.28 ± 3.85	35.28 ± 1.14
<i>Nicotiana glauca</i>	18.19 ± 1.32	1.31 ± 0.15	1.30 ± 0.12	37.94 ± 0.56	4.79 ± 0.54	8.00 ± 0.87	29.91	33.06 ± 2.95	721.51 ± 234.69	43.64 ± 3.96
<i>Ocimum basilicum</i>	72.09 ± 2.43	2.42 ± 0.09	2.18 ± 0.03	38.31 ± 0.34	4.66 ± 0.45	8.29 ± 0.84	29.1	14.56 ± 0.79	405.13 ± 34.32	118.77 ± 5.93
<i>Olea europea</i>	48.16 ± 1.21	0.80 ± 0.20	1.23 ± 0.11	41.13 ± 2.59	1.86 ± 0.8	22.08 ± 3.23	11.64	7.91 ± 1.13	406.30 ± 31.86	21.93 ± 0.82
<i>Phalaris canariensis</i>	163.35 ± 6.8	2.74 ± 0.30	1.48 ± 0.17	40.73 ± 0.53	2.84 ± 0.52	14.78 ± 2.87	17.73	33.78 ± 8.95	254.84 ± 70.34	63.97 ± 7.56
<i>Poliomintha longiflora</i>	31.63 ± 5.50	1.80 ± 0.81	1.45 ± 0.75	42.9 ± 0.24	4.89 ± 0.21	8.79 ± 0.36	30.54	29.04 ± 2.79	316.63 ± 130.27	32.99 ± 7.99
<i>Rosmarinus officinalis</i>	26.59 ± 5.54	1.18 ± 0.28	1.24 ± 0.38	47.77 ± 5.43	4.54 ± 0.24	10.57 ± 1.65	28.41	11.36 ± 7.39	336.95 ± 122.63	86.07 ± 21.27
<i>Ruta graveolens</i>	28.04 ± 2.59	0.88 ± 0.20	1.24 ± 0.20	38.86 ± 1.4	2.79 ± 0.25	13.23 ± 5.7	17.41	20.87 ± 9.55	458.62 ± 71.30	50.72 ± 11.64
<i>Salvia hispanica</i>	46.48 ± 6.70	3.52 ± 0.71	5.79 ± 0.75	44.68 ± 2.18	5.24 ± 0.62	8.59 ± 0.81	32.77	28.11 ± 4.75	479.51 ± 64.24	62.22 ± 3.93
<i>Tagetes lucida</i>	21.49 ± 3.31	1.90 ± 0.13	1.40 ± 0.04	46.19 ± 1.04	5.89 ± 0.29	7.85 ± 0.33	36.81	8.93 ± 2.40	167.40 ± 13.81	12.16 ± 1.64
<i>Tillandsia usenoides L.</i>	40.56 ± 4.67	1.20 ± 0.14	4.45 ± 0.17	44.10 ± 1.61	1.56 ± 0.71	31.32 ± 8.20	9.75	31.70 ± 24.04	936.75 ± 543.65	34.06 ± 9.45
<i>Tragia ramosa</i>	60.57 ± 10.1	0.46 ± 0.42	1.45 ± 0.10	42.68 ± 1.16	3.89 ± 0.63	11.22 ± 2.01	24.31	24.96 ± 2.87	1450.25 ± 273.89	57.87 ± 5.81

Data are means and standard deviation (n=5).

Table 3: Leaf nutrient content of the 44 species used to cure particular diseases.

earache, cholesterol, cough, respiratory headache, circulation, burning [18-20,22]. The supply of macro- and micronutrients has been found to be useful in alleviating various diseases. Several medicinal plants used to reduce blood sugar such as Mulberry leaf, soybean, black bean, Persian shallot, *Zizyphus lotus*, etc., such as in alleviating eye problem [23] and diabetes [24,25].

It may be stated here that the efficacy of the species used for different diseases is not confirmed scientifically, though few studies have been directed on the chemistry of few species. Some Mexican medicinal plants containing β -glycosides and other compounds are effective in lowering blood sugar and body weight in diabetic patients. The results reveal great variability in nutrient profile among the medicinal species. The species with high nutrients could be verified for their efficacy. In other words, these variation studies in macro and micronutrient contents of 44 medicinal plant species were studied in medicinal plants that are traditionally utilized to control diabetes and other diseases in Nuevo Leon, Mexico. It is observed that most of the species used traditionally to combat various diseases contain very high amount of macro and micronutrients and protein confirming the efficacy of these species. In this respect we have no scope to compare the contents of nutrient profile with individual disease species wise. In addition to micronutrients some major elements are necessary in higher amounts for good health. In this respect phosphorus is required for many biochemical reactions in the body, such as conversion of foods to energy, muscle contractions, nerve conduction, normal kidney functions and helps to build strong bones [38].

It is reported that most of American diets are adequate in magnesium (Mg). Magnesium is an activator of numerous enzyme systems to control carbohydrate, fat and electrolyte metabolism, nucleic acid, protein synthesis, membrane integrity and transport [39].

In the case of diabetes we want to mention the nutraceutical values of few species commonly used in combating diabetes. In this context research on nutraceutical plants is a modern trend. The nutraceutical values of various plants are well documented in the literature. It is evident from this study that the species which are commonly used to combat diabetes contain very high amount of nutrients, thereby confirming their efficacy, but the efficacy of these species should be confirmed by analyzing the tolerance level in a pharmacology laboratory. On the other hand may be cited the use of *Tragia ramosa* commonly used traditionally in blood purification contain very high amount of Fe (1250.25) confirming the role of iron as blood purifier and hemoglobin function. In the present study few medicinal plants were selected viz. *Gnaphalium canescens* (3973.55), *Tragia ramosa* (1450.25), *Tillandsia usenoides* (936.75), *Nicotiana glauca* (771.51), *Salvia hispanica* (479.51), for having very high amount of Fe which may be tested for blood purification. On the other hand, the exceptional high amount of iron in the case of *Dyssodia setifolia*, containing 3540.19 could be confirmed for its possible efficacy in blood purification, hemoglobin function, blood circulation through pharmacological study.

Research on medicinal plants could follow the following steps

- Several medicinal plants are used to alleviate a disease. There is a necessity to select species of maximum use.
- Chemical analysis of the species and selection of species with high nutrient profile.
- Analysis of chemical composition (ingredients) of the selected species.
- Final selection and confirmation of the species for its efficacy.

- Analysis of the tolerance level of the selected species at the laboratory.
- Establish a germplasm bank with high nutritional values.

Conclusion

It is assessed from the present study that most of the species used traditionally to combat various diseases contain high amount of nutrients of high nutraceutical values, thereby confirming the efficacy of these medicinal species, justifying the importance of these in ethnomedicine. Based on the analysis of macro and micro nutrients the following species *Phalaris canariensis*, *Eruca sativa*, *Ocimum basilicum*, *Tragia ramosa*, *Mentha piperita*, *Mimosa malacophylla*, *Acalypha monostachya*, *Salvia hispanica*, *Tillandsia usenoides*, *Litsea glauscens*, *Tagetes lucida*, *Erygium heterophylla*, *Dyssodia setifolia*, *Nicotiana glauca*, *Ruta graveolens*, *Olea europea*, *Equisetum hyemale*, *Rosmarinus officinalis* are selected containing a majority of each of these nutrients. In our study we obtained few species with high C/N ratio having high potentialities of the production of secondary metabolites and antioxidants exhibiting as the potential lines of research in future. It is suggested that the efficacy of the species with high nutritional values in alleviating various diseases need to be confirmed in the pharmacology laboratory via tolerance level. It may be stated here that the high amount of nutrients reported in this study in the studied medicinal plants are rarely available in food grains and vegetables. Therefore the high nutritional values of these medicinal species could serve as enriched sources of various macro- and micronutrients for our health improvement. There is a great necessity of inter-multidisciplinary research to harness high nutraceuticals reported in the present study to improve health as well as to alleviate various diseases.

Acknowledgements

Valuable technical assistance provided by Elsa Dolores González Serna and Manuel Hernández Charles is gratefully recognized. This research was funded by Consejo Nacional de Ciencia y Tecnología (CONACYT, Grant 250732, Convocatoria de Investigación Ciencia Básica 2015). Useful suggestions from two anonymous reviewers helped to improve the manuscript.

References

1. Ramirez GRG, Lozano (2014) Native shrubs: Edible foliage for small ruminants. In: Applied Botany, Ratikanta M, et al. (eds). Pustaka Publishing Company, Kolkata, India, pp: 165-180.
2. Wright IJ, Reich PB, Westoby M (2001) Strategy shifts in leaf physiology, structure and Nutrient content between species of high- and low-rainfall and high- and low-nutrient habitats. *Functional Ecology* 15: 423-434.
3. Reich A, Holbrook NM, Ewel JJ (2004) Developmental and physiological correlates of leaf size in *Hyeronima alchorneoides* (Euphorbiaceae). *Am J Bot* 91: 582-589.
4. Chapin FS (1982) The mineral nutrition of wild plants. *Annu Rev Ecol Evol Syst* 19: 233-260.
5. Chapin FS, Schulze ED, Mooney HA (1990) The ecology and economics of storage in plants. *Annu Rev Ecol Evol Syst* 21: 423-447.
6. Ackerly DD, Bazzaz FA (1995) Leaf dynamics, self-shading and carbon gain in seedlings of a tropical pioneer tree. *Oecologia* 101: 289-298.
7. Hikosaka K (1996) Effects of leaf age, nitrogen nutrition and photon flux density on the organization of the photosynthetic apparatus in leaves of a vine (*Ipomoea tricolor* Cav.) grown horizontally to avoid mutual shading of leaves. *Planta* 198: 144-150.
8. Nooden LD (1988) The phenomena of senescence and ageing. In: *Senescence and Ageing in Plants* (LD Nooden & AC Leopold Eds.). Academic Press, USA.
9. Aerts R (1996) Nutrient resorption from senescing leaves of perennials: Are there general patterns? *Journal of Ecology* 84: 597-608.
10. Negi GCS, Sing SP (1993) Leaf nitrogen dynamics with particular reference to retranslocation in evergreen and deciduous tree species in Kumaun Himalaya.

- Can J For Res 23: 349-357.
11. Rengel Z, Marschner P (2005) Nutrient availability and management in the rhizosphere: exploiting genotypic differences. *New Phytol* 168: 305-312.
 12. Lukhele MS, van Ryssen JBJ (2003) The chemical composition and potential nutritive value of the foliage of four subtropical tree species in southern Africa for ruminants. *S Afr J Anim Sci* 33: 132-141.
 13. Huerta C (1997) La herbolaria: mito o realidad. *Biodiversitas. Conabio No.* 13.
 14. Villaseñor JL (2003) Diversidad y distribución de las Magnoliophyta de México. *Interciencia* 28: 160-167.
 15. Vega-Avila E, Espejo-Serna A, Alarcón-Aguilar F, Velasco-Lezama R (2009) Cytotoxic activity of four Mexican medicinal plants. *Proc West Pharmacol Soc* 52: 78-82.
 16. Bell WH, Castetter EF (1997) The utilization of mesquite and screwbean by the aborigines in the American southwest. *University of New Mexico biological series* 5: 1-55.
 17. Hernández-Sandoval LC, González and González-Medrano F (1991) Plantas útiles de Tamaulipas, México. *Anales del Instituto de Biología de la Universidad Nacional Autónoma de México. Serie Botánica* 62: 1-38.
 18. Eduardo EC, Villarreal JAQ, Alfonso DS, Marisela PM, Laura SM, et al. (2006) Diversity and distributional patterns of legumes in Southern Nuevo León, México. *Southwest Nat* 51: 1-10.
 19. Estrada E, Villarreal JA, Cantú C, Cabral I, Scott L, et al. (2007) Ethnobotany in the Cumbres de Monterrey National Park, Nuevo León, México. *J Ethnobiol Ethnomed* 3: 8.
 20. Estrada EC, Soto BEM, Garza LM, Villarreal JAQ, Jiménez JP, et al. (2012) Plantas útiles en el centro-sur del estado de Nuevo León. *Bot Sci* 91: 381.
 21. Estrada-Castillón E, Soto-Mata BE, Garza-López M, Villarreal-Quintanilla JÁ, Jiménez-Pérez J, et al. (2012) Medicinal plants in the southern region of the State of Nuevo León, México. *J Ethnobiol Ethnomed* 8: 45.
 22. Villarreal-Quintanilla JA (2012) Plantas Medicinales de Miquihuana, Tamaulipas. *UANL* 99: 146.
 23. Kowluru RA, Kanwar M, Chan PS, Zhang JP (2008) Inhibition of retinopathy and retinal metabolic abnormalities in diabetic rats with AREDS-based micronutrients. *Arch Ophthalmol* 126: 1266-1272.
 24. Patton SR, Dolan LM, Powers SW (2007) Dietary adherence and associated glycemic control in families of young children with type 1 diabetes. *J Am Diet Assoc* 107: 46-52.
 25. Farvid MS, Homayouni F, Amiri Z, Adelmanesh F (2011) Improving neuropathy scores in type 2 diabetic patients using micronutrients supplementation. *Diabetes Res Clin Pract* 93: 86-94.
 26. Devasagayam TP, Tilak JC, Boloor KK, Sane KS, Ghaskadbi SS, et al. (2004) Free radicals and antioxidants in human health: current status and future prospects. *J Assoc Physicians India* 52: 794-804.
 27. Serafini M, Bellocchio R, Wolk A, Ekström AM (2002) Total antioxidant potential of fruit and vegetables and risk of gastric cancer. *Gastroenterology* 123: 985-991.
 28. Mandel S, Youdim MB (2004) Catechin polyphenols: neurodegeneration and neuroprotection in neurodegenerative diseases. *Free Radic Biol Med* 37: 304-317.
 29. Shoskes DA, Zeitlin SI, Shahed A, Rajfer J (1999) Quercetin in men with category III chronic prostatitis: a preliminary prospective, double-blind, placebo-controlled trial. *Urology* 54: 960-963.
 30. Shaheen SO, Sterne JA, Thompson RL, Songhurst CE, Margetts BM, et al. (2001) Dietary antioxidants and asthma in adults: population-based case-control study. *Am J Respir Crit Care Med* 164: 1823-1828.
 31. Halliwell B, Gutteridge JMC (1989) *Free radicals in Biology and Medicine*. 2nd edn. Clarendon Press, Oxford, UK.
 32. Gülçin I (2005) The antioxidant and radical scavenging activities of black pepper (*Piper nigrum*) seeds. *Int J Food Sci Nutr* 56: 491-499.
 33. Schramm DD, Karim M, Schrader HR, Holt RR, Cardetti M, et al. (2003) Honey with high levels of antioxidants can provide protection to healthy human subjects. *J Agric Food Chem* 51: 1732-1735.
 34. Lindroth RL, Osien TL, Burnhill HRH, Wood SA (2002) Effects of genotype and nutrient availability on photosensitivity of trembling aspen (*Populus tremoides* Michx.) during leaf senescence. *Biochem Sys Ecol* 30: 297-307.
 35. Pandey N, Meena RP, Rai SK, Rai SP (2011) Medicinal plants derived nutraceutical: A reemerging health aid. *International Journal of Plants and Biosciences* 2: 419-441.
 36. Cherney DJR (2000) Characterization of forages by chemical analysis. In: *Forage evaluation in ruminant nutrition*. Givens DJ, Owen E, Axford RFE, Omed HM (eds.). CAB International, Wallingford, pp: 281-300.
 37. Association of Official Analytical Chemists (2016) *Official Methods of Analysis of AOAC International*. 20th Edition, Washington, DC, USA.
 38. Ishreen L (2014) *Mineral Guide - Phosphorus requirements and dietary sources*.
 39. Seelig MS (1982) *Magnesium Requirements in Human Nutrition*. Contemporary Nutrition 7: 45-58.

Citation: Maiti R, Rodriguez HG, Kumari A (2016) Nutrient Profile of Native Woody Species and Medicinal Plants in Northeastern Mexico: A Synthesis. *J Bioprocess Biotech* 6: 283. doi:[10.4172/2155-9821.1000283](https://doi.org/10.4172/2155-9821.1000283)

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>