

## Nutritional Status of Green Mussel *Perna Viridis* at Tamil Nadu, Southwest Coast of India

Saritha K<sup>1\*</sup>, Mary D<sup>2</sup> and Patterson J<sup>2</sup>

<sup>1</sup>Suganthi Devadason Marine Research Institute, Tuticorin, Tamil Nadu, India

<sup>2</sup>St.Jude's College, Thoothoor, Tamil Nadu, India

### Abstract

The biochemical compositions of green mussel *P. viridis* were analyzed. The percentage of protein was high followed by the carbohydrate and lipid. The total essential amino acids were found to be higher than the non-essential amino acids. Seven vitamins were detected and among that vitamin A and C were found to be high. Totally six macro minerals and 2 trace minerals were reported. The result showed that marine bivalve *P. viridis* is a valuable food source for human consumption, due to its high quality protein and well-balanced nutritional composition.

**Keywords:** Mussel; Essential amino acids; Vitamins; Minerals; Biochemical compositions

### Introduction

Mussels of the genus *Perna* Philipsson belong to the Mytilidae or true mussels (Mollusca; Bivalvia; Lamellibranchia; Mytiloida; Mytilidae). This genus contains green and brown mussels and are distributed in tropical, subtropical, warm temperate and cold temperate regions, mostly from the southern hemisphere, but also from northern Africa and the northern coasts of South America [1,2]. In India both the green and brown mussels are widely distributed in the east and west coast of India along the inter-tidal zones [3]. These mussels are both ecologically and economically important throughout their ranges, and have long constituted an important source of human food [4-8].

The knowledge on nutritional composition of any edible organisms is enormously important because of the nutritive value is reflected to the biochemical contents [9]. The demand for protein rich food is increasing, especially in developing countries, stimulating the exploration of unexploited or non-traditional resources. Marine molluscs are commercially valuable species and easy to cultivate in coastal areas. Marine molluscs are important for marine ecology and human diet, since it is an important source of nutrients. Consumption of marine molluscs provides an inexpensive source of protein with a high biological value, essential minerals and vitamins. Additionally, the molluscs muscle contains little saturated fat and significant amount of Vitamin C. Molluscs is also a good source of minerals such as calcium, potassium, zinc, iron, phosphorus and copper.

Consumption of shellfish is considerably low but the bivalve mussels consumption in western coast area in India is more just like other sea foods. Shellfish contain significant amounts of "good" fats called omega-3 fatty acids. Shellfish such as mussel contains approximately 20 to 28 percent calories from fat [10]. The types of fat in mussels are also favorable. Shellfish also provide high quality protein with all the dietary essential amino acids for the maintenance and growth of human body. For this reason, shellfish should be considered a low-fat, high-protein food that can be included in a low-fat diet [10,11]. The demand for protein rich food is increasing day by day especially in developing countries with the growth of the human population. Marine fin fishes occupy an important place in human diet followed by crustaceans and molluscs to a much smaller extent [12].

As the world population is growing, the per capita consumption of seafood is also increasing rapidly. Because of health consciousness,

the modern day man is interested in taking seafood more in view of its nutritional superiority than all other sources of food accessible to him. There is no considerable study on the nutritive value of green mussel *P. viridis*. In the present study the major biochemical components like total protein, carbohydrate, lipid, amino acids, vitamins and mineral contents of green mussel meat were assessed using standard methods.

### Materials and Methods

The green mussel *P. viridis* were collected from the intertidal rocky coastal areas of Kanyakumari district in the west coast of India. Samples were brought to the laboratory and the shells were broken and the soft tissues were removed from the shell and dried at 50°C in an oven and used for biochemical analysis. The protein, lipid carbohydrate content of the samples was analyzed in triplicates. The proximate compositions of the experimental samples were determined by using standard methods; viz., protein [13], carbohydrate [14] and lipid [15]. The dried mussel meat samples were finely ground for estimating the amino acids using HPLC (Merck Hitachi L-7400) following the method of Baker and Han [16]. Fat soluble vitamin A and the water soluble vitamins B1, B2, B3, B6, and C were analysed in HPLC (Merck Hitachi L-74000) following the method described by Sadasivam and Manickam [17]. The vitamin B12 was estimated by following methods suggested in USP NF 2000 Asian edition. The minerals were estimated by following the method of Guzman and Jimenez [18].

### Results

#### Biochemical composition

The results of biochemical composition of *P. viridis* are shown in Table 1. While protein, carbohydrate and lipid content of green mussel

**\*Corresponding author:** Saritha K, Suganthi Devadason Marine Research Institute, Tuticorin, Tamil Nadu, India, Tel: 914612336488; Fax: 914612325692; E-mail: saritha23kailasam@gmail.com

**Received** August 20, 2015; **Accepted** September 21, 2015; **Published** September 28, 2015

**Citation:** Saritha K, Mary D, Patterson J (2015) Nutritional Status of Green Mussel *Perna Viridis* at Tamil Nadu, Southwest Coast of India. J Nutr Food Sci S14: 003. doi:10.4172/2155-9600.S14-003

**Copyright:** © 2015 Saritha K, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

| Proximate composition | Mean% ± SD (W/W) |
|-----------------------|------------------|
| Protein               | 36.15 ± 1        |
| Carbohydrate          | 24.54 ± 1        |
| Lipid                 | 19.720 ± 2       |

Results are means ± standard deviation of triplicates.

**Table 1:** Proximate composition of *P. viridis* meat powder.

| Essential amino acids | Mean % ± SD (W/W) | Nonessential amino acids | Mean % ± SD (W/W) |
|-----------------------|-------------------|--------------------------|-------------------|
| Phenyl alanine        | 0.993 ± 0.002     | Glycine                  | 0.813 ± 0.11      |
| Lysine                | 1.015 ± 0.005     | Serine                   | 1.354 ± 0.02      |
| Histidine             | 1.154 ± 0.011     | Glutamic acid            | 1.93 ± 0.170      |
| Methionine            | 2.145 ± 0.090     | Cystine                  | 1.134 ± 0.034     |
| Arginine              | 1.193 ± 0.04      | Glutamine                | 0.935 ± 0.02      |
| Leucine               | 1.193 ± 0.04      | Alanine                  | 0.454 ± 0.104     |
| Threonine             | 1.343 ± 0.07      | Proline                  | 0.934 ± 0.004     |
| Iso leucine           | 1.125 ± 0.021     | Asparagine               | 0.993 ± 0.01      |
| Valine                | 1.093 ± 0.002     | Tyrosine                 | 1.094 ± 0.008     |
| Tryptophan            | 1.235 ± 0.10      | Aspartic acid            | 1.215 ± 0.075     |

Results are means ± standard deviation of triplicates.

**Table 2:** Amino acid content of *P. viridis* meat powder.

were 36.15 ± 1, 24.54 ± 1 and 19.720 ± 2 % respectively. The protein content of *P. viridis* was higher than the other biochemical parameters.

### Amino acids

The percentage of essential and non-essential amino acids results are presented in Table 2. Among the ten essential amino acids methionine was found to be high as 2.145 ± 0.090% whereas phenyl alanine was found to be low as 0.993 ± 0.002%. The remaining amino acids like lysine, histidine, arginine, leucine, threonine, isoleucine, valine and tryptophan were in intermediate level. The non-essential amino acids glutamic acid level was high (1.93 ± 0.170%) and alanine level was low (0.454 ± 0.104%) and rest of the non essential amino acids such as glycine, serine, cystine, glutamine, proline, asparagines, tyrosine and aspartic acid were in intermediate level.

### Vitamins

The vitamin content of the meat of *P. viridis* is shown in Table 3. Among them, vitamin A (14.5 ± 1.1 mg/g) and C (13.6 ± 1.05 mg/g) were found in higher levels and other vitamins available in the order of > Vitamin B<sub>6</sub> > Vitamin B<sub>3</sub> > Vitamin B<sub>12</sub> > Vitamin B<sub>2</sub> > Vitamin B<sub>1</sub>.

### Minerals

Totally 6 macro minerals and 2 trace minerals (Table 4) were detected. The macro mineral calcium, potassium, sodium and iodine were found to be high. Magnesium and iron were also detected in meager level. Trace metals like zinc and copper were in trace level.

### Discussion

Seafood is an important in the diets of many individuals because of their unique nutritional composition. The shellfishes are known to be high in protein, low in fat and low in calories. Shellfishes are nutritionally valuable sources of various minerals and high quality protein [11,19-22]. Protein is essential for the sustenance of life and exists in largest quantity of all nutrients as a component of the human body [23]. Our present research revealed that the protein content is high when comparing to the carbohydrate and lipids. Therefore, the results suggest that the *P. viridis* can be considered as an alternative potential

food crop capable of providing cheap animal protein. The findings here are in agreement with the report of Catedral et al. [24] in green mussel *P. viridis*. Giese [25] reported that protein to be the dominant organic constituent in molluscs. NOAA [26] reported that the protein ranges in shellfishes ranged between 0.4 to 26.3% at different part of the world. Salaskar [27] reported that the nutritional content of *P. viridis* varied between the summer and monsoon season.

Protein value obviously reflected to the essential amino acids concentration. Present study reveals that the mussel meat have high in essential amino acids (12.489%) than that of nonessential amino acids (10.856%). In general, the shellfish has a balanced amount of all essential amino acids required for an adult per day. Present study showed that among the total 20 amino acids Methionine, threonine and tryptophan were high. The result reveals that the green mussel *P. viridis* meat is a potential source of food due to high quality protein, as well as balanced essential amino acids. Ajaya Bhaskar [28] reported that the total amino acids in *Perna viridis* are 95.76% whereas *C. madrasensis* 98.4% and in *M. casta* is 65.17%. Arularasan et al. [29] reported the percentage of essential amino acids was more (80.97%) than those non essential amino acids (15.07%) in the Gastropod, *Strombus canarium*. Babu et al. [30] reported 96.8%, of total amino acid in mesogastropod, *Bursa spinosa* was among that it is having 50.01% of essential amino acids (EAA) 50.01% and 46.79% of nonessential amino acids and 3.2% of unidentified amino acids. Palpandi [12] reported that *Nertia crepidularia* had 68.5% essential amino acids and 31.01% of non-essential amino acids. Periyasamy et al. [31] reported that *B. spirata* had 9.91 mg/g of 10 essential and nonessential amino acids. Vitamins are organic chemical compounds are essential for promoting growth, reproduction and maintenance of normal body health and function. So far flesh of fish and shell fish is not considered to be important sources of vitamin A, but high content of Vitamin A was reported in fishes and shell fishes like eel, mackerel and bivalve. *P. viridis* meat showed the dominance of vitamin A and C. Ajaya Bhaskar [28] estimated the vitamin level from three species of bivalve namely *P. viridis*, *C. madrasensis* and *M. casta* and found vitamin B<sub>1</sub>, B<sub>2</sub> and B<sub>6</sub>. Ozden and Erkan [32] reported that the vitamin B mainly present in muscle of

| Vitamins                | Mean mg/100g ± SD |
|-------------------------|-------------------|
| Vitamin A               | 14.5 ± 1.1        |
| Vitamin B <sub>6</sub>  | 3.45 ± 0.95       |
| Vitamin B <sub>12</sub> | 1.77 ± 0.07       |
| Vitamin B <sub>1</sub>  | 0.993 ± 0.00      |
| Vitamin B <sub>2</sub>  | 1.34 ± 0.09       |
| Vitamin B <sub>3</sub>  | 2.66 ± 1.06       |
| Vitamin C               | 13.6 ± 1.05       |

Results are means ± standard deviation of triplicates.

**Table 3:** Vitamin content.

| Minerals  | Mg/100g ± SD |
|-----------|--------------|
| Sodium    | 156.7 ± 3.2  |
| Calcium   | 235.9 ± 1    |
| Potassium | 156.7 ± 4.1  |
| Magnesium | 24.6 ± 1.1   |
| Iron      | 5.09 ± 0.01  |
| Zinc      | 2.93 ± 0.99  |
| Copper    | 1.225 ± 0.00 |
| Iodine    | 112.7 ± 0.8  |

Results are means ± standard deviation of triplicates.

**Table 4:** Minerals content.




Figure 1: Amino acids in standard.




Figure 2: Amino acids in *P. viridis*.

gonad and eggs, while B2 was detected in the digestive gland, gonad and eggs. Shellfish covered in the present study showed complete vitamin profile as the levels required from good health (Figures 1 and 2).

Minerals also constitute important components of hormones, enzymes and enzyme activators in human nutrition [33]. Mineral components such as sodium, potassium, magnesium, calcium, iron, phosphorus and sulphur are important for human nutrition [34]. Mineral deficiencies can cause biochemical, structural and functional pathologies changes. Present study clearly indicate that calcium was more and copper was low in *P. viridis* whereas other macro-minerals potassium, sodium and iodine were in significant level; magnesium and iron were in negligible level. Coombs [35] reported more than 40% of soluble copper and zinc was present in oyster *Ostrea edulis*. The localized Zn and Fe in lysosomes of a number of cell types were reported in *Mytilus edulis* [36]. Rajagopal et al. [37] reported the importance of Ca, Mg, and K in the human nutrition. Gopakumar reported that the

sea foods in general are excellent sources of I, Ca, P, Na, Fe, Zn and oysters are good sources of Fe and Cu. Minerals in different species of pearl shells of South China Sea were rich, in Ca, P and Zn contents [38].

## Conclusion

In general, seafood is one of the most nutritionally balanced foods. The seafood diet helps to control weight and goes a long way towards preventing heart diseases. Studies on amino acid composition of commercial seafood in India are limited. This might be due to lack of awareness on the benefits of these nutrients particularly from molluscan meat. The results of the present study provide information about the amino acids and vitamin and mineral composition and also suggest the consumption of this bivalve, *P. viridis*. Therefore the balanced and healthy diet is a prerequisite for good health. In the present study clearly indicate that the mussel *P. viridis* can be a balanced diet and contribute good nutritional status.

## Acknowledgement

The authors are thankful to Dr. J.K. Patterson Edward, Director, Suganthi Devadason Marine Research Institute, India for providing us the facilities to carry out the work.

## References

- Gosling EM (2003) Bivalve Molluscs: Biology, Ecology and Culture. Fishing News Books, Blackwell Science. 443.
- Siddall SE (1980) A classification of the genus *Perna* (Mytilidae). Bull Mar Sci 30: 858-870.
- Appukkuttan KK, Mathew J, Thomas KJ (1987) Larval rearing and spat production of the brown mussel *P. indica* Kuriakose and Nair at Vizhinjam, Southwest coast of India. Nat Sem Shellfish Re. Farming, Tuticorin. CMFRI Bull. 42: 337-343.
- Griffiths CL, Branch GM (1997) The exploitation of coastal invertebrates and seaweeds in South Africa: historical trends, ecological impacts and implications for management. Trans R Soc S Afr. 52: 121-148.
- Nichol R (1986) Analysis of midden from N44/215: hard times at Hahei? Pacifc Anthropological Records 37. In: Anderson AJ (edn) Traditional Fishing in the Pacific: Ethnological and Archaeological Papers from the 15th Pacific Science Congress, pp. 179-198.
- Parkington J (1976) Coastal settlement between the mouths of the Berg and Olifants Rivers, Cape Province. S Afr Archaeol Bull 31: 127-140.
- Swadling P (1977) The implications of shellfish exploitation for New Zealand Prehistory. Mankind 11: 11-18.
- Tomalin BJ, Kyle R (1998) Subsistence and recreational mussel (*Perna perna*) collecting in KwaZulu-Natal, South Africa: Wshing mortality and precautionary management. S Afr J Zool 33: 12-22.
- Nagabhushanam R, Mane VH (1978) Seasonal variation in the biochemical composition of *Perna viridis* at Ratnagiri on the West Coast of India. Hydrobiol 57: 69-72.
- Ersoy B, Sereflihan H (2010) The Proximate Composition and Fatty Acid Profiles of Edible Parts of Two Freshwater Mussels. TJFAS 10: 71-74.
- King I, Childs MT, Dorsett C, Ostrander JG, Monsen ER (1990) Shellfish: proximate composition, minerals, fatty acid, and sterols. JADA 90: 677-685.
- Palpandi C (2010) Studies on mollusca, biology, biochemistry and heavy metal accumulation in *Nerita (Dostia) crepidularia* Lamarck, 1822 from mangroves of Vellar estuary South east Coast of India. Ph. D., Thesis, Annamalai University India, p. 318.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the folin phenol reagent. J Biol Chem 193: 265-273.
- Dubois M, Giller KA, Hamilton JK, Roberts RA, Smith F (1956) Colorimetric method for determination of sugars and related substances. Analyst Chem 28: 350-356.
- Floch J, Lees, Sloane-Stanley GH (1956) A simple method for the isolation and purification of total lipids from animal tissues. J Biol Chem 226: 497-509.
- Baker DH, Han Y (1994) Ideal amino acid profile for broiler chicks during first three weeks post hatching. Poult Sci 73: 1441-1447.
- Sadasivam S, Manickam A (1996) Biochemical Methods. (2nd edn), New Age International (p) limited, Publishers 179-186.
- Guzman HM, Jimenez CE (1992) Concentration of coral reefs by heavy metals along the Caribbean coast of central Africa (costarica and panama) Mar Poll Bull 24: 554-561.
- Leu S, Jhaveri SN, Karakoltsidis PA, Constantinides SM (1981) Atlantic mackerel (*Scomberscombrus*, L): seasonal variation in proximate composition and distribution of chemical nutrients. Jour Foo Sci 46: 1635.
- Connor WE, Lin DS (1982) The effects of shellfish in the diet upon the plasma lipid levels in humans. Metabolism, 31: 1046-1051.
- Skonberg DI, Perkins BL (2002) Nutrient composition of green crab (*Carcinus maenus*) leg meat and claw meat. Food Chemistry 72: 401-404.
- USDA (2003) National Nutrient Database for Standard Reference, Release 16. Nutrient Data Laboratory Home Page.
- Okuzumi M, Fujii T (2000) Nutritional and Functional Properties of Squid and Cuttle Fish 35th Anniversary of Commemorative Publication p 223.
- Catedral FFD, Sayson R (1977) Proximate and mineral analysis of mussel from Himalayan, Negros. Unpublished Report, Aquaculture Depi, SEAFDEC, Tigbauan, Iloilo.
- Giese AC (1969) A new approach to the biochemical composition of the mollusc body. Oceanogr Mar Biol Ann Rev 7: 175-229.
- NOAA Technical Report NMFS 55 (1987) Proximate Composition, Energy, Fatty Acid, Sodium, and Cholesterol Content of Finfish, Shellfish, and their Products. National Marine Fisheries Service.
- Salaskar GM, Nayak VN (2011) Nutritional quality of bivalves, *crassostrea madrasensis* and *Perna viridis* in the kali estuary, Karnataka, India. Recent Research in Science and Technology 3: 06-11.
- Ajaya BD (2002) Nutritional evaluation of molluscan seafood. Ph.D. Thesis, Annamalai University, India pp. 25-56.
- Arularasan S, Lyla PS, Kesavan K, Ajmal Khan S (2010) Recieps for the Mesogastropod - *Strombus canarium*. Advan J Food Sci Technol 2: 31-35.
- Babu A, Kesavan K, Annadurai D, Rajagopal S (2010) *Bursa spinosa* - A Mesogastropod Fit for Human Consumption. Advan J Food Sci Technol 2: 79-83.
- Periyasamy N, Srinivasan M, Devanathan K, Balakrishnan S (2011) Nutritional value of gastropod *Babylonia spirata* (Linnaeus, 1758) from Thazhanguda, Southeast coast of India. Asian Paci J Trop Biomed 1: S249-S252.
- Ozden O, Erkan N (2011) A preliminary study of amino acid and mineral profiles of important and estimable 21 seafood species. British Food Journal 113: 457-469.
- Khan PA (1992) Biochemical composition, minerals (calcium and iron) and chitin content of two portunid crabs *Scylla serrata* Forskal and *Portunus pelagicus* Linnaeus available in and around the coastal regions of Bangladesh. M.Sc. thesis, Institute of Marine Sciences, Chittagong University 112.
- Erkan N, Ozden O (2007) Proximate composition and mineral contents in aqua cultured sea bass (*Dicentrarchuslabrax*), sea bream (*Sparusaurata*) analyzed by ICP-MS. Food Chemistry 102: 721-725.
- Coombs TL (1974) The nature of zinc and copper complexes in the oyster *Ostrea edulis*. Marine biology 28: 1-10.
- Lowe DE, Moore MN (1979) The cytochemical distributions of zinc (ZnII) and iron (FeII) in the common mussel *Mytilus edulis* and their relationship with lysosomes. J Mar Biol Ass UK 59: 581.
- Rajagopal S, Ajmal KS, Srinivasan M, Shanmugam A (1998) Gastropods of Parangipettai coast. Centre of Advanced study in marine biology, Parangipettai, Annamalai University, India 36.
- Zhang H, Wang D, Wang Z (1994) Analysis of Nutrition composition of oyster *Ostrea denselamellosa*. Chinese Journal of Marine Drugs.

This article was originally published in a special issue, **Food Hydrocolloids** handled by Editor(s), Dr. Srinivas Janaswamy, Purdue University, USA

Citation: Saritha K, Mary D, Patterson J (2015) Nutritional Status of Green Mussel *Perna Viridis* at Tamil Nadu, Southwest Coast of India. J Nutr Food Sci S14: 003. doi:10.4172/2155-9600.S14-003