

Obstructive Sleep Apnea in Children: Why Screen?

Kulbir Singh Coraya*

Kulbir Singh Coraya, 101 South Whiting Street, Suite# 313, Alexandria, VA-22304, USA

Editorial

Obstructive sleep apnea (OSA) is a disorder of breathing and according to the American Academy of Sleep Medicine (AASM), OSA is characterized by repetitive episodes of complete or partial upper airway obstruction occurring during sleep. These repeated and short cessations in breathing are called apneas and may be caused by airway obstruction resulting from tonsils, uvula, redundant or fatty throat tissues and even blocked nasal airway etc.

OSA can affect anyone at any age, and in children, it is thought to be most common at ages 2-7 years [1] and if left untreated, may have severe consequences including neurocognitive impairment, behavioral problems, failure to thrive, growth failure, cardiovascular effects, including cor pulmonale, ventricular dysfunction, systemic hypertension etc. [2,3]. Risk factors for OSA are adenotonsillar hypertrophy, obesity, craniofacial anomalies, neuromuscular disorders, nasal abnormalities, increased waist and neck circumference, familial history, asthma, pre-term birth and so on. Some of the symptoms include habitual snoring, disturbed sleep, enuresis, awkward sleeping posture, daytime neurobehavioral problems and decreased academic performance [2-5].

OSA is a hidden health crisis, which is not only underdiagnosed but also undertreated [4]. While there are costs involved in the diagnosis and treatment of OSA, they are considerably lower than the costs associated with leaving the disorder untreated. Failure to treat OSA may increase the likelihood of disease-associated comorbidities [4]. The gold standard for the diagnosis of OSA is a sleep study called polysomnography and it is recommended in children and adolescents with signs and symptoms of OSA [3-6]. OSA in adults is characterized by episodes of airway obstruction for at least 10 seconds or more during sleep, resulting in breathing pauses. However in children, apneas of 3-4 seconds' duration are enough to cause oxygen desaturations [5]. Therefore, polysomnographic guidelines to diagnose OSA in children are not the same as those in adults [1-8].

Treatment options for OSA include continuous positive airway pressure (CPAP), surgery, oral appliances and lifestyle changes like body posture during sleep and weight loss [4].

Because of these serious and detrimental consequences of undetected and untreated OSA, it is of paramount importance to screen children for OSA. Tonsil and adenoid hypertrophy are generally acknowledged as the most common etiological factors of OSA in children, so dentists can play an important role in identifying adenotonsillar hypertrophy during examination [1-3]. Additionally, they can incorporate a questionnaire in the patient history forms to screen for the symptoms of OSA. For instance, Pediatric Sleep Questionnaire [6] has a good diagnostic accuracy and dentists may use it as a screening tool to identify the related symptoms of OSA. High-risk patients should be referred to sleep specialists [7] so that diagnosis and treatment of OSA may be done in a timely manner.

References

1. Capua M, Ahmadi N, Shapiro C (2009) Overview of obstructive sleep apnea in children: exploring the role of dentists in diagnosis and treatment. *J Can Dent Assoc* 75: 285-289.
2. Marcus CL, Chapman D, Ward SD, McColley SA, Herrerias CT, et al. (2002) Clinical practice guideline: Diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics* 109: 704-712.
3. Marcus CL, Brooks LJ, Draper KA (2012) Diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics* 130: 714-755.
4. Frost and Sullivan (2016) Hidden health crisis costing America billions. Underdiagnosing and undertreating obstructive sleep apnea draining healthcare system. *American Academy of Sleep Medicine*.
5. Chan J, Edman JC, Koltai PJ (2004) Obstructive sleep apnea in children. *Am Fam Physician* 69: 1147-54.
6. Chervin RD, Hedger K, Dillon JE, Pituch KJ (2000) Pediatric sleep questionnaire (PSQ): validity and reliability of scales for sleep-disordered breathing, snoring, sleepiness, and behavioral problems. *Sleep Med* 1: 21-32.
7. De Luca Canto G, Singh V, Major MP, Witmans M, El-Hakim H, et al. (2014) Diagnostic capability of questionnaires and clinical examinations to assess sleep-disordered breathing in children: a systematic review and meta-analysis. *J Am Dent Assoc* 145: 165-178.
8. Marcus CL, Omlin KJ, Basinki DJ, Bailey SL, Rachal AB, et al. (1992) Normal polysomnographic values for children and adolescents. *Am Rev Respir Dis* 146: 1235-1239.

*Corresponding author: Kulbir Singh Coraya, Kulbir Singh Coraya, 101 South Whiting Street, Suite313, Alexandria, VA-22304, USA, Tel: 703-823-2998; Fax: 703-823-8875; E-mail: ksg577@yahoo.com

Received May 06, 2017; Accepted May 31, 2017; Published June 07, 2017

Citation: Coraya KS (2017) Obstructive Sleep Apnea in Children: Why Screen? *Pediatr Dent Care* 2: e106. doi: [10.4172/2573-444X.1000e106](https://doi.org/10.4172/2573-444X.1000e106)

Copyright: © 2017 Coraya KS. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.