

## On Farm Pre Harvest Agronomic Management Practices of *Aspergillus* Infection on Groundnut in Abergelle, Tigray

Gebreselassie R<sup>1\*</sup>, Dereje A<sup>2</sup> and Solomon H<sup>3</sup>

<sup>1</sup>Mekelle Agricultural Research Center, crop core process, P.O.Box 258, Mekelle, Ethiopia

<sup>2</sup>Department of Crop and Horticultural Science, Mekelle University, P.O.Box 231, Mekelle, Ethiopia

<sup>3</sup>Department of Land Resource Management and Environmental protection, Mekelle University, P.O.Box 231, Mekelle, Ethiopia

### Abstract

*Aspergillus* infection and subsequent contamination of groundnut with Aflatoxin is a major limitation in groundnut production in the study area (Tanqua Abergel, Tigray). This study was executed to evaluate the effect of fertilizer (DAP and gypsum application), tied ridging and supplementary irrigation on *Aspergillus* infection of groundnut. The experiment was laid out in Randomized Complete Block Design with three replications in two sites. DAP as a source of P, and gypsum as source of Ca, were applied at planting and pod setting stages, respectively. While Tied ridging and supplementary irrigation were applied at early flowering and during cessation of rainfall, respectively. Data on incidence and severity levels of *Aspergillus* infection on sampled kernels were recorded. The analysis of variance indicated that the integrated agronomic management practices showed a significant reduction of *Aspergillus flavus* infection on groundnut at both experimental sites. The lowest (3%) *Aspergillus flavus* infection was recorded in management practices where Supplementary irrigation+tied ridging were practiced at Hadinet. The highest *Aspergillus flavus* infection (17.3%) was recorded in the control. In Lemlem experimental site the lowest *Aspergillus flavus* infection (4.3%) was recorded in gypsum+supplementary irrigation combination, while highest (19.3%) *Aspergillus flavus* infection was recorded in control. Application of integrated agronomic management practices did not showed a significant reduction of *Aspergillus niger* infection at both experimental sites.

**Keywords:** Agronomic practices; *Aspergillus* infection; groundnut

### Introduction

Groundnut is the sixth most important oilseed crop in the world. Groundnut kernels are consumed directly as a raw, roasted or boiled or oil extracted from the kernel is used as culinary oil. The crop is also used as animal feed and industrial raw material. Its multiple use and as a source of income make the crop important besides to its cash crop for domestic markets as well as for foreign trade in several developing and developed countries. *Aflatoxin* contamination has a significant role in groundnut production. *Aflatoxin* toxic fungal metabolites mainly produced by *Aspergillus flavus* and *A. parasiticus*. Contamination of groundnut with aflatoxin occurs under pre harvest, postharvest handling and storage conditions. The main factors leading to aflatoxin contamination include poor cultural practices; use of damaged and loose shelled kernels as seed and delayed harvesting after physiological maturity aggravates biological and physical effects of aflatoxins [1].

*Aspergilli* are group of fungi exhibiting immense ecological and metabolic diversity. These include notorious pathogen such as *Aspergillus fulvus*, which produces aflatoxin, one of the most potent, naturally occurring carcinogenic compounds known to man [1].

The development and contamination of the fungus in poorly managed fields is more serious. Timely planting, adequate fertility, good weeding and insect control, supplementary irrigation, suitable plant population and hybrid selection considerably reduce aflatoxin contamination [2].

The association of mycotoxins with consumption of contaminated groundnut and other food lead many importing countries, including the EU countries, to enact regulations establishing maximum levels for aflatoxins in groundnut and groundnut products [3].

The infection of *A.flavus* can be minimized to a level where it cannot cause significant damage through the application of different pre- harvest management practices. According to Doner [3] the risk to

infection and contamination of aflatoxin can be reduced substantially with proper irrigation supply particularly from flowering to pod setting stages, soil amendments; application of proper and rapid drying after harvest.

Groundnut by now is regarded as a cash crop and is the most important oil crop. The area coverage for groundnut production is constantly increasing. Farmers in Tanqua Abergelle are becoming interested in growing groundnut because of its high price as compared to other cereal and pulse crops. As a result farmers start sharing their land for groundnut production instead of planting the whole land for sorghum and other crop production.

Despite its importance as a food and cash crop in Ethiopia in general and study area in particular the quality of groundnut is poor which is mainly caused by *Aspergillus* infection at late growth stage of the crop. In addition, little work has been done on the extent of integrated agronomic management practices which improves quality of groundnut. Moreover, late onset and early cessations of rain fall in of Tigray in general and Abergalle in particular are common phenomena. And hence, studying different agronomic management practices that could alleviate such problems and improve quality of groundnut are important to groundnut production [4-6].

**\*Corresponding author:** Mekelle Agricultural Research Center, crop core process, P.O.Box 258, Mekelle Ethiopia, Tel: +251344408030; E-mail: [gebrhe62@yahoo.com](mailto:gebrhe62@yahoo.com)

**Received:** March 07, 2014; **Accepted** May 22, 2014; **Published** June 01, 2014

**Citation:** Gebreselassie R, Dereje A, Solomon H (2014) On Farm Pre Harvest Agronomic Management Practices of *Aspergillus* Infection on Groundnut in Abergelle, Tigray. J Plant Pathol Microb 5: 228. doi:10.4172/2157-7471.1000228

**Copyright:** © 2014 Gebreselassie R, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: Moisture content retention and *Aspergillus* infection in groundnut at 41-80 cm root depth at Hadinet, Ethiopia.




Figure 2: Moisture content retention and *Aspergillus* infection in groundnut at 41-80 cm root depth Lemlem.

And hence, the major objective of the study is thus to identify pre harvest agronomic management practices that reduces *Aspergillus* infection levels.

## Materials and Methods

### Description of the study area

The study was conducted during the 2011 cropping season in Tanqua Abergelle Wereda at *Tabia Lemlem* and *Hadinet* in the central administrative Zone of Tigray, northern Ethiopia on sandy soil. Tanqua Abergelle is located 120 kms far from Mekelle at an altitude of 1500 m.a.s.l, 130 14' 06"N Latitude & 38°58'50" E longitudes (Figure 1). Agro-ecologically it is characterized as hot warm sub-moist low land (SML-4b). The mean annual rainfall and temperature range of the wereda is 350–700mm & 24-41°C, respectively (Legesse 1999). The mean maximum and minimum temperature of the wereda during the 2011 growing season was 36.4°C and 19.5°C, respectively. While the total rainfall of *Hadinet* and *Lemlem* experimental sites during the 2011 growing season were 277 mm and 314 mm respectively (Figure 2).

### Treatments and experimental design

As planting material Sedi was used in the study area during the execution of the experiment. The experimental plot was arranged in

RCBD with three replications. Different farmers field were considered as a replication. The total plot size used was 9 m<sup>2</sup> (3 mX3 m), having 60 cm distance between rows and 20 cm between plants. The net harvestable area was 5.4 m<sup>2</sup>(1.8 mx3 m) leaving one outer most row as border.

DAP (100 kg/ha ), as a source of Phosphorous was applied at planting, application of gypsum 672 kg/ha was performed at early flowering stage; while Tied ridge and application of supplementary irrigation were done at the end of rainy season. The tied ridge had 20X30 cm depth of the soil. (Table 1)

### Crop water requirement and irrigation scheduling

Meteorological data (maximum and minimum temperature, relative humidity, sunshine hours, rainfall and wind speed) were collected from National Meteorology Agency Mekelle branch. Crop water requirement was determined using modified FAO Penman Monteith method [6]. Reference evapotranspiration (ET<sub>0</sub>) was calculated based on the climatic data using CWAT software. While the other parameters were calculated as:

Crop water requirement (Etc)=Eto\*Kc (crop coefficient at each growth stage of the crop)

Gross irrigation (GI)=Net irrigation (NI)/application efficiency (Ea), Ea was taken as 85%

NI=Etc-Pe (Pe=effective rain fall), since there was no rain fall during supplementary irrigation time NI=Etc. Irrigation was started at late September for both locations. Accordingly, irrigation water was applied for the specific treatments/plots every five days as supplementary.

### Soil sample collection and analysis

At planting, soil samples were randomly collected from each field to a depth of 0-40 cm with auger. Composite soil samples were prepared from the collected samples. The samples were oven dried at 105°C for 24 hours and ground to pass through a 2 mm sieve for physical and chemical analysis. Soil samples were analyzed for relevant soil parameters of bulk density, Ca, Mg and texture, permanent wilting point, and field capacity. Samples were also collected from each plot at harvest and the moisture content was determined by oven drying at 105°C till constant weight.

### Isolation and identification of aspergillus in laboratory

Groundnut kernel seed was examined in laboratory and the incidence and severity of *Aspergillus* infection was recorded based on incidence and severity percentage as described by [7]. Hundred kernels of groundnut samples from bulk were taken randomly from each experimental plot for lab analysis. Using the blotter plate method

Code Number	Treatments combination	Code Number	Treatments combination
1	C	9	SI+TD
2	GP	10	SI+F
3	SI	11	TD+F
4	TD	12	GP+SI+TD
5	F	13	GP+SI+F
6	GP+SI	14	GP+TD+F
7	GP+TD	15	SI+TD+F
8	GP+F	16	SI+TD+F+GP

C=Control, GP=gypsum, SI=supplementary irrigation, TD=tied ridge, F=fertilizer (DAP)

Table 1: list of treatments and treatments combinations.

Soil characteristics	Location	
	Hadinet	Lemelem
pH <sup>(1,2,5)</sup>	7.04	7.06
EC <sup>(1,2,5)</sup>	0.04	0.05
OC (%)	0.26	0.14
Olsen-P (mg/kg)	6.41	3.51
Total-N (%)	0.08	0.05
Exch Na (mg/kg)	31.33	14.33
Exch K (mg/kg)	21.33	10.67
Exch Ca+mg (mlieq/lit)	6.47	6.73
Exch Ca (mlieq/lit)	1.53	1.4
Sand (%)	94	94
Silt (%)	1	1
Clay (%)	5	5
texture	sand	sand

Table 2: Soil characteristics of experimental site.

Location	Root depth (cm)	Texture	Bulk density	FC	PWP	TAW (%) FC-PWP	TAW (volume), cm	TAW mm/m
Hadinet	0-40	sand	1.12	6.10	1.25	4.90	5.51	55.1
	41-80	sand	1.22	9.60	4.28	5.35	6.53	65.31
Lemelem	0-40	sand	1.11	6.12	1.25	4.87	5.12	54.22
	41-80	sand	1.23	9.64	4.28	5.36	6.57	65.75

TAW=total available water

Table 3: Field capacity (FC), permanent wilting point (PWP), and bulk density of experimental site.

[8] the seeds were soaked in distilled water for three minutes and surface sterilized by soaking in 1% sodium hypochlorite solution for one minute and rinsed three times with distilled water so as to protect from other external contamination. Then, the kernels were subjected to examination by placing them on moistened germination paper in the sterile plastic tray at an equal distance sealed with parafilm and were grown. The seeds were incubated for 5 days at 28°C with 12 hour day light and 12 hour dark. Then kernels infected by *Aspergillus* spp (*A.fluvs* and *A.nigr*) were counted based on the color of the fungus as described by [9].

Incidence and severity of each fungus types were determined according to [7]. Incidence as the percentage of number of *Aspergillus* infected samples by the fungus to the whole sample size as:

$$I = \frac{n}{N} \times 100, \text{ n represents number of } \textit{Aspergillus} \text{ infected sample and N represent the total sample size}$$

The severity (%) of each fungus was expressed as a percentage of infected grains per 100 plated seeds as:

$$S (\%) = \frac{n}{N} \times 100,$$

Where n is the number of infected seeds and N is the total planted seeds

### Statistical analysis

Analysis of variance (ANOVA) was carried out using Genstat software 13<sup>th</sup> version. Treatment means showing significant differences at 5% level of significance were compared using Duncan's Multiple Range Test (DMRT). As the value consisted of small whole numbers and range between 0 and 30% square root transformation method was employed before statistical analysis. Percentage data on fungal infection were transformed using square root method as described by [10-12].

## Result and Discussion

### Soil Characteristics

Results of soil analysis for soil sample taken before planting indicated that the proportion of sand, silt and clay was 94%, 1% and 5%, respectively. As a result the textural classes of the experimental soils are classed as sandy. The pH, values of soil in Hadinet and Lemlem was 7.04 and 7.06 (Table 2) respectively which indicated that the experimental soils are almost neutral and within the ideal pH range value for groundnut production. In addition the EC of Hadinet and Lemlem were 0.04 and 0.05 mmhos/cm (Table 2), respectively indicating that the sites were non saline.

Total available water for the experimental areas was 55.1 mm and 54.22 mm at 0-40 cm root depth at Hadinet and Lemlem, respectively (Table 3). While the total available water at a depth of 41- 80 cm was 65.31 and 65.75 mm/m at Hadinet and Lemlem experimental sites, respectively (Table 3). This result also agrees with the results of [6] where it is described that the values of total available soil water of sandy soil ranges between 50 mm–110 mm/m (Table 3).

### Soil water content at harvest

Soil moisture content was measured at harvest. Comparison among different management options in retaining water and their effect on improving yield of groundnut as well as impact in minimizing *Aspergillus* infection were determined. Analysis of variance on soil moisture determination at harvesting in Hadinet indicated that no significant difference ( $P < 5\%$ ) was observed among the different management options at a soil depth of (0–40 cm). But significant difference (at  $P < 5\%$ ) was observed at soil depth of 41-80 cm (Table

Treatments Mgt options)	Moisture content (%)			
	Hadinet		Lemelem	
	Soil depth (cm)		Soil depth (cm)	
	0-40	41-80	0-40	41-80
C	1.87	3.06a	3.93a	4.32a
F	3.57	5.63ab	7.30b	8.85bc
GP	4.16	6.05b	7.37b	8.35ab
GP+F	3.12	8.86cd	10.09cde	9.73bc
GP+SI+TD	5.15	9.04d	12.61ef	14.23de
GP+SI	5.07	7.35bcd	8.50bc	9.86bc
GP+SI+F	4.23	6.43bcd	9.62bcd	9.74bc
GP+TD	3.22	6.13b	8.02bc	8.94bc
GP+TD+F	3.78	6.19b	9.20bc	10.16bcd
SI	5.51	7.55bcd	11.93def	12.22bcde
SI+F	4.32	7.70bcd	10.15cde	11.58bcde
SI+TD	6.46	6.66bcd	13.68f	14.31de
SI+TD+F	5.60	8.92d	11.96def	13.10cde
SI+TD+F+GP	4.34	7.31bcd	12.91f	14.63e
TD	4.03	6.90bcd	7.04b	8.30ab
TD+F	3.39	5.29ab	9.07bc	10.87bcde
DMRTa 0.5	ns	2.718	2.690	4.267
CV %	35.9	23.9	16.8	24.2

<sup>a</sup>Any means having the common letter are not significantly different at the 5% level of significance

Table 4: Effect of Agronomic management options on moisture retention in different soil depth at harvesting.

No.	Trt	Incidence (yes/no), 1=yes ,0=no			
		A.flavus		A.niger	
		Hadinet	Lemelm	Hadinet	Lemelm
1	C	1	1	1	1
2	GP	1	1	1	1
3	SI	0	1	1	1
4	TD	1	1	1	1
5	F	1	1	1	1
6	GP+SI	1	1	1	1
7	GP+TD	1	1	1	1
8	GP+F	1	1	1	1
9	SI+TD	1	1	1	1
10	SI+F	1	1	1	1
11	TD+F	1	1	1	1
12	GP+SI+TD	1	1	1	1
13	GP+SI+F	1	1	1	1
14	GP+TD+F	1	1	1	1
15	SI+TD+F	1	1	1	1
16	SI+TD+F+GP	1	1	1	0
n/N*100		15/16*100	16/16*100	16/16*100	15/16*100
(%)		93%	100%	100%	93%

Table 5: Incidence of *A.flavus* and *A.niger*.

Treatments	Hadinet		Lemlem	
	A.flavus Infection (%)	A.niger Infection (%)	A.flavus Infection (%)	A.niger Infection (%)
C	17.3d	5.7	19.3f	6.3
F	9.7bcd	6.3	7.7e	5.0
GP	7.0abc	5.0	8.0e	4
GP+F	10.0cd	5.3	7.3de	2.0
GP+SI+TD	4.3abc	3.0	4.7ab	4.0
GP+SI	6.7abc	3.3	5.7abc	2.3
GP+SI+F	10.3cd	4.0	5.0ab	1.7
GP+TD	5.7abc	3.7	7.3de	2.3
GP+TD+F	10.0bcd	3.3	8.0e	3.7
SI	4.0ab	2.7	6.0bcd	2.3
SI+F	5.0abc	2.3	6.7cde	2.7
SI+TD	3.0a	1.3	4.3a	2.7
SI+TD+F	6.0abc	3.7	5.0ab	1.7
SI+TD+F+GP	10.3cd	6.3	5.3abc	2.3
TD	7.7abc	2.7	7.7e	4.0
TD+F	7.3abc	6.0	5.0ab	2.7
DMRTa 0.05	1.238	ns	0.2918	ns
CV %	27.5	33.5	6.5	32.5

Table 6: Effect of integrated Agronomic management practices on *A.flavus*, and *A.niger* infection (severity) of groundnut seed at Hadinet and Lemlem experimental site means having the common letter are not significantly different at the 5% level of significance.

4). The integrated management options showed an effect on moisture content percentage at both soil depth in Lemlem experimental site. The highest moisture content (9.04%) was recorded in Hadinet in (41–80 cm) soil depth, where gypsum+supplementary irrigation+tied ridging management options were practiced while the lowest moisture content was observed at the control (Table 4). In Lemlem experimental site the highest (13.68%) and the lowest (3.93 %) moisture content

were recorded in supplementary irrigation+tied ridging management options and control at soil depth of 0–40 cm, respectively this could be due to the better rain fall in amount and distribution than Hadinet. While in a soil depth of 41–80 cm 14.63 percent moisture content was recorded in gypsum+supplementary irrigation+tied ridging and fertilizer combination management options were applied (Table 4).

### Identification and incidence of Aspergillus species

Seeds collected from the experimental areas produced through application of integrated management options were examined for identification and incidence of fungus. Aspergilli were isolated from groundnut seed samples from both experimental sites. These include *A. flavus* and *A. niger* in both experimental sites. The incidence of *A. flavus* was 93% and 100% at Hadinet and Lemlem, respectively (Table 5). The incidence of *A. niger* was 100% and 93% at Hadinet and Lemlem, respectively (Table 5). *Aspergillus flavus*, however was, the dominant fungi (Table 5). This result was also in agreement with the findings of [11,12] which showed that *A. flavus* was the main species occurring in peanut seed.

### Effect of integrated agronomic management practices on infection of groundnut seed by aspergillus species

Groundnut seed infection by *Aspergillus* species can occur through various conditions. Aspergillus infection can occur under pre and post-harvest handling and storage conditions [13]. Study indicated that management practices minimize aflatoxin contamination caused by Aspergillus species. Breeding of resistant varieties, good agricultural production, processing, and handling and storage practices are some of the different management options. But it has been reported by [14] that little success was achieved in the development of groundnut varieties with resistance to aflatoxin contamination. In line with this an experiment was conducted using different management options. Results of the different integrated agronomic management options in minimization of Aspergillus species infection on groundnut are presented below (Table 6).

### Effect of soil fertility amendment practices on aspergillus species infection

Application of fertilizer (DAP) at planting as a means of reducing *A.flavus* and *A.niger* infection of groundnut showed a significant effect ( $P < 5\%$ ) over the control at Lemlem (Table 6). Significance difference on application of gypsum as a source of calcium over control was observed at both experimental site. From this result it can be suggested that the role of Calcium in reducing aspergillus infection is highly emphasized. Application of calcium in groundnut production alleviates problems like aborted and shriveled kernel seed of peanut and high percentage of ‘pops’ which are major sources for aspergillus infection. [15] reported that application of gypsum as a source of calcium is crucial to have good quality with totally sound matured groundnut seed.

### Effect of moisture retention practices and aspergillus species infection

Analysis of variance indicated that the integrated management options of soil moisture conservation practices applied at the early flowering stage resulted in a significant effect ( $P < 5\%$ ) in *Aspergillus flavus* infection at both experimental areas. However, the integrated management options did not affect the level of infection in *A. niger* at both locations. The lowest infection levels (3.0 and 4.3%) were recorded in SI+TD at Hadinet and Lemlem, respectively. But the highest severity levels (17.3 and 19.3%) were recorded in control at Hadinet and Lemlem experimental site, respectively (Table 6).






Figure 3: Map of Tanqua Abergela Woreda.




Figure 4: Mean monthly rain fall (mm) and maximum and minimum temperature of Abergelle Wereda during the 2011 growing season.

This result also agreed with findings of [16] who found that there was higher infection of peanut by *Aspergillus* and subsequent higher contamination where there was moisture stress during the last 20-40 days to growth period of the crop. In similar manner the role of irrigation on minimizing *Aspergillus* infection were highly emphasized. Finding on irrigation frequency and sowing date on groundnut reveals that infection of *Aspergillus* increased as the frequency declined from 7 day to non (rain fed) [17]. As indicated in (Table 6) no significance difference in *A.flavus* infection was observed between the two soil moisture amendments options (tied and supplementary irrigation) managements options implemented. In addition combination of all the soil moisture retention amendment practices (TD and SI) had no any significant difference as compared with the double and triple combination of soil fertility amendment practices (Table 6). Un like Lemlem application of fertilizer, gypsum+fertilizer, gypsum+supplementary+fertilizer, gypsum+tied+fertilizer and supplementary irrigation+fertilizer, gypsum+tied ridge combinations did not showed a significant over the control (Table 6). This might occur due to termite a problem which is serious problem in the areas well as other factors which result in moisture difference in the experimental plots.

## Moisture content at harvesting and aspergillus infection of groundnut

Statistical analysis indicated that moisture content and *Aspergillus* infection have inverse relations (Figure 3 and 4). Moisture content increased from 3% at control to 9.04% in supplementary irrigation+tied ridge+gypsum combinations at Hadinet (Figure 3). The infestation level of *Aspergillus* decreased from 17% at control to 3% in supplementary irrigation and tied ridge combination at Hadinet (Figure 3). In Lemlem the moisture content ranged from 4.32% in control to 14.63% in supplementary irrigation, tied ridge, Fertilizer (DAP) and gypsum combinations. As a result the infection level of *Aspergillus* decreased from 19.3% in control (without intervention of any agronomic practices) to 4.3% in supplementary irrigation and tied ridge combination (Figure 4).

## Conclusion

Application of gypsum as a source of calcium at early flowering stage significantly lowered *Aspergillus* infection compared over the control at Hadinet and Lemlem experimental sites.

Application of DAP fertilizer significantly lowered *Aspergillus* infection only at Lemlem experimental site indicating that moisture has a direct relationship with *Aspergillus* infection.

Application of soil moisture conservation practice (tied ridging) at early flowering stage of groundnut and supplementary irrigation at the cessation of rain fall significantly lowered *Aspergillus* infection compared to control at Hadinet and lemlem experimental site. At Hadinet experimental site no significant effect was observed between tied ridge and supplementary irrigation application as means of reducing *aspergillus* infection. This indicated that based on the availability of these practices, they can serve as a means of reducing *aspergillus* infection.

Though there was no consistency in reducing the infestation level of *Aspergillus flavus* over all agronomic combination practices, combination of supplementary irrigation, tied ridge, fertilizer and gypsum reduced the level of *Aspergillus flavus* infestation by 68% over the control.

Supplementary irrigation and tied ridge combination was effective and consistent (as compared to the other agronomic practices) agronomic practices that have significantly reduced level of *Aspergillus* infection over the control. Seeds samples were infested by *Aspergillus flavus* and *Aspergillus niger*. *Aspergillus flavus* was the dominantly occurred fungus in both locations.

## Recommendation

Soil moisture and fertility amendment practices reduce the level *Aspergillus flavus* infection level but to what level of aflatoxin these agronomic practices did reduce has to be considered for future research direction in verifying the present investigation across years and locations.

The experiment showed that in areas where water for supplementary irrigation is not available application of tied ridge become as an alternative option in reducing *Aspergillus flavus* infection.

## References

- UNIDO (2010) Capacity Building For Aflatoxin Management In Groundnuts In Malawi. Ministry Of Industry and Trade, and Ministry Of Agriculture and Food Security: Government Coordinating Agency.

2. Erick L (2009) Minimizing Aflatoxine in Corn. Mississippi state university Extension service.
3. Steven J, Luz BDR (2008) Agriculture and Rural Development. Barrier, Catalyst, or Distraction Standards, Competitiveness, and Africa's Groundnut.
4. Dorner JW (2008) Management and prevention of mycotoxins in peanuts. Food Addit Contam Part A Chem Anal Control Expo Risk Assess 25: 203-208.
5. Legesse Y (1999) Agroecology of zone of Tigray land use planning division of Tigray Buaraue of Agriculture and natural resource.
6. Allen RG, Pereira LS, Raes D, Smith M (1998) Crop evapotranspiration: guideline for computing crop water requirement. Irrigation and drainage paper number 56.FAO, Rome.
7. González HH1, Resnik SL, Boca RT, Marasas WF (1995) Mycoflora of Argentinian corn harvested in the main production area in 1990. Mycopathologia 130: 29-36.
8. ICRISAT (2005) Aspergillus flavus seed infection and aflatoxin estimation by ELISA and Aflatoxin management in groundnut.
9. Pitt JI, Hocking AD, Glenn DR (1983) An improved medium for the detection of Aspergillus flavus and A. parasiticus. J Appl Bacteriol 54: 109-114.
10. Gomez AK, Gomez AA (1984): Statistical Procedures for Agricultural Research. Second edition. John Wiley and Sons Inter-Science Publications. New York.
11. Idress HA, Laith TAA, Muftah AN, Ibrahim AAB, Maziah Z, et al. (2010) Screening of fungi associated with commercial grains and animal feeds. *World Applied Science Journal*, 9: 746- 756.
12. Sultan Y1, Magan N (2010) Mycotoxigenic fungi in peanuts from different geographic regions of Egypt. *Mycotoxin Res* 26: 133-140.
13. Mehan VK, Mayee CD, Jayanthi S, McDonald D (1991) Pre-harvest seed infection by Aspergillus flavus group of fungi and subsequent aflatoxin contamination in groundnuts in relation to soil types. *Plant Soil* 136: 239-248.
14. Waliyar F, Ba A, Hama H, Bonkougou S, Bosc JP (1994) Sources of resistance to Aspergillus flavus and aflatoxin contamination in groundnut genotypes in West Africa. *Plant Disease*, 78: 704-708.
15. Florence RJ (2011) Fertilization of Peanut (*Arachis hypogaea* L.) with Calcium: Influence of Source, Rate, and Leaching on Yield and Seed Quality. Msc. thesis, Auburn University, Auburn, Alabama.
16. Cole RJ, Sanders TJ, Dorner JW, Blankenship PD (1989) Environmental conditions required to induce pre-harvest concentration in groundnut. Summary of six years research.
17. Craufurd PW, Prasad PVV, Waliyar F, Taheri A (2006) Drought, pod yield, pre-harvest Aspergillus infection and aflatoxin contamination on peanut in Niger. *Field Crops Research* 98: 20-29.

**Citation:** Gebreselassie R, Dereje A, Solomon H (2014) On Farm Pre Harvest Agronomic Management Practices of Aspergillus Infection on Groundnut in Abergelle, Tigray. J Plant Pathol Microb 5: 228. doi:[10.4172/2157-7471.1000228](https://doi.org/10.4172/2157-7471.1000228)

### Submit your next manuscript and get advantages of OMICS Group submissions

#### Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

#### Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: [www.editorialmanager.com/](http://www.editorialmanager.com/)

