

## On the Generation of Efficient Methodologies for Cell-Based Toxicity Screening by Comparative Protocol Analysis and Optimization

Sepideh Abolpour Mofrad<sup>1,2</sup>, Katharina Kuenzel<sup>1,2</sup>, Oliver Friedrich<sup>1,2</sup> and Daniel F Gilbert<sup>1,2\*</sup>

<sup>1</sup>Institute of Medical Biotechnology, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany

<sup>2</sup>Erlangen Graduate School in Advanced Optical Technologies (SAOT), Friedrich-Alexander-Universität Erlangen-Nürnberg Erlangen, Germany

### Abstract

In cell biology in general and in high-throughput cell-based screening approaches in particular - e.g. for *in vitro*-based toxicity assessment - various methodologies or protocols are reported in the literature. In the case that a cell-based screening assay, for toxicity evaluation or for assessing another biological question, is to be established for the first time in a research lab, the researcher has to select a number of different protocols from the literature and to create an optimized protocol for the intended use. This is typically required, because optimized protocols, generated following comparative protocol analysis and optimization, are mostly rarely available. In another study, which we refer to as a showcase, we conducted a comparative analysis of three different protocols for neuronal NT2 cell differentiation, available in the literature. From this comparison, we generated an improved and optimized method, allowing for neuronal NT2 differentiation in monolayer cultures with high yield of NT2-N cells, allowing for systematic *in vitro*-based primary screening for developmental toxicants and neuro-toxicants at different stages of maturation. In this commentary, to prevent the same experiments from being repeatedly conducted in different labs around the world and at different times over and over again, we suggest generation of advanced and efficient methods by comparative protocol analysis and optimization, for application in a variety of research fields, related to cell and single cell biology.

**Keywords:** *In vitro* DNT testing; NT2 human pluripotent stem cells; Protocol optimization; High-throughput screening

### Showcase: *In vitro*-Based DNT Testing

Everyday products such as cosmetics, pharmaceuticals, textiles, detergents, plastic or pesticides, contain organic and inorganic chemical substances, that are potentially harmful to humans and the ecosystems. The number and volume of worldwide traded and registered chemicals has dramatically increased in recent years. Simultaneously, the incidence of neurological diseases, including mental retardation and autism, as well as developmental and learning disorders or hyperactivity disorder and attention deficit, has also increased [1-3]. The developing central nervous system (CNS) is particularly susceptible to damage by chemicals and therefore, assessment of the chemicals surrounding us on a daily basis for adverse effects on the maturing CNS - also referred to as developmental neurotoxicity (DNT) - is critical for human health and wealth [4-6]. Currently, as of January 2017, the CAS Registry (Chemical Abstracts Service), the world's largest database of chemical substances, lists more than 126 million unique organic and inorganic chemicals. Of these chemicals, only very few representatives have been evaluated for DNT in recent years [7,8]. The reason for this is possibly because existing guidelines for toxicity evaluation mostly involve animal experiments that are expensive, low in throughput, of poor predictive quality, often not reproducible and, because there is no legal obligation for alternative DNT standardized testing [9-12]. Hence, potentially harmful effects of many of the tens of thousands of chemicals contained in everyday products, still remain unknown.

### The Potential of Stem Cells for *In vitro*-Based DNT Testing

*In vitro*-based DNT testing using e.g. stem cells, such as induced pluripotent stem (iPS) cells or human embryonic stem (ES) cells, as well as human pluripotent teratocarcinoma NTERA-2 (NT-2) stem

cells with their almost unlimited proliferative potential, may provide a powerful alternative to animal experimentation [13]. However, the large variety of methodologies and protocols reported in the literature can be challenging when it comes to selecting the appropriate approach for maintenance and application of stem cells for *in vitro*-based toxicity assessment. For example, human pluripotent teratocarcinoma NTERA-2 (NT2) cells are increasingly considered as an appropriate model for such *in vitro*-based DNT screening approaches [14-19]. NT2 stem cells exhibit many aspects of human embryonic neural stem cells [20]. Upon treatment with retinoic acid (RA), NT2 cells differentiate into post-mitotic cells, showing properties of cells of the central nervous system, also referred to as NT2-N cells [21-26]. These cells are committed towards neurons, astrocytes, and oligodendrocytes, which are the three main cell lineages of the CNS, and have extensively been used for drug *in vitro* screening and for studying human neurogenesis, brain regeneration, and terminal differentiation [20].

### Spoiled for Choice - Which Protocol to Choose for a Cell-Based Application?

There are numerous approaches reported in the literature on neuronal differentiation of NT2 cells in aggregate or suspension culture, as well as in monolayer cultures [21,25-32]. These protocols


**\*Corresponding author:** Dr. Daniel Gilbert, Institute of Medical Biotechnology, Erlangen Graduate School in Advanced Optical Technologies (SAOT), Friedrich-Alexander-Universität Erlangen-Nürnberg Erlangen, Germany, Tel: +49 9131 850; E-mail: [daniel.gilbert@fau.de](mailto:daniel.gilbert@fau.de)

**Received** January 24, 2017; **Accepted** February 01, 2017; **Published** February 03, 2017

**Citation:** Mofrad SA, Kuenzel K, Friedrich O, Gilbert DF (2017) On the Generation of Efficient Methodologies for Cell-Based Toxicity Screening by Comparative Protocol Analysis and Optimization. Single Cell Biol 6: 157. doi:10.4172/2168-9431.1000157

**Copyright:** © 2017 Mofrad SA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Sepideh Abolpour Mofrad performed the present work in fulfillment of the requirements for obtaining the degree Dr. rer. biol. hum. at the Friedrich-Alexander-Universität of Erlangen-Nürnberg (FAU).



**Figure 1:** Application of the optimized protocol for neuronal differentiation of NT2-cells in monolayer cultures, generated from comparative protocol analysis. A). Transmission light images of non-differentiated NT2 cells. B). Differentiated NT2-N cells. Formation of interconnected clusters as indicated by white arrows indicates successful differentiation. Scale bar: 200  $\mu$ m.

mainly differ with respect to the presence or absence and exposure time of a differentiation stimulus (e.g. RA or 22R-hydroxycholesterol), the number and concentration, as well as the incubation time of mitotic inhibitors (cytosine arabinoside, fluorodeoxyuridine and uridine), or the frequency of media exchange, as well as procedures for purification of neuronal cells. The published studies report strongly varying differentiation results including expression of neuronal markers, morphological characteristics of differentiated cells and also efficiency - even from comparable approaches. For example, in the original study published by Andrews in 1984, a large proportion of >50% differentiated cells upon treatment with 10  $\mu$ M RA, is reported, whereas the studies by Pleasure et al. and Stewart et al. mention proportions of approx. 5% and 20% successfully differentiated cells, respectively [21,24,26].

### Protocol Comparison and Optimization

In the case that a cell-based screening assay for toxicity evaluation or for assessing another biological question is to be established for the first time in a research lab, the researcher needs to select a number of different protocols from the literature and apply and compare them, as well as subsequently generate an optimized protocol for the intended use, because data from comparative protocol analysis and optimization are typically rarely available. As this was the case with regard to methods describing differentiation of NT2 cells in monolayer culture at the time we aimed to work with NT2 cells in our lab, we conducted a comparative analysis of three different protocols for neuronal NT2 cell differentiation available in the literature. From our comparative data, we generated an improved and optimized method, allowing for neuronal NT2 differentiation in monolayer cultures with high yield of NT2-N cells, allowing for systematic *in vitro*-based primary screening for developmental toxicants and neuro-toxicants at different stages of maturation [33]. Figure 1 depicts images of non-differentiated NT2 as well as differentiated NT2-N cells obtained from our improved methodology.

### The Value of Comparative Protocol Analysis and Negative Results

Despite the fact that there are publications available in the literature on comparative protocol analysis and optimization, not only in the

context of stem cell differentiation, but also from other fields of cell biology, such literature is typically only rarely available. This is probably due to the fact that many of the required experiments often generate a large amount of negative results that are typically not published or publishable. However, data from comparative protocol analysis and optimization as well as negative results can be of high value to researchers aiming to establish a specific method in their laboratory, as the provided information may prevent the same experiments from being repeatedly conducted in different labs around the world and at different times over and over again. We thus, suggest generation of advanced and efficient methods by comparative protocol analysis and optimization, not only in the context of cell-based toxicity screening, but also for other research fields related to cell and single cell biology. This can eliminate the need for repetitive experiments, increase efficiency and contribute to furthering the applicability of cell-based screening methodologies.

### References

1. Colborn T (2004) Neurodevelopment and endocrine disruption. *Environ Health Perspect* 112: 944-949.
2. Herbert MR (2010) Contributions of the environment and environmentally vulnerable physiology to autism spectrum disorders. *Curr Opin Neurol* 23: 103-110.
3. Rauh VA, Garfinkel R, Perera FP, Andrews HF, Hoepner L, et al. (2006) Impact of prenatal chlorpyrifos exposure on neurodevelopment in the first 3 years of life among inner-city children. *Pediatrics* 118: e1845-1859.
4. Grandjean P, Landrigan PJ (2006) Developmental neurotoxicity of industrial chemicals. *Lancet* 368: 2167-2178.
5. Grandjean P, Landrigan PJ (2014) Neurobehavioural effects of developmental toxicity. *The Lancet Neurology* 13: 330-338.
6. Rice D, Barone S (2000) Critical periods of vulnerability for the developing nervous system: evidence from humans and animal models. *Environ Health Perspect* 108: 511-533.
7. Makris SL, Raffaele K, Allen S, Bowers WJ, Hasset U (2009) A retrospective performance assessment of the developmental neurotoxicity study in support of OECD test guideline 426. *Environmental health perspectives* 117: 17-25.
8. Middaugh LD, Dow-Edwards D, Li AA (2003) Neurobehavioral assessment: a survey of use and value in safety assessment studies. *Toxicol Sci* 76: 250-261.
9. Cooper RL, Lamb JC, Barlow SM (2006) A tiered approach to life stages testing for agricultural chemical safety assessment. *Crit Rev Toxicol* 36: 69-98.
10. Doe JE, Boobis AR, Blacker A, Dellarco V, Doerrer NG, et al. (2006) A tiered approach to systemic toxicity testing for agricultural chemical safety assessment. *Crit Rev Toxicol* 36: 37-68.

Sepideh Abolpour Mofrad performed the present work in fulfillment of the requirements for obtaining the degree Dr. rer. biol. hum. at the Friedrich-Alexander University of Erlangen-Nürnberg (FAU).

11. OECD (1997) Test Guideline 424. OECD Guideline for testing of Chemicals. Neurotoxicity Study in Rodents.
12. OECD (2007) Test Guideline 426. OECD Guideline for testing of Chemicals. Developmental Neurotoxicity Study.
13. NRC (2007) Toxicity testing in the 21st Century: A Vision and a Strategy. (1stedn), Washington DC, USA: National Academy Press.
14. Couillard-Despres S, Quehl E, Altendorfer K, Bentley K, Brady AM, et al. (2008) Human in vitro reporter model of neuronal development and early differentiation processes. *BMC neuroscience* 9: 31.
15. Hill EJ, Woehrling EK, Prince M, Coleman MD (2008) Differentiating human NT2/D1 neurospheres as a versatile in vitro 3D model system for developmental neurotoxicity testing. *Toxicology* 249: 243-250.
16. Hou Z, Zhang J, Schwartz MP, Stewart R, Page CD, et al. (2013) A human pluripotent stem cell platform for assessing developmental neural toxicity screening. *Stem Cell Res Ther* 4: S12.
17. Laurenza I, Pallocca G, Mennecozzi M, Scelfo B, Pamies D, et al. (2013) A human pluripotent carcinoma stem cell-based model for in vitro developmental neurotoxicity testing: Effects of methylmercury, lead and aluminum evaluated by gene expression studies. *Int J Dev Neurosci* 31: 679-691.
18. Menzner AK, Abolpour Mofrad S, Friedrich O, Gilbert DF (2015) Towards in vitro DT/DNT testing: Assaying chemical susceptibility in early differentiating NT2 cells. *Toxicology* 338: 69-76.
19. Pallocca G, Fabbri M, Sacco MG, Gribaldo L, Pamies D, et al. (2013) miRNA expression profiling in a human stem cell-based model as a tool for developmental neurotoxicity testing. *Cell Biol Toxicol* 29: 239-257.
20. Haile Y, Fu W, Shi B, Westaway D, Baker G, et al. (2014) Characterization of the NT2-derived neuronal and astrocytic cell lines as alternative in vitro models for primary human neurons and astrocytes. *J Neurosci Res* 92: 1187-1198.
21. Andrews PW (1984) Retinoic acid induces neuronal differentiation of a cloned human embryonal carcinoma cell line in vitro. *Dev Biol* 103: 285-293.
22. Coyne L, Shan M, Przyborski SA, Hirakawa R, Halliwell RF (2011) Neuropharmacological properties of neurons derived from human stem cells. *Neurochem Int* 59:404-412.
23. Kuenzel K, Friedrich O, Gilbert DF (2016) A recombinant human pluripotent stem cell line stably expressing halide-sensitive YFP-I152L for GABAAR and GlyR-targeted high-throughput drug screening and toxicity testing. *Front Mol Neurosci* 9: 51.
24. Pleasure SJ, Lee VM (1993) NTera 2 cells: A human cell line which displays characteristics expected of a human committed neuronal progenitor cell. *J Neurosci Res* 35: 585-602.
25. Pleasure SJ, Page C, Lee VM (1992) Pure, postmitotic, polarized human neurons derived from NTera 2 cells provide a system for expressing exogenous proteins in terminally differentiated neurons. *J Neurosci* 12: 1802-1815.
26. Stewart R, Coyne L, Lako M, Halliwell RF, Przyborski SA (2004) Human embryonal carcinoma stem cells expressing green fluorescent protein form functioning neurons in vitro: A research tool for co-culture studies. *Stem Cells Dev* 13: 646-657.
27. Hsu TC, Liu KK, Chang HC, Hwang E, Chao JI (2014) Labeling of neuronal differentiation and neuron cells with biocompatible fluorescent nanodiamonds. *Scientific reports* 4: 5004.
28. Megiorni F, Mora B, Indovina P, Mazzilli MC (2005) Expression of neuronal markers during NTera2/cloneD1 differentiation by cell aggregation method. *Neurosci Lett* 373: 105-109.
29. Paquet-Durand F, Tan S, Bicker G (2003) Turning teratocarcinoma cells into neurons: rapid differentiation of NT-2 cells in floating spheres. *Brain Res Dev Brain Res* 142: 161-167.
30. Podrygajlo G, Tegenge MA, Gierse A (2009) Cellular phenotypes of human model neurons (NT2) after differentiation in aggregate culture. *Cell Tissue Res* 336: 439-452.
31. Saporta S, Spencer E, Shamekh R (2014) Stable differentiation of NT2 cells. Google Patents.
32. Yao ZX, Han Z, Xu J, Greeson J, Lecanu L, et al. (2007) 22R-Hydroxycholesterol induces differentiation of human NT2 precursor (NTera2/D1 teratocarcinoma) cells. *Neuroscience* 148: 441-453.
33. Abolpour Mofrad S, Kuenzel K, Friedrich O, Gilbert DF (2016) Optimizing neuronal differentiation of human pluripotent NT2 stem cells in monolayer cultures. *Dev Growth Differ* 58: 664-676.