

# Orthogonal Left Derivations of Semi-Prime Rings

Ali Al-Hachami KH\*

Department of Mathematics, Wasit University, Al Kut, Wasit, Iraq

## Abstract

In this paper we show a few outcomes concerning two remaining deductions on a semi-prime ring are displayed. These outcomes are identified with an outcome which is motivated by Posner's hypothesis. This outcome affirms that if  $R$  is a 2-torsion free semi-prime ring,  $\delta$  and  $g$  are non-zero remaining inductions of  $R$  with the end goal that  $g$  is a surjective on  $R$ , and  $g(y)\delta(x)=g(x)\delta(y)$  for all  $x,y \in R$ . At that point  $\delta$   $g$  can't be a non-zero left derivation. A thought of orthogonal left derivations emerges here.

**Keywords:** Left derivation; Orthogonal left derivations; Prime ring; Semi-prime ring

## Introduction

All through  $R$  will speak to a cooperative ring.  $R$  is said to be 2-torsion free if  $2x=0, x \in R$  implies  $x=0$  [1]. Review that  $R$  is prime if  $xRy=0$  implies  $x=0$  or  $y=0$ , and  $R$  is semi-prime if  $xRx=0$  suggests  $x=0$ . Ref. [2], characterized the accompanying thought. An added substance mapping  $\delta:R \rightarrow R$  is known as a left inference if  $\delta(xy)=x\delta(y)+y\delta(x)$  holds for all  $x,y \in R$ . Different properties of left deductions can be found in refs. [3-8].

Two additive mapping  $\delta, g:R \rightarrow R$  is said to be orthogonal if:

$$(x)Rg(y)=0=g(y)R\delta(x) \text{ for all } x,y \in R.$$

Brešar and Vukman [9] presented the idea of orthogonality for two inductions  $\delta$  and  $g$  on a semi-prime ring, and they introduced a few important and adequate conditions for  $\delta$  and  $g$  to be orthogonal. In ref. [10] the creators presented orthogonal summed up inferences on a semi-prime ring and they introduced a few outcomes concerning two summed up determinations on a semi-prime ring. Their outcomes are a speculation of after effects of Brešar and Vukman in ref. [9]. What's more [11], in the creators presented orthogonal  $(\sigma, \tau)$ -determinations and orthogonal summed up  $(, \tau)$ -deductions. Their outcomes dreamy a few aftereffects of Brešar and Vukman [9]. In this paper, our point is to give similar consequences of Brešar and Vukman to orthogonal left derivations [9].

For a generalized semi-prime ring  $R$  and a perfect  $U$  of  $R$ , it is outstanding that the left and right annihilators of  $U$  in  $R$  agree [12]. We indicate the annihilator of  $U$  by  $\text{Ann}(U)$ . Take note of that  $U \cap \text{Ann}(U)=0$  and  $U \oplus \text{Ann}(U)$  is a fundamental perfect of  $R$  [12].

## Materials and Methods

In the accompanying, we give the documentation of orthogonal left derivations.

### Definition 2.1

Left derivations  $\delta$  and  $g$  are called orthogonal if,

$$(x)Rg(y)=0=g(y)R\delta(x), \text{ for all } x,y \in R. \quad (1)$$

Clearly a non-zero remaining deduction cannot be orthogonal on itself.

Give us a chance to consider a straightforward case of the non-zero orthogonal left derivations.

**Example:** Give  $S$  a chance to be a prime ring and set  $R=S \oplus S$ . At

that point  $R$  is a semi-prime ring. Give  $\delta$  and  $g$  a chance to be two non-zero remaining deductions of  $S$ . At that point the maps  $\delta_1$  and  $g_1$  from  $R$  to  $R$ , which are characterized by:

$\delta_1((x,y))=(\delta(x),0)$  and  $g_1((x,y))=(0,g(y))$ , for all  $x,y \in S$ , are non-zero left derivations of  $R$ .

Then  $\delta_1$  and  $g_1$  are orthogonal.

Presently, to get the primary outcomes, we require the accompanying lemmas:

### Lemma 2.2 (Lemma 1 [9])

Give  $R$  a chance to be a 2-torsion free semi-prime ring and  $a,b$  the components of  $R$ . At that point the accompanying conditions are proportional:

- (i)  $axb=0$ , for all  $x \in R$ .
- (ii)  $bxa=0$ , for all  $x \in R$ .
- (iii)  $axb+bxa=0$ , for all  $x \in R$ .

On the off chance that one of these conditions is satisfied then abdominal muscle  $ab=ba=0$

### Lemma 2.3 (Lemma 2.2 [9])

Give  $R$  a chance to be a semi-prime ring. What's more, assume that added substance mappings  $f$  and  $h$  of  $R$  into itself fulfill satisfy  $f(x)Rh(x)=0$ , for all  $x \in R$ . Then  $f(x)Rh(y)=0$ , for all  $x,y \in R$ .

## Results and Discussion

In Theorem 3.1, we will demonstrate that If  $\delta$  and  $g$  are orthogonal left derivations of a 2-torsion free semi-prime ring  $R$ , then there exists a fundamental perfect  $E$  of  $R$ , with the end goal that the confinements of  $\delta$  and  $g$  to  $E$  are fitting direct wholes.

**\*Corresponding author:** Ali Al-Hachami KH, Department of Mathematics, Wasit university, Al Kut, Wasit, Iraq, Tel: 009647832315025; E-mail: [alhachamia@uowasit.edu.iq](mailto:alhachamia@uowasit.edu.iq)

Received January 21, 2017; Accepted June 27, 2017; Published July 02, 2017

**Citation:** Ali Al-Hachami KH (2017) Orthogonal Left Derivations of Semi-Prime Rings. J Generalized Lie Theory Appl 11: 270. doi: [10.4172/1736-4337.1000270](https://doi.org/10.4172/1736-4337.1000270)

**Copyright:** © 2017 Ali Al-Hachami KH. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Theorem 3.1**

Give R a chance to be a 2-torsion free semi-prime ring. Give  $\delta$  and  $g$  a chance to be left derivations of R. At that point the accompanying conditions are equal:

- (i)  $\delta$  and  $g$  are orthogonal.
- (ii) There exist standards  $E_1$  and  $E_2$  of R with the end goal that  $E_1 \cap E_2 = 0$  and  $E = E_1 \oplus E_2$  is a basic perfect of R.
  - (a) maps R into  $E_1$  and  $g$  maps R into  $E_2$ .
  - (b) The restriction of  $\delta$  to  $E = E_1 \oplus E_2$  is a direct sum  $\delta_1 \oplus \delta_2$ , where  $\delta_1: E_1 \rightarrow E_1$  is a left derivation of  $E_1$  and  $\delta_2: E_2 \rightarrow E_2$  is zero. If  $\delta_1 = 0$  then  $\delta = 0$ .
  - (c) The restriction of  $g$  to  $E = E_1 \oplus E_2$  is a direct sum  $g_1 \oplus g_2$ , where  $g_1: E_1 \rightarrow E_1$  is zero and  $g_2: E_2 \rightarrow E_2$  is a left derivation of  $E_2$ . If  $g_2 = 0$  then  $g = 0$ .

In Theorem 3.2, we give a few vital and adequate conditions for the orthogonality of two left derivations.

**Theorem 3.2**

Give R a chance to be a 2-torsion free semi-prime ring. Let  $\delta$  and  $g$  be left derivations of R such that  $g$  is a surjective on R, and  $g(y)\delta(x) = g(x)\delta(y)$ , for all  $x, y \in R$ . Then  $\delta$  and  $g$  are orthogonal if and only if one of the following conditions holds:

- (i)  $g(x)\delta(x) = 0$ , for all  $x \in R$ .
- (ii)  $\delta(x)g(x) = 0$ , for all  $x \in R$ .
- (iii)  $g(x)\delta(x) + \delta(x)g(x) = 0$ , for all  $x \in R$ .
- (iv)  $\delta g = 0$ .
- (v)  $g\delta = 0$ .
- (vi)  $\delta g + g\delta = 0$ .
- (vii)  $\delta g$  is a left derivation.
- (viii)  $g\delta$  is a left derivation.
- (ix) There exist  $a, b$  in R such that  $(\delta g)(x) = xa + xb$ , for all  $x \in R$ .

For the evidence of the Theorem 3.1 and Theorem 3.2, we require the accompanying lemma:

**Lemma 3.3**

Give R a chance to be a 2-torsion free semi-prime ring. Give  $\delta$  and  $g$  a chance to be left derivations of R. In the event that  $\delta$  and  $g$  are orthogonal then the accompanying relations hold.

- (i)  $g(x)\delta(x) = 0$ , for all  $x \in R$ .
- (ii)  $\delta(x)g(x) = 0$ , for all  $x \in R$ .
- (iii)  $g(x)\delta(x) + \delta(x)g(x) = 0$ , for all  $x \in R$ .
- (iv)  $\delta g = 0$ .
- (v)  $g\delta = 0$ .
- (vi)  $\delta g + g\delta = 0$ .

**Proof:** (i) By the hypothesis we have  $\delta(x)Rg(x) = 0$ , for all  $x \in R$ . By Lemma 2.2, we get  $g(x)\delta(x) = 0$ , for all  $x \in R$ .

- (ii) By the hypothesis we have  $\delta(x)Rg(x) = 0$ , for all  $x \in R$ .

By Lemma 2.2, we get  $\delta(x)g(x) = 0$ , for all  $x \in R$ .

(iii) By the hypothesis we have  $\delta(x)Rg(x) = 0$ , for all  $x \in R$ .

By Lemma 2.2, we get  $\delta(x)g(x) = g(x)\delta(x) = 0$ , for all  $x \in R$ .

Thus  $g(x)\delta(x) + \delta(x)g(x) = 0$ , for all  $x \in R$ .

(iv) We have  $\delta(x)yg(z) = 0$ , for all  $x, y, z \in R$ . Hence,

$$\begin{aligned} 0 &= \delta(\delta(x)yg(z)) \\ &= \delta(x)\delta(yg(z)) + yg(z)\delta^2(x) \\ &= \delta(x)y(\delta g)(z) + \delta(x)g(z)\delta(y) + yg(z)\delta^2(x) \end{aligned}$$

The second two summands are zero since  $\delta$  and  $g$  are orthogonal. Therefore, this relation reduces to  $\delta(x)y(\delta g)(z) = 0$ , where  $x, y, z$  are arbitrary elements in R. But then also  $(\delta g)(z)R(\delta g)(z) = 0$ , for all  $z \in R$ . Since R is semi-prime, we get  $(\delta g)(z) = 0$ .

The second two summands are zero since  $\delta$  and  $g$  are orthogonal. In this manner, this connection decreases to  $\delta(x)y(\delta)(z) = 0$ , where  $x, y$  and  $z$  are discretionary components in R. Be that as it may, then likewise  $(\delta g)(z)R(\delta g)(z) = 0$ , for all  $z \in R$ . Since R is semi-prime, we get  $(\delta g)(z) = 0$ .

(v) By a similar way in (iv) we get the result.

(vi) From (iv) and (v), Lemma 2.4, we have  $\delta g + g\delta = 0$ .

We need the following lemma to proof Theorem 3.2.

**Lemma 3.4**

Let R be a 2-torsion free semi-prime ring. Let  $\delta$  and  $g$  be left derivations of R such that  $g$  is a surjective on R, and  $g(y)\delta(x) = g(x)\delta(y)$ , for all  $x, y \in R$ . Then  $\delta$  and  $g$  are orthogonal if and only if  $g(y)\delta(x) + g(x)\delta(y) = 0$ , for all  $x, y \in R$ .

**Proof:** Suppose that  $g(y)\delta(x) + g(x)\delta(y) = 0$ , for all  $x, y \in R$ . By the assumption, we have  $g(y)\delta(x) = 0$ , for all  $x, y \in R$ . Since R is 2-torsion free, we have  $g(y)\delta(x) = 0$ , for all  $x, y \in R$ . Take  $x = g(z)x$  in the above relation, where  $z$  in R, we get,

$$\begin{aligned} 0 &= g(y)\delta(g(z)x) \\ &= g(y)g(z)\delta(x) + g(y)x\delta(g(z)) \\ &= g(y)x\delta(g(z)), \text{ for all } x, y, z \in R. \end{aligned}$$

Since  $g$  is surjective, we get  $g(y)x\delta(z) = 0$ , for all  $x, y, z \in R$ .

Then  $g(y)R\delta(z) = 0$ , for all  $y, z \in R$ . Using Lemma 2.2, we see that  $\delta$  and  $g$  are orthogonal.

Conversely, if  $\delta$  and  $g$  are orthogonal, we have,

$$\delta(x)Rg(y) = 0, \text{ for all } x, y \in R.$$

By Lemma 2.2, we get  $g(y)\delta(x) = g(x)\delta(y) = 0$ , for all  $x, y \in R$ .

Thus  $g(y)\delta(x) + g(x)\delta(y) = 0$ , for all  $x, y \in R$ .

Let  $\delta$  and  $g$  be left derivations of any ring R. By a direct computation, we verify the following identities:

$$(\delta g)(xy) = x(\delta g)(y) + g(y)\delta(x) + g(x)\delta(y) + y(\delta g)(x) \tag{2}$$

$$(g\delta)(xy) = x(g\delta)(y) + \delta(y)g(x) + \delta(x)g(y) + y(g\delta)(x) \tag{3}$$

We now have enough information's to prove Theorem 3.1.

**Proof of Theorem 3.1:** (i)  $\Rightarrow$  (ii). Let  $E_1$  be an ideal of R generated by all  $\delta(x)$ ,  $x \in R$ , and let  $E_2$  be  $\text{Ann}(E_1)$ , the annihilator of  $E_1$ . From eqn. (1) we see that  $g(x)$ ,  $x \in E_2$ , for all  $x \in R$ . Whenever  $E_1$  is an ideal in a semi-prime ring we have  $E_1 \cap E_2 = 0$  and  $E = E_1 \oplus E_2$  is an essential ideal. Thus (a) and (b) are proved.

Our next goal is to show that  $\delta$  is zero on  $E_2$ . Take  $e_2 \in E_2$ . Then  $e_1 e_2 = 0$ , for all  $e_1 \in E_1$ . Hence  $0 = \delta(e_1 e_2) = e_1 \delta(e_2) + e_2 \delta(e_1)$ .

It is obvious from the definition of  $E$  that  $\delta$  leaves  $E_1$  invariant, hence  $e_2 \delta(e_1) = 0$ . Then the relation above reduces to  $e_1 \delta(e_2) = 0$ . Since in a semi-prime ring the left and right and two-sided annihilators of an ideal coincide, then we have  $\delta(e_2) \in \text{Ann}(E_1) = E_2$ . But on the other hand  $\delta(e_2)$  belongs to the set of generating elements of  $E_1$ . Thus  $\delta(e_2) \in E_1 \cap E_2 = 0$ , which means that  $\delta$  is zero on  $E_2$ .

As we have mentioned above  $\delta$  leaves  $E_1$  invariant. Therefore we may define a mapping  $\delta_1: E_1 \rightarrow E_1$  as a restriction of  $\delta$  to  $E_1$ . Suppose that  $\delta_1 \neq 0$ . Then  $\delta$  is zero on  $E = E_1 \oplus E_2$ . Take  $e \in E$  and  $x \in R$ . We have  $\delta(ex) = e\delta(x) + x\delta(e)$ . But  $\delta(ex) = \delta(e) = 0$  since  $ex, e \in R$ .

Consequently  $e\delta(x) = 0$ , for all  $x \in R$ . Thus  $\delta(x) \in \text{Ann}(E)$ . But ideal  $E$  is essential and therefore  $\text{Ann}(E) = 0$ . Hence  $\delta(x) = 0$ , for all  $x \in R$ . Then (c) is thereby proved.

It remains to prove (d). First we show that  $g$  is zero on  $E_1$ . Take  $x, y, z \in R$  and set  $e_1 = x \delta(y) z$ . Then  $g(e_1) = g(x\delta(y)z) = xg(\delta(y)z) + \delta(y)zg(x) = x\delta(y)g(z) + xz(g\delta)(y) + \delta(y)zg(x)$ .

Since  $\delta$  and  $g$  are orthogonal we have  $\delta(y)g(z) = 0$ ,  $\delta(y)zg(x) = 0$  and  $g\delta = 0$  by Lemma 3.3. Hence  $g(e_1) = 0$ . In a similar fashion we see that  $g(x\delta(y)) = 0$ ,  $g(\delta(y)z) = 0$  and  $g(\delta(y)) = 0$  by Lemma 3.3. Then  $g$  is zero on  $E_1$ . Recall that  $g$  maps  $R$  into  $E_2$ . In particular, it leaves  $E_2$  invariant. Thus, we may define  $g_2: E_2 \rightarrow E_2$  as a restriction of  $g$  to  $E_2$ . The proof that  $g_2 = 0$  implies  $g = 0$  is the same as the proof that  $\delta_1 = 0$  implies  $\delta = 0$ .

(ii)  $\Rightarrow$  (i). Clear.

**Proof of Theorem 3.2:** " $\delta$  and  $g$  are orthogonal"  $\Rightarrow$  (i), (ii), (iii), (iv), (v) and (vi) are proved by Lemma 3.

(i)  $\Rightarrow$  " $\delta$  and  $g$  are orthogonal". A linearization of  $g(x)\delta(x) = 0$  gives,  $g(x)\delta(y) + g(y)\delta(x) = 0$ , for all  $x, y \in R$ .

Hence  $\delta$  and  $g$  are orthogonal by Lemma 4.

(ii)  $\Rightarrow$  " $\delta$  and  $g$  are orthogonal". A linearization of  $\delta(x)g(x) = 0$  gives,  $(x)g(y) + \delta(y)g(x) = 0$ , for all  $x, y \in R$ .

Left multiplication by  $g(y)$  in the above relation gives,

$$g(y)\delta(x)g(y) + g(y)\delta(y)g(x) = 0, \text{ for all } x, y \in R.$$

By the assumption, we get,

$$g(x)\delta(y)g(y) + g(y)\delta(y)g(x) = 0, \text{ for all } x, y \in R.$$

Hence,

$$g(y)\delta(y)g(x) = 0, \text{ for all } x, y \in R.$$

Since  $g$  is surjective, we get,

$$g(y)\delta(y)x = 0,$$

where  $x, y$  are arbitrary elements in  $R$ .

Since  $R$  is semi-prime, we get,

$$g(y)\delta(y) = 0, \text{ for all } y \in R.$$

Therefore, by (i), Theorem 3.2, we get the result.

(iii)  $\Rightarrow$  " $\delta$  and  $g$  are orthogonal". Suppose that  $g(x)\delta(x) + \delta(x)g(x) = 0$ , for all  $x \in R$ . Then,

$$g(x)\delta(x) = -\delta(x)g(x),$$

for all  $x \in R(\cdot)$ ,

A linearization of  $g(x)\delta(x) + \delta(x)g(x) = 0$  gives,

$$g(x)\delta(y) + g(y)\delta(x) + \delta(x)g(y) + \delta(y)g(x) = 0, \text{ for all } x, y \in R.$$

By the assumption, we have  $2g(y)\delta(x) + \delta(x)g(y) + \delta(y)g(x) = 0$ , for all  $x, y \in R$ . Left multiplication by  $g(y)$  in the above relation, we get  $2g(y)g(x)\delta(x) + g(y)\delta(x)g(y) + g(y)\delta(y)g(x) = 0$ , for all  $x, y \in R$ . By (.) and the assumption, we obtain  $2g(y)g(x)\delta(y) + g(x)\delta(y)g(y) - \delta(y)g(y)g(x) = 0$ , for all  $x, y \in R$ .

Hence  $2g(y)g(x)\delta(y) + [g(x), \delta(y)g(y)] = 0$ . Take  $g(x) = \delta(y)g(y)$  in the above relation, we get  $2g(y)g(x)\delta(y) = 0$ , for all  $x, y \in R$ . Since  $R$  is 2-torsion free and  $g$  is surjective, we have  $g(y)x\delta(y) = 0$ , for all  $x, y \in R$ . Then  $g(y)R\delta(y) = 0$ , for all  $y \in R$ .

By Lemma 2.3, we then have  $g(y)R\delta(z) = 0$ , for all  $y, z \in R$ .

Using Lemma 2.2, we see that  $\delta$  and  $g$  are orthogonal.

(iv)  $\Rightarrow$  " $\delta$  and  $g$  are orthogonal". Suppose that  $\delta g = 0$ . According to eqn. (2), we have,

$$g(y)\delta(x) + g(x)\delta(y) = 0, \text{ for all } x, y \in R.$$

Hence, we get  $\delta$  and  $g$  are orthogonal by Lemma 3.4.

(v)  $\Rightarrow$  " $\delta$  and  $g$  are orthogonal". Suppose that  $g\delta = 0$ . According to eqn. (3), we have,

$$\delta(y)g(x) + \delta(x)g(y) = 0, \text{ for all } x, y \in R.$$

Take  $y = x$  in the above relation, we get,

$$2\delta(x)g(x) = 0, \text{ for all } x \in R.$$

Since  $R$  is 2-torsion free, we have,

$$\delta(x)g(x) = 0, \text{ for all } x \in R.$$

Therefore, by (ii), Theorem 3.2, we get the result.

(vi)  $\Rightarrow$  " $\delta$  and  $g$  are orthogonal". If  $\delta$  and  $g$  are any left derivations then we have by eqns. (2) and (3) that,

$$(\delta g + g\delta)(xy) = x(\delta g + g\delta)(y) + g(y)\delta(x) + g(x)\delta(y) + \delta(y)g(x) + \delta(x)g(y) + y(\delta g + g\delta)(x)$$

Thus, if  $\delta g + g\delta = 0$ , then the above relation reduces to,

$$g(y)\delta(x) + g(x)\delta(y) + \delta(y)g(x) + \delta(x)g(y) = 0, \text{ for all } x, y \in R.$$

Take  $x = y$  in the above relation, then we have  $2(g(x)\delta(x) + \delta(x)g(x)) = 0$ , for all  $x \in R$ . Since  $R$  is 2-torsion free, we have  $g(x)\delta(x) + \delta(x)g(x) = 0$ , for all  $x \in R$ . Therefore, by (iii), Theorem 3.2, we get the result.

(iv)  $\Rightarrow$  (vii). Clear.

(vii)  $\Rightarrow$  " $\delta$  and  $g$  are orthogonal". Since  $\delta g$  is a left derivation. We have,

$$(\delta g)(xy) = x(\delta g)(y) + y(\delta g)(x), \text{ for all } x, y \in R. \text{ Comparing this express with eqn. (2), we obtain } g(y)\delta(x) + g(x)\delta(y) = 0, \text{ for all } x, y \in R.$$

Now apply Lemma 3.4.

(v)  $\Rightarrow$  (viii). Clear.

(viii)  $\Rightarrow$  " $\delta$  and  $g$  are orthogonal". Since  $g\delta$  is a left derivation. We have,

$$(g\delta)(xy) = x(g\delta)(y) + y(g\delta)(x), \text{ for all } x, y \in R. \text{ Comparing this express with eqn. (3), we obtain } \delta(y)g(x) + \delta(x)g(y) = 0, \text{ for all } x, y \in R.$$

Let  $y=x$  in the above relation, we get  $2\delta(x)g(x)=0$ , for all  $x \in R$ .

Since  $R$  is a 2-torsion free, we have  $\delta(x)g(x)=0$ , for all  $x \in R$ .

Therefore, by (ii), Theorem 3.2, we get the result.

(iv)  $\Rightarrow$  (ix). Clear.

(ix)  $\Rightarrow$  " $\delta$  and  $g$  are orthogonal". For every  $x, y \in R$  we have  $(\delta g)(x y)=xya+xyb$ .

That is,  $x(\delta g)(y)+g(y)\delta(x)+g(x)\delta(y)+y(\delta g)(x)=xya+xyb$ .

Using  $(\delta g)(x)=xa+xb$  and  $(\delta g)(y)=ya+yb$ ,

we get  $2g(x)\delta(y)+yx(a+b)=0$ . Replacing  $x$  by  $y$ ,  $x$  in the above relation yields that  $y\{2g(x)\delta(y)+yx(a+b)\}+2xg(y)\delta(y)=0$

Then we have  $2xg(y)\delta(y)=0$ , for all  $x, y \in R$ .

Since  $R$  is 2-torsion free, we have  $x g(y)\delta(y)=0$ , where  $x, y$  are arbitrary elements in  $R$ . Since  $R$  is semi-prime, we get  $g(y)\delta(y)=0$ , for all  $y \in R$ .

Therefore, by (i), Theorem 3.2, we get the result.

A notable consequence of Posner [1] states that, if  $R$  is a prime ring of trademark not 2,  $\delta$  and  $g$  are non-zero inductions of  $R$ , then  $\delta g$  cannot be a derivation. The outcome which is enlivened by a hypothesis of E. Posner, states that, if  $R$  is a 2-torsion free semi-prime ring,  $\delta$  and  $g$  are non-zero left derivations of  $R$  such an extent that  $g$  is a surjective on  $R$ , and  $g(y)\delta(x)=g(x)\delta(y)$ , for all  $x, y \in R$ . At that point  $\delta g$  cannot be a non-zero left derivation. One can consider (vii) and (iv), Theorem 3.2 as a proof of this outcome.

We now express a few outcomes of Theorem 3.2.

### Corollary 3.3

Give  $R$  a chance to be a prime ring of trademark not equivalent 2. Give  $\delta$  and  $g$  be left derivations of  $R$  with the end goal that  $g$  is a surjective on  $R$ , and  $g(y)\delta(x)=g(x)\delta(y)$ , for all  $x, y \in R$ . On the off chance that  $\delta$  and  $g$  are fulfill one of the states of Theorem 3.2, then either  $\delta=0$  or  $g=0$ .

Since a non-zero left derivation cannot be orthogonal on itself we see that (i), Theorem 3.2 yield the accompanying outcome.

### Corollary 3.4

Let  $R$  be a 2-torsion free semi-prime ring. And let  $\delta$  be a left derivation of  $R$  such that  $\delta$  is a surjective on  $R$ , and  $\delta(y)\delta(x)=\delta(x)\delta(y)$ , for all  $x, y \in R$ . If  $\delta^2(x)=0$ , for all  $x \in R$ , then  $\delta=0$ .

According to (vii), Theorem 3.2, we have,

### Corollary 3.5

Let  $R$  be a 2-torsion free semi-prime ring. And let  $\delta$  be a left derivation of  $R$  such that  $\delta$  is a surjective on  $R$ , and  $\delta(y)\delta(x)=\delta(x)\delta(y)$ , for all  $x, y \in R$ . If  $\delta^2$  is also a left derivation, then  $\delta=0$ .

Similarly, using (ix), Theorem 3.2, we obtain,

### Corollary 3.6

Let  $R$  be a 2-torsion free semi-prime ring. And let  $\delta$  be a left derivation of  $R$  such that  $\delta$  is a surjective on  $R$ , and  $\delta(y)\delta(x)=\delta(x)\delta(y)$ , for all  $x, y \in R$ . If there exist  $a, b \in R$  such that  $\delta^2(x)=xa+xb$ , for all  $x \in R$ , then  $\delta=0$ .

It is normal to inquire as to whether there is any association

between left derivations  $\delta$  and  $g$  of a ring  $R$ , If  $\delta^2=g^2$  or if  $\delta^2(x)=g(x)^2$ , for each  $x \in R$ . In the accompanying hypotheses, we give certifiable answer of this question.

### Theorem 3.7

Let  $R$  be a 2-torsion free semi-prime ring. Let  $\delta$  and  $g$  be left derivations of  $R$  such that  $g$  is a surjective on  $R$ , and  $g(y)\delta(x)=g(x)\delta(y)$ , for all  $x, y \in R$ . If  $\delta^2=g^2$ , then  $\delta+g$  and  $\delta-g$  are orthogonal.

**Proof:** From  $\delta^2=g^2$  it follows immediately that,  $(\delta+g)(\delta-g)+(\delta-g)(\delta+g)=0$ . Hence  $\delta+g$  and  $\delta-g$  are orthogonal by (vi), Theorem 3.2.

### Corollary 3.8

Let  $R$  be a prime ring of characteristic not equal 2. Let  $\delta$  and  $g$  be left derivations of  $R$  such that  $g$  is a surjective on  $R$ , and  $g(y)\delta(x)=g(x)\delta(y)$ , for all  $x, y \in R$ . If  $\delta^2=g^2$  then either  $\delta=g$  or  $\delta=-g$ .

### Theorem 3.9

Let  $R$  be a 2-torsion free semi-prime ring. Let  $\delta$  and  $g$  be left derivations of  $R$  such that  $g$  is a surjective on  $R$ , and  $g(y)\delta(x)=g(x)\delta(y)$ , for all  $x, y \in R$ . If  $\delta(x)^2=g(x)^2$ , for all  $x \in R$ , then  $\delta+g$  and  $\delta-g$  are orthogonal.

**Proof:** Note that  $(\delta+g)(x)(\delta-g)(x) + (\delta-g)(x)(\delta+g)(x)=0$ , for all  $x \in R$ . Now apply (iii), Theorem 3.2.

### Corollary 3.10

Let  $R$  be a prime ring of characteristic not equal 2. Let  $\delta$  and  $g$  be left derivations of  $R$  such that  $g$  is a surjective on  $R$ , and  $g(y)\delta(x)=g(x)\delta(y)$ , for all  $x, y \in R$ . If  $\delta(x)^2=g(x)^2$ , for all  $x \in R$ , then either  $\delta=g$  or  $\delta=-g$ .

## References

1. Posner E (1957) Derivations in Prime Rings. Proc Amer Math Soc 8: 1093-1100.
2. Brešar M, Vukman J (1990) On Left Derivations and Related Mappings. Proc Amer Math Soc 110(1): 7-16.
3. Ashraf M (2005) On Left  $(\theta, \phi)$ -Derivations in Prime Rings. Arch Math (Brno) 41: 157-166.
4. Ashraf M, Rehman N (2000) On Lie Ideal and Jordan Left Derivations of Prime Rings. Arch Math (Brno) 36(3): 201-206.
5. Ashraf M, Rehman N, Shaker A (2001) On Jordan Left Derivations of Lie Ideals in Prime Rings. Southeast Asian Bull Math 25(3): 379-382.
6. Deng Q (1992) On Jordan Left Derivations. Math J Okayama Univ 34: 145-147.
7. Vukman J (1997) Jordan Left Derivations on Semiprime Rings. Math J Okayama Uni 39: 1-6.
8. Zaidi SMA, Ashraf M, Al S (2004) On Jordan Ideals and Left  $(\theta, \theta)$  Derivations in Prime Rings. IJMMS 37: 1957-1964.
9. Brešar M, Vukman J (1991) Orthogonal Derivations and An Extension of A Theorem of Posner. Radovi Matematički 5: 237-246.
10. Argaç N, Nakajima A, Albaş E (2004) On Orthogonal Generalized Derivations of Semiprime Rings. Turk J Math 28: 185-194.
11. Lbaşı G, Aydin N (2007) Orthogonal Generalized  $(\sigma, \tau)$ -Derivations of Semiprime Rings. Sibirsk Mat Zh 48(6): 1222-1227.
12. Kharchenko VK (1991) Automorphisms and Derivations of Associative Rings. Kluwer Academic Publishers.