

Palladium-Catalyzed Regioselective 2-Carboethoxyethylation of 1H-Indoles By C-H Activation: One-Step Synthesis of Ethyl 2-(1H-Indol-2-Yl) Acetates

Yi Zhang^{1,2}, Jiang Wang², Yong Nian^{1,2}, Hongbin Sun¹ and Hong Liu^{1,2*}

¹State Key Laboratory of Natural Medicines, China Pharmaceutical University, 24 Tongjiaxiang, Nanjing 210009, P.R. China

²CAS Key Laboratory of Receptor Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, 555 Zuchongzhi Road, Shanghai 201203, P. R. China

Abstract

An efficient and convenient method was developed for the one-step synthesis of various substituted ethyl 2-(1H-indol-2-yl) acetates via a palladium-catalyzed regioselective cascade C-H activation reaction. Importantly, this practical approach can be carried out with readily accessible starting materials and exhibits excellent functional group compatibility.

Keywords: Ethyl 2-(1H-Indol-2-yl) Acetates; C-H Activation; Palladium; Norbornene

Introduction


2-substituted indoles are important structural motifs present in diverse biologically active molecules [1] and are precursors for a wide variety of alkaloids, such as vindoline [2], vindorosine [2], ellipticine [3], etc. Among them, 2-(1H-indol-2-yl)acetic acids constitute a valuable class of building blocks for natural product and natural product analogue syntheses (Figure 1) [4-7], combinatorial [8], diversity-oriented syntheses [9,10], and medicinal chemistry [11-18]. In addition, they can serve as highly attractive precursors for various chemical transformations, such as diazomethylation to 1-diazo-3-(2-indolyl)-2-propanone [19] and reduction to 2-(2-hydroxyethyl) indole [20]. Therefore, the development of efficient synthetic methods for these compounds has received much attention.

Hydrolysis of ethyl 2-(1H-indol-2-yl) acetates is an important route to 2-(1H-indol-2-yl) acetic acids. Accordingly, the literature describes several preparations of ethyl 2-(1H-indol-2-yl) acetates. For example, Capuano et al. demonstrated that the intramolecular Wittig reaction of 2-[(ω-alkoxycarbonyl) amino] benzyltri-phenylphosphonium salts produced 2-(1H-indol-2-yl)-acetate with a 78% yield [21]. Furthermore, Moody et al. reported that reductive cyclisation of the ethyl 4-(2-nitrophenyl)-acetoacetate using titanium(III) chloride in aqueous acetone gave 2-(1H-indol-2-yl) acetate with a 75% yield [22]. Despite producing good yields, these methods suffer from indispensable multi-step pre-transformations of commercially available starting materials, and are therefore of limited synthetic scope [23]. Wilkens et al. described a one-pot reaction of *N*-phenylhydroxylamine, benzaldehyde and ethyl 2,3-butadienoate followed by hydrolysis-mediated production of 2-(1H-indol-2-yl)acetate [24]. In spite of a 49% yield, this approach also had its drawbacks, including the use of unstable and relatively rare reactant (i.e. allenes) that might limit their broad applications. Osornio YM et al. and Guerrero MA et al. demonstrated that the direct intermolecular oxidative radical alkylation of indole under xanthate-mediated radical conditions afforded 2-(1H-indol-2-yl) acetate in 60% yield [25,26]. Nevertheless, the radical mechanism of this approach might limit its broad synthetic applications.

If indoles can be directly functionalized at the C2-position, the preparation of the target compound class would be more facile. Owing to the development of the transition-metal catalyzed C-H activation

chemistry, methods for regioselective direct C2/C3-alkenylation [27-33], alkynylation [34-41], cyano [42,43] and arylation [44-52] of indole nucleus have been well developed to date. However, regioselective C2-alkylation of indole is more challenging.

The Catellani reaction, a palladium-catalyzed norbornene-mediated cascade reaction, has been modified to achieve the direct functionalization of indoles [53-55]. Recently, Bressy et al. reported an intramolecular direct arylation of indoles based on modified Catellani conditions [56]. More recently, Jiao et al. reported a direct 2-alkylation reaction of indoles relying on a Pd(II)-catalyzed norbornene-mediated direct alkylation method for indoles, which regioselectively installs an alkyl group to the C2-position of free *N*-H indoles [57,58]. However, the original publication did not report the use of 2-bromoacetate as an alkylating reagent. To the best of our knowledge, a straight-forward approach for direct C2-alkoxycarbonylalkylation of indoles has not yet been well established. Given its high C2-regioselectivity and excellent



*Corresponding author: Hong Liu, State Key Laboratory of Natural Medicines, China Pharmaceutical University, 24 Tongjiaxiang, Nanjing 210009, P.R. China, Tel: 862150807042; E-mail: hbsun2000@yahoo.com; hliu@mail.shcnc.ac.cn

Received December 27, 2013; Accepted January 22, 2014; Published January 29, 2014

Citation: Zhang Y, Wang J, Nian Y, Sun H, Liu H (2014) Palladium-Catalyzed Regioselective 2-Carboethoxyethylation of 1H-Indoles By C-H Activation: One-Step Synthesis of Ethyl 2-(1H-Indol-2-Yl) Acetates. Organic Chem Curr Res 3: 124. doi:10.4172/2161-0401.1000124

Copyright: © 2014 Zhang Y, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

functional group tolerance, we envisioned to apply Bach's method to the synthesis of 2-(1H-indol-2-yl) acetates by employing indoles and ethyl 2-bromoacetate as the starting materials. As part of our continuing effort to assemble indole-based drug scaffolds [59-61], we here present our findings on one-step synthesis of various substituted ethyl 2-(1H-indol-2-yl)acetates via a palladium-catalyzed regioselective cascade C-H activation reaction. Various substituents are tolerated in this system in moderate to good yields. Our protocol highlights a facile one-step transformation from easily available starting material and excellent functional group compatibility.

Experimental

Unless otherwise noted, the reagents (chemicals) were purchased from commercial sources, and used without further purification. Water was deionized before used. Analytical thin layer chromatography (TLC) was HSGF 254 (0.15-0.2 mm thickness). Compound spots were visualized by UV light (254 nm). Column chromatography was performed on silica gel FCP 200-300. NMR spectra were run on 300 or 400 MHz instrument. Chemical shifts were reported in parts per million (ppm, δ) downfield from tetramethylsilane. Proton coupling patterns are described as singlet (s), doublet (d), triplet (t), quartet (q), multiplet (m), and broad (br). Low- and high-resolution mass spectra (LRMS and HRMS) were measured on spectrometer.

General procedure for synthesis of Ethyl 2-(1H-indol-2-yl) acetates

A vial equipped with a magnetic stir bar and a rubber stopper was charged with Pd(PhCN)₂Cl₂ (65 mg, 0.171 mmol), indole substrate (1.71 mmol), norbornene (321 mg, 3.41 mmol), NaHCO₃ (574 mg, 6.84 mmol) and capped with septa. The vial was evacuated and backfilled with argon and the process was repeated three times. A solution of water in DMF (0.5 M) was added via syringe as the solvent. Under argon, ethyl bromoacetate (0.41 mL, 3.41 mmol) was added via syringe, and then the resulting mixture was stirred at room temperature for 10 minutes. After that, the reaction mixture was then placed in a preheated oil bath at 70°C for appropriate time and vigorous stirring was applied. The reaction was monitored by TLC. Upon completion, the reaction mixture was cooled to room temperature, diluted with ethyl acetate, washed with water (twice) and brine (once), dried over Na₂SO₄, and concentrated. The residue was directly submitted to flash column chromatography (by dry loading) to afford the ethyl 2-(1H-indol-2-yl) acetates products as yellow oils.


Results and Discussion

We initiated our study by reacting indole with ethyl bromoacetate in the presence of norbornene and Pd(MeCN)₂Cl₂. However, the direct application of the reported optimized reaction conditions yielded diethyl 2,2'-(1H-indole-2,3-diyl) diacetate, a 2,3-disubstitued-indole product, as the major product.

According to the postulated catalytic cycle of Bach's approach [57,58], and distincts from the previous Catellani reaction, the arylpalladium(II) species in the terminating step undergoes proto-depalladation (i.e. hydrogenolysis), instead of a further palladium catalyzed carbon functionalization, such as Heck, Suzuki and Sonagahira reactions. Consequently, we speculated that for more active alkyl halide substrates, such as ethyl bromoacetate, it might be possible to quench the further substitution reaction and improve the yields of mono-2-substitued-indole products by the use of different

combination of solvent and base. Furthermore, we postulated that the phosphine ligands might have an effect on the rate of the substitution process as well.

With the aforementioned considerations in mind, we screened different reaction conditions for optimal results (Table 1). We conducted the first runs (Table 1, entries 1-6) employing the reported optimized Pd source [Pd(OAc)₂], base (K₂CO₃), reaction temperature (70°C) and time (14 hours) in different solvents (Table 1, entries 1-3). Notably, the higher water content in the solvents was found to significantly increase the selectivity for the mono-2-substitued-indole product 1 (Table 1), but led to a compromised conversion rate. These results suggest that water might serve as the hydrogen source in the catalytic system, which accelerates the norbornene-mediated cascade reaction by hydrolysing the final arylpalladium(II) species. As DMF/5 M water produced a better yield than the other solvents, it was selected as the solvent in the following tests. The effect of the Pd source was then investigated (Table 1, entries 7-10). Palladium species other than Pd(MeCN)₂Cl₂ and Pd(PhCN)₂Cl₂ were substantially less effective. Further screening of catalysts did not lead to better yields and demonstrated that the reaction may not proceed well in the presence of phosphine ligands which may reduce the palladium(II) species to palladium(0) species and interferes the normal palladium(II-IV) catalytic cycle [62]. Pd(PhCN)₂Cl₂ was chosen as the preferred catalyst because it provided higher conversion rates than the other catalysts. Given the possibility that excessive water in the solvent might hydrolyse the arylpalladium(II) species formed in the first oxidative addition step and lower the solubility of the intermediates, resulting in a cessation of the catalytic cycle and undesired conversion rate, we switched the solvent back to DMF with 0.5M H₂O. Finally, a brief screen of bases showed an interesting correlation between the alkalinity of the base and the yield of the 2,3-disubstitued-indole byproduct 2 (Table 1, entries




Entry	Pd Source	Solvent	Base	Yield (%) ^b	
				1	2
1	Pd(OAc) ₂	DMA / 0.5 M H ₂ O	K ₂ CO ₃	trace ^c	56
2	Pd(OAc) ₂	DMF / 0.5 M H ₂ O	K ₂ CO ₃	7	62
3	Pd(OAc) ₂	DMA / 5 M H ₂ O	K ₂ CO ₃	19	26
4	Pd(OAc) ₂	DMF / 5 M H ₂ O	K ₂ CO ₃	34	0
5	Pd(OAc) ₂	DMF / 2 M H ₂ O	K ₂ CO ₃	24	10
6	Pd(OAc) ₂	CH ₃ CN / 5 M H ₂ O	K ₂ CO ₃	15	21
7	Pd(Ph ₃ P) ₂ Cl ₂	DMF / 5 M H ₂ O	K ₂ CO ₃	trace ^c	0
8	Pd(dppf) ₂ Cl ₂ ·CH ₂ Cl ₂	DMF / 5 M H ₂ O	K ₂ CO ₃	trace ^c	0
9	Pd(MeCN) ₂ Cl ₂	DMF / 5 M H ₂ O	K ₂ CO ₃	46	0
10	Pd(PhCN) ₂ Cl ₂	DMF / 5 M H ₂ O	K ₂ CO ₃	52	0
11	Pd(PhCN) ₂ Cl ₂	DMF / 0.5 M H ₂ O	K ₂ CO ₃	8	79
12	Pd(PhCN) ₂ Cl ₂	DMF / 0.5 M H ₂ O	KHCO ₃	trace ^c	67
13	Pd(PhCN) ₂ Cl ₂	DMF / 0.5 M H ₂ O	NaHCO ₃	85	7
14	Pd(PhCN) ₂ Cl ₂	DMF / 0.5 M H ₂ O	KOAc	58	0
15	Pd(PhCN) ₂ Cl ₂	DMF / 0.5 M H ₂ O	NaOAc	42	0

^aGeneral reaction conditions: Pd (0.171 mmol), indole (1.71 mmol), norbornene (3.41 mmol), base (6.84 mmol), ethyl bromoacetate (3.41 mmol), solvent (8 mL, total), 70 °C, 14 h. ^bIsolated yield. ^cMonitored by LC-MS and TLC

Table 1: Optimization of the catalysis conditions^a.

11-15). Specifically, the yields of the 2,3-disubstituted-indole products increased significantly as the alkalinity of the base increased, with NaHCO₃ affording the target products in desirable yields. On the other hand, insufficient basicity led to undesired conversion rates (Table 1, entries 14 and 15). Thus, DMF with 0.5M H₂O as the solvent, NaHCO₃ as the base, a reaction temperature of 70 °C, and a reaction time of 14 hours were selected as the optimal conditions. It was noteworthy that in all tests, the other possible substituted byproducts, N- and 3-substituted products 3, 4 (Table 1) were not detected.

Encouraged by these initial results, we next proceeded to examine the general utility of the Pd(PhCN)₂Cl₂/norbornene catalytic system for synthesis of a wide range of ethyl 2-(1H-indol-2-yl) acetates (Table 2). Indoles with electron-donating and electron-drawing substituents at the various positions smoothly participated in the C2-ethoxycarbonylmethylation reaction. Four regioisomeric methylindoles effectively participated in the cascade reactions to afford equally good levels of yields (Table 2, entries 1-4). The reaction worked well with different halogens on various positions on the benzene ring




Entry	R ₁	Product	Yield ^b
1	4-Me (5a)	6a	67%
2	5-Me (5b)	6b	71%
3	6-Me (5c)	6c	69%
4	7-Me (5d)	6d	65%
5	5-OMe (5e)	6e	80%
6	4-F (5f)	6f	75%
7	5-F (5g)	6g	83%
8	6-F (5h)	6h	82%
9	7-F (5i)	6i	78%
10	5-Cl (5j)	6j	85%
11	5-Br (5k)	6k	84%
12	5-I (5l)	6l	71%
13	5-CN (5m)	6m	89%
14	5-NO ₂ (5n)	6n	91%

15			58%
16			35%
17			trace ^c

^aReaction conditions: Pd(PhCN)₂Cl₂ (0.171 mol), indole (1.71 mmol), norbornene (3.41 mmol), NaHCO₃ (6.84 mmol), ethyl bromoacetate (3.41 mmol), DMF with 0.5 M H₂O (8 mL, total), 70 °C, 14 h. ^bIsolated yield. ^cMonitored by LC-MS

Table 2: Palladium-catalyzed synthesis of various ethyl 2-(1H-indol-2-yl) acetates^a.



(Table 2, entries 6-12), which has three main advantages. First, halogen substituents could be used to adjust the polarity, lipophilicity, and metabolic stability of dyes or pharmaceuticals. Second, halogens allow further modification using cross-coupling reactions for the elaboration of molecule libraries. Third, halogen substituents on various positions can serve as invaluable building blocks for enrichment of Structure-Activity Relationships in medicinal chemistry. It was noteworthy to mention that 5-iodoindole, a potential substrate for classic Catellani reaction, was readily tolerated in our catalytic system and gave the desired product 6l in good yield. Interestingly, electron-deficient indoles reacted better than their electron-rich counterparts to give good yields of 2-(1H-indol-2-yl) acetates under the optimal conditions (Table 2, entries 6-15). Moreover, the reaction could be extended to functionalized azindoles to give the expected product 6o in modest yield. We also noted that a more bulky 3-position blocked the substrate, 3-methyl-indole, produced the desired product 3 in a 35% yield (Table 2, entry 16). Control experiment showed that this catalytic system did not work in the absence of norbornene. By contrast, N-methylindole failed to participate in the reaction and remained mostly unchanged even after extension of reaction time (Table 2, entry 17). These results are consistent with Bach and Jiao's latest work regarding the mechanism of regioselective Pd-catalyzed, norbornene-mediated alkylation of indoles. It has been proved that N1-norbornene type palladacycle rather than originally proposed C3-norbornene type palladacycle is formed as the key intermediate in this catalytic cycle [58].

On the basis of above observations, recent publications on the mechanism of the Catellani reaction [63] and extensive mechanistic work by Jiao et al. [58], a plausible mechanism for the observed reaction is proposed in Scheme 1. The norbornene-mediated cascade C-H activation process proceeds as follows: a. N1-position direct palladation; b. *syn*-aminopalladation of norbornene; c. irreversible palladacycle formation, leading to C2-position ortho-C-H palladation; d. oxidative addition with an alkyl halide to generate palladium IV species; e. reductive elimination of the palladium IV species; f. norbornene expulsion; and g. release of the 2-alkyl indole product and

regeneration the Pd(II) species (Scheme 1). The 2-monosubstituted product of this catalytic cycle might undergo further electrophilic substitution process [56] to afford the 2,3-disubstituted-indole product.

Conclusion

To conclude, a practical Pd(PhCN)₂Cl₂/norbornene catalytic system for the versatile and one-step synthesis of ethyl 2-(1H-indol-2-yl)acetates has been developed. This practical method features the ability to rapidly and efficiently synthesize various substituted ethyl 2-(1H-indol-2-yl)acetates as precursors to a class of valuable synthetic building blocks, 2-(1H-indol-2-yl)acetic acids. In addition, the protocol exhibits excellent functional group compatibility, leading to valuable derivatives that are not readily available by conventional methods. The present reaction also enables a new access to 2,3-dialkoxy-carbonylalkylated indole derivatives which can be applied to indole fused heterocycles and indole alkaloid synthesis.

Acknowledgements

We gratefully acknowledge financial support from the National Natural Science Foundation of China (Grants 21021063, 91229204, and 81025017), National S&T Major Projects (2012ZX09103101-072, 2012ZX09301001-005, and 2013ZX09507-001), Sponsored by Program of Shanghai Subject Chief Scientist (Grant 12XD1407100).

References

- Katritzky AR, Li J, Stevens CV (1995) Facile Synthesis of 2-Substituted Indoles and Indolo[3,2-b]carbazoles from 2-(Benzotriazol-1-ylmethyl)indole. *J Org Chem* 60: 3401-3404.
- Kuehne ME, Podhorez DE, Mulamba T, Bornmann WG (1987) Biomimetic alkaloid syntheses. 15. Enantioselective syntheses with epichlorohydrin: total syntheses of (+)-, (-) and (+,-)-vindoline and a synthesis of (-)-vindorosine. *J Org Chem* 52: 347-353.
- Modi SP, Zayed AH, Archer S (1989) Synthesis of 6-substituted 7H-pyrido[4,3-c]carbazoles. *J Org Chem* 54: 3084-3087.
- Martin CL, Overman LE, Rohde JM (2008) Total synthesis of (+/-)-actinophyllic acid. *J Am Chem Soc* 130: 7568-7569.
- Sundberg RJ, Hong J, Smith SQ, Sabat M, Tabakovic I (1998) Synthesis and Oxidative Fragmentation of Catharanthine Analogs. Comparison to the Fragmentation - Coupling of Catharanthine and Vindoline. *Tetrahedron* 54: 6259-6292.
- Gul W, Hammond NL, Yousaf M, Peng J, Holley A, et al. (2007) Chemical transformation and biological studies of marine sesquiterpene (S)-(+)-curcuphenol and its analogs. *Biochim Biophys Acta* 1770: 1513-1519.
- Sripha K, Zlotos DP, Holzgrabe U, Ruchirawat S (2011) A New Synthetic Approach to Pentacyclic Ring, 6,7,14,15-Tetrahydro[1,5]diazocino[1,2-a:6,5-a']-8-dihydrodiindole. *ETEROCYCLES* 83: 2627-2633.
- Boger DL, Goldberg J, Satoh S, Ambroise Y, Cohen SB, et al. (2000) Non-Amide-Based Combinatorial Libraries Derived from N-Boc-Iminodiacetic Acid: Solution-Phase Synthesis of Piperazinone Libraries with Activity Against LEF-1/ β -Catenin-Mediated Transcription. *Helv Chim Acta* 83: 1825-1845.
- Royer D, Wong YS, Plé S, Chiaroni A, Diker K, et al. (2008) Diastereodivergence and appendage diversity in the multicomponent synthesis of aryl-pyrrolo-tetrahydrocarbazoles. *Tetrahedron* 64: 9607-9618.
- Waldmann H, Kühn M, Liu W, Kumar K (2008) Reagent-controlled domino synthesis of skeletally-diverse compound collections. *Chem Commun (Camb)* : 1211-1213.
- Méndez-Andino J, Colson AO, Denton D, Mitchell MC, Cross-Doersen D, et al. (2007) MCH-R1 antagonists based on an arginine scaffold: SAR studies on the amino-terminus. *Bioorg Med Chem Lett* 17: 832-835.
- Potter AJ, Ray S, Gueritz L, Nunns CL, Bryant CJ, et al. (2010) Structure-guided design of alpha-amino acid-derived Pin1 inhibitors. *Bioorg Med Chem Lett* 20: 586-590.
- Weber D, Berger C, Eickelmann P, Antel J, Kessler H (2003) Design of selective peptidomimetic agonists for the human orphan receptor BRS-3. *J Med Chem* 46: 1918-1930.
- Harikrishnan LS, Srivastava N, Kayser LE, Nirschl DS, Kumaragurubaran K, et al. (2012) Identification and optimization of small molecule antagonists of vasoactive intestinal peptide receptor-1 (VIPR1). *Bioorg Med Chem Lett* 22: 2287-2290.
- Yang K, Wang Q, Su L, Fang H, Wang X, et al. (2009) Design and synthesis of novel chloramphenicol amine derivatives as potent aminopeptidase N (APN/CD13) inhibitors. *Bioorg Med Chem* 17: 3810-3817.
- Rode HB, Sos ML, Grütter C, Heynck S, Simard JR, et al. (2011) Synthesis and biological evaluation of 7-substituted-1-(3-bromophenylamino)isoquinoline-4-carbonitriles as inhibitors of myosin light chain kinase and epidermal growth factor receptor. *Bioorg Med Chem* 19: 429-439.
- McInnis CE, Blackwell HE (2011) Thiolactone modulators of quorum sensing revealed through library design and screening. *Bioorg Med Chem* 19: 4820-4828.
- Meng Q, Zhao B, Xu Q, Xu X, Deng G, et al. (2012) Indole-propionic acid derivatives as potent, S1P3-sparing and EAE efficacious sphingosine-1-phosphate 1 (S1P1) receptor agonists. *Bioorg Med Chem Lett* 22: 2794-2797.
- Salim M, Capretta A (2000) Intramolecular Carbenoid Insertions: the Reactions of α -Diazoketones Derived from Pyrrolyl and Indolyl Carboxylic Acids with Rhodium(II) Acetate. *Tetrahedron* 56: 8063-8069.
- Sripha K, Zlotos DP, Buller S, Mohr K (2003) 6,7,14,15-Tetrahydro-15aH-azocino[1,2-a:6,5-b']diindole. Synthesis of a novel pentacyclic ring system. *Tetrahedron Lett* 44: 7183-7186.
- Capuano L, Ahlhelm A, Hartmann H (1986) New Syntheses of 2-Acylbenzofurans, 2-Acylindoles, 2-Indolylcarboxylates, and 2-Quinolones by Intramolecular Wittig Reaction. *Chem Ber* 119: 2069-2074.
- Moody CJ, Rahimtoola KF (1990) Diels-Alder reactivity of pyrano[4,3-b]indol-3-ones, indole 2,3-quinodimethane analogues. *J Chem Soc, Perkin Trans 1* 673-679.
- Attia MI, Guclu D, Hertlein B, Julius JP, Witt-Enderby A, et al. (2007) Synthesis, NMR conformational analysis and pharmacological evaluation of 7,7a,13,14-tetrahydro-6H-cyclobuta[b]pyrimido[1,2-a : 3,4-a']diindole analogues as melatonin receptor ligands. *Organic & Biomolecular Chemistry* 5: 2129-2137.
- Wilkens J, Kühling A, Blechert S (1987) Hetero-cope rearrangements - v1 short and stereoselective syntheses of 2-vinylindoles by a tandem-process. *Tetrahedron*: 43, 3237-3246.
- Osornio YM, Cruz-Almanza R, Jiménez-Montaño V, Miranda LD (2003) Efficient, intermolecular, oxidative radical alkylation of heteroaromatic systems under "tin-free" conditions. *Chem Commun (Camb)* : 2316-2317.
- Guerrero MA, Miranda LD (2006) Et3B-Mediated radical alkylation of pyrroles and indoles. *Tetrahedron Letters* 47: 2517-2520.
- Grimster NP, Gauntlett C, Godfrey CR, Gaunt MJ (2005) Palladium-catalyzed intermolecular alkenylation of indoles by solvent-controlled regioselective C-H functionalization. *Angew Chem Int Ed Engl* 44: 3125-3129.
- García-Rubia A, Gómez Arrayás R, Carretero JC (2009) Palladium(II)-catalyzed regioselective direct C2 alkenylation of indoles and pyrroles assisted by the N-(2-pyridyl)sulfonyl protecting group. *Angew Chem Int Ed Engl* 48: 6511-6515.
- Ding Z, Yoshikai N (2012) Mild and efficient C2-alkenylation of indoles with alkynes catalyzed by a cobalt complex. *Angew Chem Int Ed Engl* 51: 4698-4701.
- Kandukuri SR, Schiffner JA, Oestreich M (2012) Aerobic palladium(II)-catalyzed 5-endo-trig cyclization: an entry into the diastereoselective C-2 alkenylation of indoles with tri- and tetrasubstituted double bonds. *Angew Chem Int Ed Engl* 51: 1265-1269.
- Baran PS, Richter JM (2004) Direct coupling of indoles with carbonyl compounds: short, enantioselective, gram-scale synthetic entry into the hapalindole and fischerindole alkaloid families. *J Am Chem Soc* 126: 7450-7451.
- Pan S, Ryu N, Shibata T (2012) Ir(III)-catalyzed C-H bond alkylation of C2-position of indole with alkenes: selective synthesis of linear or branched 2-alkylindoles. *J Am Chem Soc* 134: 17474-17477.
- Byers JH, Campbell JE, Knapp FH, Thissell JG (1999) Radical aromatic substitution via atom-transfer addition. *Tetrahedron Lett* 40: 2677-2680.

34. Gu Y, Wang X (2009) Direct palladium-catalyzed C-3 alkynylation of indoles. *Tetrahedron Lett* 50: 763-766.
35. Brand JP, Charpentier J, Waser J (2009) Direct alkynylation of indole and pyrrole heterocycles. *Angew Chem Int Ed Engl* 48: 9346-9349.
36. Brand JP, Chevalley C, Waser J (2011) One-pot gold-catalyzed synthesis of 3-silylethynyl indoles from unprotected o-alkynylanilines. *Beilstein J Org Chem* 7: 565-569.
37. Brand JP, Chevalley C, Scopelliti R, Waser J (2012) Ethynyl benziodoxolones for the direct alkynylation of heterocycles: structural requirement, improved procedure for pyrroles, and insights into the mechanism. *Chemistry* 18: 5655-5666.
38. de Haro T, Nevado C (2010) Gold-catalyzed ethynylation of arenes. *J Am Chem Soc* 132: 1512-1513.
39. Yang L, Zhao L, Li CJ (2010) Palladium-catalyzed direct oxidative Heck-Cassar-Sonogashira type alkynylation of indoles with alkynes under oxygen. *Chem Commun (Camb)* 46: 4184-4186.
40. Ruano JL, Alemán J, Marzo L, Alvarado C, Tortosa M, et al. (2012) Arylsulfonylacetylenes as alkynylating reagents of Csp²-H bonds activated with lithium bases. *Angew Chem Int Ed Engl* 51: 2712-2716.
41. García Ruano JL, Alemán J, Marzo L, Alvarado C, Tortosa M, et al. (2012) Expanding the scope of arylsulfonylacetylenes as alkynylating reagents and mechanistic insights in the formation of Csp²-Csp and Csp³-Csp bonds from organolithiums. *Chemistry* 18: 8414-8422.
42. Yan G, Kuang C, Zhang Y, Wang J (2010) Palladium-catalyzed direct cyanation of indoles with K(4)[Fe(CN)(6)]. *Org Lett* 12: 1052-1055.
43. Ghosh K, Sarkar AR, Samadder A, Khuda-Bukhsh AR (2012) Pyridinium-based fluororeceptors as practical chemosensors for hydrogen pyrophosphate (HP2O7(3-)) in semiaqueous and aqueous environments. *Org Lett* 14: 4314-4317.
44. Lane BS, Sames D (2004) Direct C-H bond arylation: selective palladium-catalyzed C2-arylation of N-substituted indoles. *Org Lett* 6: 2897-2900.
45. Wang X, Lane BS, Sames D (2005) Direct C-arylation of free (NH)-indoles and pyrroles catalyzed by Ar-Rh(III) complexes assembled in situ. *J Am Chem Soc* 127: 4996-4997.
46. Lane BS, Brown MA, Sames D (2005) Direct palladium-catalyzed C-2 and C-3 arylation of indoles: a mechanistic rationale for regioselectivity. *J Am Chem Soc* 127: 8050-8057.
47. Deprez NR, Kalyani D, Krause A, Sanford MS (2006) Room temperature palladium-catalyzed 2-arylation of indoles. *J Am Chem Soc* 128: 4972-4973.
48. Stuart DR, Fagnou K (2007) The catalytic cross-coupling of unactivated arenes. *Science* 316: 1172-1175.
49. Lebrasseur N, Larrosa I (2008) Room temperature and phosphine free palladium catalyzed direct C-2 arylation of indoles. *J Am Chem Soc* 130: 2926-2927.
50. Phipps RJ, Grimster NP, Gaunt MJ (2008) Cu(II)-catalyzed direct and site-selective arylation of indoles under mild conditions. *J Am Chem Soc* 130: 8172-8174.
51. Cornella J, Lu P, Larrosa I (2009) Intermolecular decarboxylative direct C-3 arylation of indoles with benzoic acids. *Org Lett* 11: 5506-5509.
52. Potavathi S, Pereira KC, Gorelsky SI, Pike A, LeBris AP, et al. (2010) Regioselective oxidative arylation of indoles bearing N-alkyl protecting groups: dual C-H functionalization via a concerted metalation-deprotonation mechanism. *J Am Chem Soc* 132: 14676-14681.
53. Catellani M (2008) Novel methods of aromatic functionalization using palladium and norbornene as a unique catalytic system. *Top Organomet Chem* 14: 21-53.
54. Catellani M, Motti E, Della Ca' N (2008) Catalytic sequential reactions involving palladacycle-directed aryl coupling steps. *Acc Chem Res* 41: 1512-1522.
55. Martins A, Mariampillai B, Lautens M (2010) Synthesis in the key of Catellani: norbornene-mediated ortho C-H functionalization. *Top Curr Chem* 292: 1-33.
56. Bressy C, Alberico D, Lautens M (2005) A route to annulated indoles via a palladium-catalyzed tandem alkylation/direct arylation reaction. *J Am Chem Soc* 127: 13148-13149.
57. Jiao L, Bach T (2011) Palladium-catalyzed direct 2-alkylation of indoles by norbornene-mediated regioselective cascade C-H activation. *J Am Chem Soc* 133: 12990-12993.
58. Jiao L, Herdtweck E, Bach T (2012) Pd(II)-catalyzed regioselective 2-alkylation of indoles via a norbornene-mediated C-H activation: mechanism and applications. *J Am Chem Soc* 134: 14563-14572.
59. Ye D, Wang J, Zhang X, Zhou Y, Ding X, et al. (2009) Gold-catalyzed intramolecular hydroamination of terminal alkynes in aqueous media: efficient and regioselective synthesis of indole-1-carboxamides. *Green Chem* 11: 1201-1208.
60. Wang J, Zhou S, Lin D, Ding X, Jiang H, et al. (2011) Highly diastereo- and enantioselective synthesis of syn-β-substituted tryptophans via asymmetric Michael addition of a chiral equivalent of nucleophilic glycine and sulfonylindoles. *Chem Commun* 47: 8355-8357.
61. Feng E, Zhou Y, Zhao F, Chen X, Zhang L, et al. (2012) Gold-catalyzed tandem reaction in water: an efficient and convenient synthesis of fused polycyclic indoles. *Green Chem* 14: 1888-1895.
62. Tolnai GL, Ganss S, Brand JP, Waser J (2013) C2-selective direct alkynylation of indoles. *Org Lett* 15: 112-115.
63. Maestri G, Motti E, Della Ca' N, Malacria M, Derat E, et al. (2011) Of the ortho effect in palladium/norbornene-catalyzed reactions: a theoretical investigation. *J Am Chem Soc* 133: 8574-8585.

Citation: Zhang Y, Wang J, Nian Y, Sun H, Liu H (2014) Palladium-Catalyzed Regioselective 2-Carboethoxyethylation of 1H-Indoles By C-H Activation: One-Step Synthesis of Ethyl 2-(1H-Indol-2-Yl) Acetates. *Organic Chem Curr Res* 3: 124. doi:[10.4172/2161-0401.1000124](https://doi.org/10.4172/2161-0401.1000124)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscripts as E- mail: editor.organicchemistry@omicsonline.org