

Physical Activity and Sedentary Behaviors as Risk Factors of Obesity among Rural Adolescents

Nasser Alqahtani*, Jane Scott and Shahid Ullah

Flinders University, Adelaide, Australia

*Corresponding author: Nasser Alqahtani, Ministry of Health, Adelaide, Riyadh PO Box 42248, Saudi Arabia, Tel: +966530102200; E-mail: nasadiet23@gmail.com

Received date: January 17, 2015, Accepted date: January 29, 2015, Published date: February 5, 2015

Copyright: © 2015 Alqahtani N, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Objective: The aim of this study is to explore relationship between Physical activity and sedentary behavior (screen time) and obesity among adolescents in rural areas.

Methodology: Cross-sectional study was conducted in 2011 using a multistage randomization method. It surveyed 370 from rural areas of Riyadh region in Saudi Arabia.

Results: This study found a strong association between the moderate to vigorous METs and overweight and obesity among male participants ($p < 0.001$). Prolonged screen time has strong association with overweight in males ($p = 0.01$) and females ($p < 0.001$). Similar association found between screen time and obesity in males and females ($p < 0.001$).

Conclusion: low physical activity and prolonged screen time are the main predictors for overweight or obesity or both of them in rural adolescents.

Keywords: Overweight; Obesity; Adolescents; Saudi arabia

Introduction

Overweight and obesity are descriptions of excessive body fat [1]. The WHO estimates that, globally, approximately 20 million children under age five are overweight [2], and Kosti and Panagiotakos (2006) have estimated that about 155 million children and adolescents are overweight; that is, one in every 10 children globally are afflicted with overweight problems. Saudi Arabia is experiencing an alarming prevalence of overweight and obesity as one in every three Saudis is obese [3].

Reports by the Saudi Diabetes and Endocrinology Society (SDS) indicate that in 2012 about 70 % of Saudi adults were obese, a statistic which is consistent with the predictions of the Saudi Obesity Research Centre (SORC) [4,5]. In the last two decades several cross-sectional studies have found that the average body fat of Saudi adolescents has been increasing nationally [6-9]. The proportion of Saudi adolescents who are overweight or obese is now very high, their combined prevalence being 47 % [10,11].

Obesity in adolescents has been associated with chronic illnesses such as cardiovascular disease [12,13], pre-diabetes [14] and Type 2 diabetes [15,16]. Moreover, obese adolescents are at serious risk of sleep apnoea, joint problems, and social and psychological problems [17-19].

So far, research into obesity in Saudi Arabia has focussed mostly on males in urban cities and there are no national data about the prevalence of overweight and obesity in rural areas; additionally, relatively few studies have investigated overweight and obesity in adolescent females. In a cross-sectional study in urban area, the

researchers noting that the general lack of participation in physical activity was a predictor of obesity in adolescents (OR=1.35; 95% CI 1.06-1.94). This study aim to evaluates the influences physical activity, and screen time on overweight and obesity in adolescents in rural districts of Saudi Arabia.

Methodology

This is a cross-sectional study conducted among adolescents in Riyadh region, Saudi Arabia in 2011. Schools and participants were selected using multistage randomization method. Thirty five eligible participants were chosen from twenty five elementary and secondary schools. This project surveyed 370 school children (196 males and 174 females) aged 14 to 19 years old.

physical activity and sedentary behaviour were measured using the physical- activity questionnaire which has been designed and validated for use with Saudi Arabian adolescents [20]. This study was approved by the Social and Behavioral Research Ethics Committee of Flinders University (Project No 4793: 11 May 2010).

Data-entry and statistical analyses were conducted using Excel 2010 and SPSS software version 19.0 (SPSS, Inc., Chicago IL.) and STATA version 12.0 (StataCorp. 2011). The two-sided tests were performed for all analyses and the level of significance was set at $P < 0.05$. Where appropriate. Descriptive statistics were expressed as median and Interquartile ranges (IQR) were reported for the data.

Results

Table 1 show that there was no significant association between weight-status and age amongst rural male and female participants.

Age and gender	Normal%	OW%	Obese%	P*
Males				
14-15	51.8	15.7	32.5	0.8
16-17	58.4	18.2	23.4	
18-19	55.6	16.7	27.8	
Females				
14-15	56.9	27.6	14.5	0.92
16-17	55.9	25.4	18.6	
18-19	51.3	28.2	20.5	

P* values are based on Chi-square test between urban and rural areas; P values of less than 0.05 were considered statistically significant.

Table 1: Weight status of participants by age group and gender

Table 2 shows the median and IQR of screen time and the MET of moderate and vigorous physical activities for normal, overweight, and obese adolescents in rural areas. Normal weight males spent less time on screen activities and more time exercising than did overweight and obese males ($p < 0.001$), but there were no particular differences

between overweight and obese males. In the same regions, normal weight females spent less time on screen-based activities and more time on METs than did overweight and obese females ($p < 0.001$) but there were no differences between overweight and obese females.

Screen time and physical activity	(N=370)			P*
	Normal (n=205)	OW (n=80)	Obese (n=85)	
Male	(n=108)	(n=33)	(n=55)	
Screen time (hours per day)	1.8 ^a (1.2-2.4)	2.6 ^b (1.8-2.9)	4.3 ^c (2.8-5.1)	<0.001
Moderate METs per week	3360 ^a (2137-6063)	1260 ^b (420-2152)	1260 ^b (840-1620)	<0.001
Vigorous METs per week	6160 ^a (3918-11116)	2310 ^b (770-3946)	2310 ^b (1540-2970)	<0.001
Total moderate to vigorous METs per week	9520 ^a (6056-7180)	3570 ^b (1190-6098)	3570 ^b (2380-4590)	<0.001
Female	(n=97)	(n=47)	(n=30)	
Screen time (hours per day)	1.4 ^a (1.0-2.0)	3.0 ^b (1.7-3.2)	2.6 ^b (1.9-4.0)	<0.001
Moderate METs per week	1260 ^a (840-1890)	630 ^b (630-1050)	652 ^b (630-1102)	<0.001
Vigorous METs per week	2131 ^a (1540-3465)	1155 ^b (1155-1925)	1196 ^b (1155-2021)	<0.001
Total moderate to vigorous METs per week	3570 ^a (2380-5355)	1785 ^b (1685-2975)	1848 ^b (1785-3123)	<0.001

*P values are based on Kruskal-Wallis Test.

Differing superscript letters denote significantly different column proportions at the 0.05 level. Absence of superscript letters denotes no significant difference between column proportions.

Table 2: Median (Inter-quartile range) screen time and physical activities for normal, overweight and obese participants by gender.

In order to determine more precisely the net independent effects that any particular factor had on weight status. The multivariate multinomial logistic regression model was used in this study [21,22] to help explain the confounding effects of the factors in the model. Tables 3 show the final stage of multivariate multinomial logistic regression model. Among rural males, total moderate to vigorous Kilo METs was

inversely associated with the risk of both overweight (OR=0.84; 95% CI 0.75-0.95; $p = 0.01$) and obesity (OR=0.71; 95% CI 0.56-0.90; $p < 0.01$). In addition, the amount of time spent on screen-based activities was directly associated with risk of overweight (OR=2.58; 95% CI 1.19-5.58; $p = 0.02$) and obesity (OR=9.63; 3.70-25.06; $p < 0.001$). Finally, the amount of time spent on screen-based activities was

directly associated with the risk of overweight among rural females (OR=4.75 95% CI 1.96-11.47; P<0.01) and obesity (OR=7.41 95% CI 2.49-22.05; p<0.001).

	Male (n=196)						Female (n=174)					
	Overweight ^b			Obese ^c			Overweight ^b			Obese ^c		
	OR	95% CI	P	OR	95% CI	P	OR	95% CI	P	OR	95% CI	P
Total Kilo METs per week	0.84	0.75-0.95	0.01	0.71	0.56-0.90	<0.01	0.79	0.58-1.06	0.12	0.82	0.47-1.44	0.50
Screen time (hours per day)	2.58	1.19-5.58	0.02	9.63	3.70-25.06	<0.001	4.75	1.96-11.47	<0.01	7.41	2.49-22.05	<0.001

^a the reference is normal weight equivalent to an adult BMI of 18.50 to 24.99 kg/m²
^b overweight equivalent to an adult BMI of 25.00 to 29.99 kg/m²
^c obesity equivalent to an adult BMI of equal or more than 30 kg/m²

Table 3: Multinomial logistic regression model (Odds ratios and 95% confidence intervals) of dietary factors, total Kilo METs, and screen time for different weight status for male and female rural adolescents.

Discussion

Physical activity

Health research has consistently found that the general increase in overweight and obesity has an association with low and/or reduced levels of physical activity by children [23-28]. Also, as noted by Patrick et al. among adolescents in the USA low physical activity has been associated with an increased risk of having a high BMI [24].

Physical activity guidelines recommend that adolescents should participate in moderate to vigorous activity for at least 60 minutes every day [29-31].

The results of this study show that there is a low prevalence of physical activity among Saudi adolescents. Only 29% of male and 25% of female participants were participating in moderate or vigorous activity for at least one hour daily. The proportion is similar to that reported in two earlier Saudi studies which reported that only 30% of adolescents aged 12 to 20 years met the requirements for physical activity [32,33].

There are several possible explanations why Saudi adolescents have low levels of physical activity. Firstly, young Saudis prefer to use cars instead of walking, even for short distances such as going to and from school [34]. For cultural and religious reasons female students cannot practice any physical activity in their schools. However, in April 2013, the Ministry of Education allowed female students in private schools to play sports and games such as volleyball and soccer [35].

This study found that the female participants perform significantly less physical activity in comparison to the male participants, this result confirming other studies of Arab children and adolescents [36-38]. A survey of 9,433 Saudi adolescents by Collison et al in 2010 found that males were spending significantly more time in exercise than females (p 0<0.05)[36]. In a cross-sectional study among Palestinian children, male children participated more in physical activities than females [37], and in the United Arab Emirates, the amount of physical activity among female adolescents was very low compared to their peers in the UK, Sweden and Australia [38].

This study found a strong association between the moderate to vigorous METs and overweight and obesity. The risk of overweight and obesity can be reduced by performing moderate to vigorous activity.

Insufficient physical activity (less than 30 minutes per week) by Saudi male adolescents in an earlier Saudi study was found to be associated with obesity (OR=1.6; 95% CI 1.01- 2.62) [39]. This confirms the results of a survey conducted in the South province by Mahfouz in 2008, the researchers noting that the general lack of participation in physical activity was a predictor of obesity in adolescents (OR=1.35; 95% CI 1.06-1.94)[40]. In a Spanish cross-sectional study among 2,859 adolescents, high levels of moderate to vigorous exercise (more than 15 minutes per day) were associated with lower abdominal obesity [41].

Screen time

While the WHO does not have any guidelines for screen time, other health organisations have recommendations for the maximum time young people should spend watching screens of any sort. The Australian Department of Health and Ageing (ADHA), and other international authorities such as the American Academy of Paediatrics (AAP), recommend that children and adolescents should not spend more than two hours per day viewing TV [31, 42]. Based on this recommendation, only 25% of male and 22% of female participants in this study met the recommendation of less than or equal to 2 hours of watching TV. The average time participants spent viewing television was 2.4 hours per day, a figure similar to the 2.5 hours reported in a study of adolescents aged 14 to 16 years in the United Arab Emirates [38]. A survey of Italian adolescents reported an average of 2.8 hours per day [43].

While the average hours the Saudi participants spent watching television was similar to, or somewhat lower than, other studies, the proportion of children watching more than the recommended two hours of television is higher than that reported in other surveys. For instance, in this study 53% of subjects watched more than two hours of television per day, a figure higher than the 47.6% reported for adolescents in the Greek part of Cyprus [44]. In Finland, 44% of male and 48% of female adolescents had reported watching TV more than

two hours daily [45]. In a Chinese study among adolescents aged 13-18, 44.3% of male and 34.7% female participants spent more than two hours of watching TV per day [46]. From the 1999 to 2006 American National Health and Nutrition Examination Survey (NHANES), the prevalence of the USA adolescent who watch TV for more than 2 hours/day was 29% [47].

This study found a strong association between screen time and weight, the risk of overweight and obesity being directly associated with the amount of screen time. However it is not the lack of movement that is the only causal factor, and previous research has concluded that sedentary behaviour is associated with other obesogenic practices such as consuming high calories snack and sweetened carbonated drinks [48, 49]. That is, people tend to snack on high-calorie food while watching television, foods which are highly advertised on TV.

Conclusion

This research found that, among rural Saudi adolescents, low physical activity and prolonged screen time are the common predictors for overweight or obesity or both of them.

This research provides, for the first time, an exploration of the determinants of overweight and obesity among adolescents in rural areas. These results are valuable because they will provide the basis for action to help resolve this health issue - at least among this age group.

References

1. Fricker J (1988) A positive correlation between energy intake and body mass index in a population of 1312 overweight subjects. *International Journal of Obesity* 13: 673-681.
2. WHO (2010) Childhood overweight and obesity. World Health Organization [website].
3. Al-Nozha MM, Arafah MR, Al-Mazrou YY, Al-Maatouq MA, Khan NB, et al. (2004) Coronary artery disease in Saudi Arabia. *Saudi Med J* 25: 1165-1171.
4. SDES (2013) Obesity management.
5. Al-Hazzaa HM (2007) Rising trends in BMI of Saudi adolescents: evidence from three national cross sectional studies. *Asia Pac J Clin Nutr* 16: 462-466.
6. Al-Hazzaa HM (2007) Prevalence and trends in obesity among school boys in Central Saudi Arabia between 1988 and 2005. *Saudi Med J* 28: 1569-1574.
7. Al-Hazzaa HM (2011) Physical activity, sedentary behaviors and dietary habits among Saudi adolescents relative to age, gender and region. *International Journal of Behavioral Nutrition and Physical Activity* 8: 140.
8. Abalkhail B (2002) Overweight and obesity among Saudi Arabian children and adolescents between 1994 and 2000. *East Mediterr Health J* 8: 470-479.
9. Khalid Mel-H (2008) Is high-altitude environment a risk factor for childhood overweight and obesity in Saudi Arabia? *Wilderness Environ Med* 19: 157-163.
10. Al-Dossary SS, Sarkis PE, Hassan A, Ezz El Regal M, Fouada AE (2010) Obesity in Saudi children: a dangerous reality. *East Mediterr Health J* 16: 1003-1008.
11. Freedman DS, Mei Z, Srinivasan SR, Berenson GS, Dietz WH (2007) Cardiovascular risk factors and excess adiposity among overweight children and adolescents: the Bogalusa Heart Study. *J Pediatr* 150: 12-17.
12. Kopelman P (2007) Health risks associated with overweight and obesity. *Obes Rev* 8 Suppl 1: 13-17.
13. Li C, Ford ES, Zhao G, Mokdad AH (2009) Prevalence of pre-diabetes and its association with clustering of cardiometabolic risk factors and hyperinsulinemia among U.S. adolescents: National Health and Nutrition Examination Survey 2005-2006. *Diabetes Care* 32: 342-347.
14. (2011) Control of Diabetes and Prevention, National diabetes fact sheet: national estimates and general information on diabetes and prediabetes in the United States, 2011. US Department of Health and Human Services.
15. [No authors listed] (2005) Fight childhood obesity to help prevent diabetes, say WHO & IDF. *Cent Eur J Public Health* 13: 39.
16. Daniels SR, Arnett DK, Eckel RH, Gidding SS, Hayman LL, et al. (2005) Overweight in children and adolescents: pathophysiology, consequences, prevention, and treatment. *Circulation* 111: 1999-2012.
17. Dietz WH (2004) Overweight in childhood and adolescence. *N Engl J Med* 350: 855-857.
18. Adair LS (2008) Child and adolescent obesity: epidemiology and developmental perspectives. *Physiol Behav* 94: 8-16.
19. Al-Ahmadi, M. and H. Al-Hazzaa (2004) Validity of a self-reported questionnaire for youth 15-25 years: Comparison with accelerometer, pedometer and heart rate telemetry. *Saudi Sports Medicine Journal* 7: 2-14.
20. Böhning D (1992) Multinomial logistic regression algorithm. *Annals of the Institute of Statistical Mathematics* 44:197-200.
21. Anderson J (1982) Logistic regression. *Handbook of Statistics*. North-Holland, New York pp: 169-191.
22. Janssen I, Katzmarzyk PT, Boyce WF, King MA, Pickett W (2004) Overweight and obesity in Canadian adolescents and their associations with dietary habits and physical activity patterns. *J Adolesc Health* 35: 360-367.
23. Patrick K, Norman GJ, Calfas KJ, Sallis JF, Zabinski MF, et al. (2004) Diet, physical activity, and sedentary behaviors as risk factors for overweight in adolescence. *Arch Pediatr Adolesc Med* 158: 385-390.
24. Sallis JF, Prochaska JJ, Taylor WC (2000) A review of correlates of physical activity of children and adolescents. *Med Sci Sports Exerc* 32: 963-975.
25. Janssen I, Katzmarzyk PT, Boyce WF, Vereecken C, Mulvihill C, et al. (2005) Comparison of overweight and obesity prevalence in school-aged youth from 34 countries and their relationships with physical activity and dietary patterns. *Obes Rev* 6: 123-132.
26. Al-Hazzaa HM (2013) A Cross-Cultural Comparison of Health Behaviors between Saudi and British Adolescents Living in Urban Areas: Gender by Country Analyses. *International Journal of Environmental Research and Public Health* 10: 6701-6720.
27. White J, R Jago (2012) Prospective Associations Between Physical Activity and Obesity Among Adolescent Girls: Racial Differences and Implications for Prevention. *Physical Activity and Obesity Among Girls*. *Arch Pediatr Adolesc Med* 166: 522-527.
28. (2008) Physical Activity Guidelines Advisory Committee, Physical activity guidelines advisory committee report, 2008. Washington, DC: US Department of Health and Human Services.
29. Tremblay MS, Warburton DE, Janssen I, Paterson DH, Latimer AE, et al. (2011) New Canadian physical activity guidelines. *Appl Physiol Nutr Metab* 36: 36-46.
30. (2004) Department of Health and Ageing, Australia's Physical Activity Recommendations for 5-12 year olds, D.o.H.a. Ageing, Editor.
31. Al-Hazzaa HM (2002) Physical activity, fitness and fatness among Saudi children and adolescents: implications for cardiovascular health. *Saudi Med J* 23: 144-150.
32. Al-Hazzaa HM (2004) Prevalence of physical inactivity in Saudi Arabia: a brief review. *East Mediterr Health J* 10: 663-670.
33. Al-Hazzaa HM (2006) School backpack. How much load do Saudi school boys carry on their shoulders? *Saudi Med J* 27: 1567-1571.
34. NBCNEWS, Saudi Arabia relaxes ban on school sports for girls. 2013.
35. Collison KS, Zaidi MZ, Subhani SN, Al-Rubeaan K, Shoukri M, et al. (2010) Sugar-sweetened carbonated beverage consumption correlates with BMI, waist circumference, and poor dietary choices in school children. *BMC Public Health* 10: 234.

36. Al Sabbah H (2007) Food habits and physical activity patterns among Palestinian adolescents: findings from the national study of Palestinian schoolchildren (HBSC-WBG2004). *Public Health Nutrition* 10: 739-746.
37. Henry CJ, Lightowler HJ, Al-Hourani HM (2004) Physical activity and levels of inactivity in adolescent females ages 11-16 years in the United Arab Emirates. *Am J Hum Biol* 16: 346-353.
38. Al-Rukban MO (2003) Obesity among Saudi male adolescents in Riyadh, Saudi Arabia. *Saudi Med J* 24: 27-33.
39. Mahfouz AA, Abdelmoneim I, Khan MY, Daffalla AA, Diab MM, et al. (2008) Obesity and related behaviors among adolescent school boys in Abha City, Southwestern Saudi Arabia. *J Trop Pediatr* 54: 120-124.
40. Ortega FB, Tresaco B, Ruiz JR, Moreno LA, Martin-Matillas M, et al. (2007) Cardiorespiratory fitness and sedentary activities are associated with adiposity in adolescents. *Obesity (Silver Spring)* 15: 1589-1599.
41. American Academy of Pediatrics Committee on Public Education. (2001) American Academy of Pediatrics: Children, adolescents, and television. *Pediatrics* 107: 423-426.
42. Patriarca A, Di Giuseppe G, Albano L, Marinelli P, Angelillo IF (2009) Use of television, videogames, and computer among children and adolescents in Italy. *BMC Public Health* 9: 139.
43. Loucaides CA, Jago R, Theophanous M (2011) Physical activity and sedentary behaviours in Greek-Cypriot children and adolescents: a cross-sectional study. *Int J Behav Nutr Phys Act* 8: 90.
44. Tammelin T, Ekelund U, Remes J, Näyhä S (2007) Physical activity and sedentary behaviors among Finnish youth. *Med Sci Sports Exerc* 39: 1067-1074.
45. Cui Z, Hardy LL, Dibley MJ, Bauman A (2011) Temporal trends and recent correlates in sedentary behaviours in Chinese children. *Int J Behav Nutr Phys Act* 8: 93.
46. Fulton JE, Wang X, Yore MM, Carlson SA, Galuska DA, et al. (2009) Television viewing, computer use, and BMI among U.S. children and adolescents. *J Phys Act Health* 6 Suppl 1: S28-35.
47. Matheson DM, Killen JD, Wang Y, Varady A, Robinson TN (2004) Children's food consumption during television viewing. *Am J Clin Nutr* 79: 1088-1094.
48. Vereecken CA, Todd J, Roberts C, Mulvihill C, Maes L (2006) Television viewing behaviour and associations with food habits in different countries. *Public Health Nutr* 9: 244-250.