

## Physical Exercise Improves Health Domains in HIV Patients: 'Lifting a Burden'

Trevor Archer\*

Department of Psychology, University of Gothenburg, Box 500, S-405 30 Gothenburg, Sweden

### Editorial

Progressive behavioral and neurological dysfunctions are associated with human immunodeficiency virus type 1 (HIV-1) that involve impairments within several areas of neurocognitive expression [1-3], brain integrity [4-7] and motor and everyday behavioral functioning [8,9]. Due to these functional and cerebral dysregulations the availability of possible therapeutics is easily reconciled with existing demand [10]. Regular physical exercise offers protection against adverse inflammatory, neurobehavioral, epigenetic and metabolic outcomes of trauma with the promise of disease prevention [11-14]. Several of the neurological and immune system disorder states arising from neurodevelopmental dysregulations, toxicity exposures, perinatal 'accidents', infections or inflammatory, neuroimmuno-excitotoxicity 'accidents' may be alleviated through planned schedules of physical exercise and/or activity [15,16], as well as problems arising from the encroachments of immunosenescence [17] with Exercise programs, both clinical, laboratory and health-promotional exercise/activity programs linked to anti-apoptotic, anti-excitotoxic and immune defense-reinforcing influences [18-20] that provide improvements in affective status, neurocognition, quality-of life, motor propensities and physical activity [21]. It has been shown that exercise adherence, involving propensity-to-engage-in and compliance-with, offers a dynamic and fluctuating construct modulated by the episodic nature of HIV and its multi-morbidity aspect among HIV patients [22]. For example, among Nigerian HIV-patients, pulmonary functioning and cardiorespiratory and symptoms of depressiveness were ameliorated [23]. In a longitudinal interventional study of exercise influence upon brain volume (integrity) and cognitive performance of exercise or sedentary HIV-patients [24], it was found that exercising patients produced better executive, but not motor, function, and greater levels of brain integrity. Furthermore, exercise was linked to improved neurocognitive performance and everyday functioning, as estimated by the instrumental activities of daily living in older HIV+ patients [25-27].

Distal symmetrical poly-neuropathy (DSP), associated with two thirds of HIV cases, involves deficits in mobility, quality-of-life and psychological performance [28-31]. Physical exercise interventions have generally given measureable improvements among those domains [32,33]. Not least of all, aerobic exercise, for the most part, has provided enhanced muscle strength and cardiovascular/metabolic improvements independent of specific exercise regimes [34,35]. Mkandla et al. [36] observed that HIV-patients presenting pain, numbness and muscular weakness due to neuropathy experienced improved quality-of-life following exercise interventions; similar improvements were seen with those HIV patients suffering from chronic inflammation, lack of fitness and metabolic vulnerability [37]. A six-week program of peer-led exercise in AmaXhosa HIV patients led to some pain reduction [38] whereas among HIV-patients living in the deep south of the United States, where physical exertion was observed a huge hindrance rendering most participants sedentary, benefits to physical performance were forthcoming [39,40]. Pedro et al. [41] studied parameters of cardiovascular functioning, including heart

rate variability, submaximal values of oxygen uptake, heart rate, peak speed and peak oxygen uptake following sixteen weeks of concurrent aerobic exercise in HIV patients. They found that this exercise schedule was effective in providing improved cardiovascular fitness and higher endurance performance. Similarly, Brown et al. [42] assessed the efficacy of a physiotherapy-led. Group-rehabilitation intervention program involving physical exercise to implement function in HIV patients with regard to referrals, adherence and outcomes; they obtained improvements in the six-minute walk test, strength in triceps and biceps, latisimus dorsi, shoulder-press, chest-press and leg-press, and the physical, emotional and functional subscales, with higher levels of valid expectancies. Finally, the importance of estimating interventional adherence, compliance and propensity to maintain an 'exercise-habit' cannot be sufficiently stressed upon [43-45].

In conclusion, exercise promotes improvements within a multiplicity of parameters of health domains and may well contribute to a significant increase in the well-being of individuals afflicted with HIV. In this respect, the intervention holds sufficient promise of a high level of quality-of-life together with an amelioration of disease burden.

### References

1. Houston E, Lyons T, Wolfe B, Rolfsen N, Williams M, et al. (2016) Assessing Implicit Cognition Among Patients Lost to Follow-up for HIV Care: A Preliminary Study. *Open AIDS J* 10: 83-92.
2. Katrina DH, Katherine FC, Paige LW, Renee S, Kathleen MM, et al. (2016) Contributions of Disease Severity, Psychosocial Factors, and Cognition to Behavioral Functioning in US Youth Perinatally Exposed to HIV. *AIDS Behav*.
3. Spies G, Fennema-Notestine C, Cherner M, Seedat S (2016) Changes in cognitive function in women with HIV infection and early life stress. *AIDS Care* 11: 1-10.
4. Kesby JP, Markou A, Semenova S, TMARC Group (2016a) Effects of HIV/TAT protein expression and chronic selegiline treatment on spatial memory, reversal learning and neurotransmitter levels in mice. *Behav Brain Res* 311: 131-40.
5. Kesby JP, Markou A, Semenova S (2016b) The effects of HIV-1 regulatory TAT protein expression on brain reward function, response to psychostimulants and delay-dependent memory in mice. *Neuropharmacology* 109: 205-15.
6. Soontornniyomkij V, Kesby JP, Morgan EE, Bischoff-Grethe A, Minassian A, et al. (2016) Effects of HIV and Methamphetamine on Brain and Behavior: Evidence from Human Studies and Animal Models. *J Neuroimmune Pharmacol* 3: 495-510.
7. Su T, Wit FW, Caan MW, Schouten J, Prins M, et al. (2016) White matter hyperintensities in relation to cognition in HIV-infected men with sustained suppressed viral load on cART. *AIDS*.

\*Corresponding author: Archer T, Department of Psychology, University of Gothenburg, Box 500, S-405 30 Gothenburg, Sweden, Tel: +46 31 7864694; E-mail: [trevor.archer@psy.gu.se](mailto:trevor.archer@psy.gu.se)

Received August 17, 2016; Accepted August 18, 2016; Published August 25, 2016

Citation: Archer T (2016) Physical Exercise Improves Health Domains in HIV Patients: 'Lifting a Burden'. *HIV Curr Res* 1: e102.

Copyright: © 2016 Archer T. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

8. McGuire JL, Gill AJ, Douglas SD, Kolson DL, CNS HIV Antiretroviral Therapy Effects Research (CHARTER) Group (2016) The complement system, neuronal injury, and cognitive function in horizontally-acquired HIV-infected youth. *J Neurovirol*.
9. Wendelken LA, Jahanshad N, Rosen HJ, Busovaca E, Allen I, et al. (2016) ApoE  $\epsilon$ 4 is Associated with Cognition, Brain Integrity and Atrophy in HIV Over Age 60. *J Acquir Immune Defic Syndr*.
10. Archer T (2016b) Aspects of cognition/health failure by HIV-infected individuals: amelioration through exercise. *HIV: Current Research* 1:1.
11. Cechinel LR, Basso CG, Bertoldi K, Schallenger B, de Meireles LC, et al. (2016) Treadmill exercise induces age and protocol-dependent epigenetic changes in prefrontal cortex of Wistar rats. *Behav Brain Res* 313: 82-87.
12. Kim TK, Lee JE, Kim JE, Park JY, Choi J, et al. (2016) G9a-mediated regulation of OXT and AVP expression in the basolateral amygdala mediates stress-induced lasting behavioral depression and its reversal by exercise. *Mol Neurobiol* 53: 2843-2856.
13. Okudan N, Belvirani M (2016) Effects of exercise training on hepatic oxidative stress and antioxidant status in aged rats. *Arch Physiol Biochem* 16: 1-6.
14. Belviranli M, Okudan N, Revan S, Balci S, Gokbel H (2016) Repeated Supramaximal Exercise-Induced Oxidative Stress: Effect of  $\beta$ -Alanine Plus Creatine Supplementation. *Asian J Sports Med* 7: e26843.
15. Archer T, Garcia D (2015) Exercise and dietary restriction for promotion of neurohealth benefits. *Health* 7: 136-152.
16. Archer T, Garcia D (2016) Exercise influences upon stress-resilience and resilient health. *Kenkyu J Neuropsychol & Stress Managem* 1: 100101.
17. Archer T, Fredriksson A, Schütz E, Kostrzewa RM (2011) Influence of physical exercise on neuroimmunological functioning and health: aging and stress. *Neurotox Res* 20: 69-83.
18. Archer T (2015) Exercise as therapy: health and well-being. *J Intellect Disabil-Diagnos Treatm* 3: 76-81.
19. Archer T (2016a) The influence of physical exercise on well-being and health.
20. Archer T, Kostrzewa RM (2012) Physical exercise alleviates ADHD symptoms: regional deficits and developmental trajectory. *Neurotox Res* 18: 195-209.
21. Archer J, Josefsson T, Lindvall M (2014) Effects of physical exercise on depressive symptoms and biomarkers in depression. *CNS Neurol Disord Drug Targets* 13: 1640-1653.
22. Simonik A, Vader K, Ellis D, Kesbian D, Leung P, et al. (2016) Are you ready? Exploring readiness to engage in exercise among people living with HIV and multimorbidity in Toronto, Canada: a qualitative study. *BMJ Open* 6: e010029.
23. Aweto HA, Aiyegbusi AI, Ugonabo AJ, Adeyemo TA (2016) Effects of Aerobic Exercise on the Pulmonary Functions, Respiratory Symptoms and Psychological Status of People Living With HIV. *J Res Health Sci* 16: 17-21.
24. Ortega M, Baker LM, Vaida F, Paul R, Basco B, et al. (2015) Physical Activity Affects Brain Integrity in HIV+ Individuals. *J Int Neuropsychol Soc* 21: 880-9.
25. Fazeli PL, Marquine MJ, Dufour C, Henry BL, Montoya J, et al. (2015) Physical Activity is Associated with Better Neurocognitive and Everyday Functioning Among Older Adults with HIV Disease. *AIDS Behav* 19: 1470-7.
26. Fazeli PL, Woods SP, Heaton RK, Umlauf A, Gouaux B, et al. (2014) An active lifestyle is associated with better neurocognitive functioning in adults living with HIV infection. *J Neurovirol* 20: 233-42.
27. Dufour CA, Marquine MJ, Fazeli PL, Henry BL, Ellis RJ, et al. (2013) Physical exercise is associated with less neurocognitive impairment among HIV-infected adults. *J Neurovirol* 19: 410-7.
28. Dudgeon WD, Phillips KD, Carson JA, Brewer RB, Durstine JL, et al. (2006) Counteracting muscle wasting in HIV-infected individuals. *HIV Med* 7: 299-310.
29. Jaggars JR, Dudgeon WD, Burgess S, Phillips KD, Blair SN, et al. (2014a) Psychological correlates of HIV-related symptom distress. *J Assoc Nurses AIDS Care* 25: 309-17.
30. Jaggars JR, Prasad VK, Dudgeon WD, Blair SN, Sui X, et al. (2016b) Associations between physical activity and sedentary time on components of metabolic syndrome among adults with HIV. *AIDS Care* 26: 1387-92.
31. Wirth MD, Jaggars JR, Dudgeon WD, Hébert JR, Youngstedt SD, et al. (2016) Association of Markers of Inflammation with Sleep and Physical Activity Among People Living with HIV or AIDS. *AIDS Behav* 19: 1098-107.
32. Dudgeon WD, Jaggars JR, Phillips KD, Durstine JL, Burgess SE, et al. (2012) Moderate-Intensity Exercise Improves Body Composition and Improves Physiological Markers of Stress in HIV-Infected Men. *ISRN AIDS*.
33. Dudgeon WD, Phillips KD, Durstine JL, Burgess SE, Lyerly GW, et al. (2010) Individual exercise sessions alter circulating hormones and cytokines in HIV-infected men. *Appl Physiol Nutr Metab* 35: 560-8.
34. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, et al. (2011) American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 43: 1334-59.
35. O'Brien KK, Tynan AM, Nixon SA, Glazier RH (2016) Effectiveness of aerobic exercise for adults living with HIV: systematic review and meta-analysis using the Cochrane Collaboration protocol. *BMC Infect Dis* 16: 182.
36. Mkandla K, Myezwa H, Musenge E (2016) The effects of progressive-resisted exercises on muscle strength and health-related quality of life in persons with HIV-related poly-neuropathy in Zimbabwe. *AIDS Care* 28: 639-43.
37. Cutrono SE, Lewis JE, Perry A, Signorile J, Tiozzo E, et al. (2016) The Effect of a Community-Based Exercise Program on Inflammation, Metabolic Risk, and Fitness Levels Among Persons Living with HIV/AIDS. *AIDS Behav* 20: 1123-31.
38. Parker R, Jelsma J, Stein DJ (2016) Managing Pain in Women Living With HIV/AIDS: A Randomized Controlled Trial Testing the Effect of a Six-Week Peer-Led Exercise and Education Intervention. *J Nerv Ment Dis*.
39. Rehm KE, Konkle-Parker D (2016a) Physical Activity Levels And Perceived Exercise Benefits And Barriers In HIV+ Women Living In Mississippi: 2527 Board #50 June 3, 11: 00 AM - 12: 30 PM. *Med Sci Sports Exerc* 48(5 Suppl 1): 698.
40. Rehm KE, Konkle-Parker D (2016b) Physical activity levels and perceived benefits and barriers to physical activity in HIV-infected women living in the deep south of the United States. *AIDS Care* 28: 1205-10.
41. Pedro RE, Guariglia DA, Okuno NM, Deminice R, Peres SB, et al. (2016) Effects Of 16 Weeks Of Concurrent Training On Resting Heart Rate Variability And Cardiorespiratory Fitness In People Living With HIV/AIDS Using Antiretroviral Therapy: A Randomized Clinical Trial. *J Strength Cond Res*.
42. Brown D, Claffey A, Harding R (2016) Evaluation of a physiotherapy-led group rehabilitation intervention for adults living with HIV: referrals, adherence and outcomes. *AIDS Care* 5: 1-11.
43. Archiropoli A, Ginossar T, Wilcox B, Avila M, Hill R, et al. (2016) Factors of interpersonal communication and behavioral health on medication self-efficacy and medication adherence. *AIDS Care* 20: 1-8.
44. Garcia D, Medeiros da Costa Daniele T, Archer (2016) A brief measure to predict exercise behavior: the Archer-Garcia ratio.
45. Peres SB, Guariglia DA, Pedro RE, Candido N, Melo BP, et al. (2016) Effects Of Concurrent Training In People Living With HIV/aids: A Randomized Clinical Trial: 2572 Board #95 June 3, 9: 30 AM - 11: 00 AM. *Med Sci Sports Exerc* 48(5 Suppl 1): 713.

**Citation:** Archer T (2016) Physical Exercise Improves Health Domains in HIV Patients: 'Lifting a Burden'. *HIV Curr Res* 1: e102.

### OMICS International: Publication Benefits & Features

#### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

#### Special features:

- 700+ Open Access Journals
- 50,000+ Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: [www.omicsonline.org/submission/](http://www.omicsonline.org/submission/)