

Pigments and Pathogenesis

Beatrice Saviola*

Basic Medical Sciences, College of Osteopathic Medicine, Western University of Health Sciences, USA

*Corresponding author: Beatrice Saviola, Basic Medical Sciences, College of Osteopathic Medicine, Western University of Health Sciences, 309 E. Second St. Pomona CA 91766, USA, Tel: +1 909 469-5373; Fax: +1 909 469-5698; E-mail: bsaviola@westernu.edu

Rec date: April 11, 2014, Acc date: September 25, 2014, Pub date: October 10, 2014

Copyright: © 2014 Saviola B. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Abstract

In recent years pigments have been identified in human nutrition to have a positive effect on human health and reduction to oxidative stress exposure. In the media it has become common wisdom that colourful food is naturally better to consume for humans and animals. Now recently it has been shown that pigments aid microbial species as well, and conversely these microbial pigments may result in more morbidity and mortality for the human host infected by these colourful microbes. Similar pigments that are available for consumption in food are also present in many bacterial species. Presumably these pigments aid the bacteria in their survival in the environment and within a human or animal host. Importantly, interference with the production of certain microbial pigments results in some bacterial strains that are more susceptible to environmental stressors and the host immune system. These studies seem to indicate a role of pigments for in vivo survival by microbial species.

Keywords: Bacterial pathogenesis, Carotenoid, Mycobacteria, Pigment

Pigments have evolved in nature to serve a variety of purposes. In higher animals colour can signal sexual fitness as brightly coloured birds attest to [1]. But pigments can serve a more fundamental purpose that includes protection from oxidative stress, absorption of light in plants and the retina in animals, and protection from harmful UV light [1]. The appearance to have colour may be incidental to the chemistry of these pigments which contain varying amounts of conjugated double bonds [1]. These double bonds serve to absorb light which leads to the phenomenon of colour. In addition, conjugated double bonds are efficient antioxidants, reducing oxidative stress inside cells that can result from UV light exposure or other stresses [2]. For bacteria where there are no visualization systems, pigments functioning in other capacities is only logical [1].

Carotenoids may have evolved to reinforce bacterial membranes. Conjugated double bonds provide rigidity, that when inserted into a membrane serve to stabilize it. Carotenoids seem to have evolved within archaeobacteria serving mechanical functions [1]. In addition carotenoids present within the ecto-skeleton of crustaceans and similar animals serve to protect and reinforce the various proteins present within these structures. Carotenoids complexed with proteins in certain ova can also prevent protein digestion from native proteases [1].

In pathogenic bacteria pigments have been implicated in bacterial survival and virulence. The gram positive pathogen *Staphylococcus aureus* is named such because it constitutively produces a golden yellow pigment [3]. When genes encoding synthetic enzymes responsible for the production of staphyloxanthin are deleted, the bacterium appears colourless. More significantly these mutant bacteria are more susceptible to killing via host immune cells, presumably due to less resistance to reactive oxygen species produced within the phagosomes of these cells [4]. In addition these mutant strains are

much less virulent in a mouse model of skin abscess or during disseminated infections. In a study where the genes responsible for carotenoid production were transferred to a group *A streptococcus*, this new recombinant streptococcal strain was both pigmented and produced enlarged skin lesions in a mouse model of infection [5]. A cholesterol synthesis inhibitor was revealed to inhibit carotenoid synthesis in *S. aureus* when placed in the growth media of the microorganism [6]. The bacteria grown in the presence of the cholesterol synthesis inhibitor also had increased sensitivity to immune cell killing and reactive oxygen species [6].

While some species of bacteria produce pigment constitutively, others produce pigment in response to environmental stresses. *Synechococcus* species accumulate carotenoids in response to low iron stress [7]. Methanol exposure can induce pigment production in *Acinetobacter wofii* [8]. Oxidative stress may possibly be increased due to methanol exposure and carotenoids may decrease the effects of this stress. *Vibrio cholera* makes melanin under hyperosmotic stress as well as acidity [9,10]. This response of increased melanin may aid survival in aquatic environments as well as within the upper gastrointestinal tracts of infected humans. In addition, pigments have been shown to combat mutagenesis and act as scavengers of reactive oxygen species thus protecting DNA [11]. Norbixin is a carotenoid from seeds of a tropical shrub from Brazil. This pigment supplied to *Escherichia coli* protects these microbes from damage due to UV light exposure as well as reactive oxygen species [2].

The genus *Mycobacterium* includes many bacteria that produce pigments [12-14]. The Runyon classification system groups mycobacteria based on growth rate and pigment production [15]. Carotenoid pigment production in mycobacteria was previously thought to only be produced constitutively or in response to UV light exposure [16]. Interestingly, recently it has been shown that mycobacteria also produce pigment in response to acidic stress at pHs 5.0-6.0 [17]. The pigment has a similar absorption profile to other carotenoids and *Mycobacterium smegmatis* carotenoid production genes were upregulated by acidic stress at pH 5.5 and 6.0 [17]. This

pigment production may protect mycobacteria from acidic stress or concurrent oxidative stress found in the environment or in the host.

Some species of mycobacteria produce pigment constitutively and are designated as scotochromogens [18]. Other mycobacteria such as *M. smegmatis* and *Mycobacterium goodii* produce pigment only after extended growth on agar plates of 7-10 days [15,19]. These species are considered to be late pigment producers. Recently both these species have been shown to produce pigment at 3 days growth on agar plates at pH 5.5 and 6.0 but not 7.0 (Figure 1). Perhaps the late formation of the pigment is actually in response to an increase in acidity in the agar media due to extended incubation and metabolism by the mycobacterial cells [17]. As the mycobacteria metabolize sugars present within the media, they also produce excess protons which cause the media to acidify. Other rapid growing mycobacteria were also shown to produce pigment in response to acidic pH at 5.5-6.0, including *Mycobacterium abscessus*, *Mycobacterium fortuitum* and *Mycobacterium chelonae* [17]. *M. chelonae* and *M. abscessus* were also exposed to pH 5.0, however only *M. abscessus* continued to produce the pigment and *M. chelonae* ceased production of the pigment possibly due to the toxic effects of the more extreme pH on pigment production [17]. Several slow growing species such as *Mycobacterium avium* intracellulare and *Mycobacterium avium* also produce pigment in response to acidic stress at pH 5.5-6.0 [17,20, 21]. As these environmental mycobacterial species may encounter acidity in stagnant water in bogs and swamps, rivers, streams, brooks, and certain soils, pigment production may protect mycobacteria against environmental stress. Pigment production may aid in the resistance to acidity and help these mycobacteria survive.



Figure 1: *M. smegmatis* were patched onto 7H10 agar media containing ADC (albumen dextrose, and sodium chloride) at pH 5.5 (left) or pH 7.0 (right) and were incubated for 72 hours at 37°C. The bacteria on the left plate appear to contain a bold o yellow colour

The promoter driving expression of carotenoid synthesis genes has regions that respond transcriptionally to exposure to light in *Mycobacterium marinum* [22,23]. In *M. smegmatis*, exposure to acidity transcriptionally upregulates the homologue of this promoter region [17]. In addition, at least for *M. smegmatis*, pigment production is thought to be controlled by the sigma factor SigF [24,25]. In *M. smegmatis* a deletion in the sigF gene resulted in a bacterium which not only cannot produce pigments, but a mutant

mycobacterium sensitive to oxidative stress as well [24,26]. Likely the presence or absence of carotenoid compounds dictates resistance to oxidative stress. In addition, lack of sigF increases transformation efficiency indicating the cell envelope has become potentially more porous [24]. Possibly carotenoid compounds lead to stabilization of the cell envelope, as well as making this structure less traversable. With the lack of stabilizing carotenoids, DNA may be able to travel through the cell wall/membrane more easily. SigF may also regulate other cell wall/membrane components that are altered within the mutant mycobacterium, and it remains to be determined what contribution carotenoid compounds make to cell wall/membrane stability. In *Mycobacterium tuberculosis* sigF is thought to control genes that aid in persistence and survival in late stage disease and a deletion in the sigF gene results in decreased mycobacterial burden *in vivo* [27].

Many mycobacteria including *M. tuberculosis* inhabit the phagosomes of macrophages [28-30]. When rapid and slow growing environmental mycobacteria enter the human body they are phagocytosed by macrophages where the mycobacteria often reside within the phagosomes. Resistant individuals may quickly eliminate these mycobacteria via many antibacterial mechanisms which include an increase in acidity to pH 5.0 and possibly rebounding to pH 6.0-6.5 [30]. In addition reactive oxygen intermediates are produced that also can damage these mycobacteria. In susceptible individuals these antibacterial mechanisms may be weakened and resistance mechanisms possessed by the environmental mycobacteria may allow for the successful replication of the pathogens *in vivo*. Pre-priming via environmental acidity to induce carotenoid compound induction may increase resistance to acidity *in vivo*, as well as resistance to oxidative stress. In fact, recently when carotenoid production was inhibited in *M. smegmatis* by a sigF disruption, it did increase sensitivity of these mycobacteria to oxidative stress [24,26]. Carotenoids are well known to detoxify oxidative stress due to their double bonds [1]. In addition carotenoids may stabilize the cellular membrane/cell wall of mycobacteria to decrease penetration of all types of environmental stresses including acidity and reactive oxygen intermediates [1].

Previously *Mycobacterium tuberculosis* was found to produce a pigment in response to para-aminobenzoic acid (PABA), however this was thought to be a metabolite of PABA [31]. Another pigment was also observed in response to para-aminosalicylic acid exposure [32]. In addition, *M. tuberculosis* seems to produce a non-carotenoid pigment that is cell wall associated, and this is produced due to long term growth in anaerobic culture [33]. These pigments may increase survivability under stressful conditions of low oxygen by serving to stabilize the outer membrane and the cell wall. It remains to be seen if *M. tuberculosis* also produces carotenoid pigments in response to acidic stress.

Pigment production in bacterial pathogens may increase their virulence. These pigments have been shown to increase resistance to oxidative stress, killing by immune cells, and mutagenesis. Pigments seem to increase virulence of pathogens by increasing invasiveness, survival in immune cells, and size of local abscesses. In fact in *M. marinum* pigmentation is linked to a locus important in intracellular survival [22]. Thus the finding that pigments in environmental mycobacteria are produced under conditions of acidity which can be found in the phagosomes of macrophages or the centers of caseating granulomas indicate that pigments may be important for pathogenesis [17]. However it is well known that mycobacteria are inherently resistant to a variety of environmental insults and stressors due a thick and waxy cell wall and possibly other genetic factors. Pigments may

augment this resistance in certain circumstances possibly to increase survivability in the face of microbe-external insults. In the future we may be able to inhibit environmental mycobacterial growth in vivo by inhibiting pigment production such as has been shown for *S. aureus* [6].

References

- Vershinin A (1999) Biological functions of carotenoids--diversity and evolution. *Biofactors* 10: 99-104.
- Júnior AC, Asad LM, Oliveira EB, Kovary K, Asad NR, et al. (2005) Antigenotoxic and antimutagenic potential of an annatto pigment (norbixin) against oxidative stress. *Genet Mol Res* 4: 94-99.
- Lan L, Cheng A, Dunman PM, Missiakas D, He C (2010) Golden pigment production and virulence gene expression are affected by metabolisms in *Staphylococcus aureus*. *J Bacteriol* 192: 3068-3077.
- Liu GY, Essex A, Buchanan JT, Datta V, Hoffman HM, et al. (2005) *Staphylococcus aureus* Golden Pigment Impairs Neutrophil Killing and Promotes Virulence Through its Antioxidant Activity. *J. Exp. Med.* 202: 209-215.
- Singer AJ, Talan DA (2014) Management of skin abscesses in the era of methicillin-resistant *Staphylococcus aureus*. *N Engl J Med* 370: 1039-1047.
- Liu CI, Liu GY, Song Y, Yin F, Hensler ME, et al. (2008) A cholesterol biosynthesis inhibitor blocks *Staphylococcus aureus* virulence. *Science* 319: 1391-1394.
- Ivanov AG, Krol M, Selstam E, Sane PV, Sveshnikov D, et al. (2007) The induction of CP43' by iron-stress in *Synechococcus* sp. PCC 7942 is associated with carotenoid accumulation and enhanced fatty acid unsaturation. *Biochim Biophys Acta* 1767: 807-813.
- Ghosh A, Goyal A, Jain RK (2007) Study of methanol-induced phenotypic changes in a novel strain of *Acinetobacter lwoffii*. *Arch Microbiol* 188: 533-539.
- Liu GY, Nizet V (2009) Color me bad: microbial pigments as virulence factors. *Trends Microbiol* 17: 406-413.
- Coyne VE, al-Harhi L (1992) Induction of melanin biosynthesis in *Vibrio cholerae*. *Appl Environ Microbiol* 58: 2861-2865.
- Geng J, Yu SB, Wan X, Wang XJ, Shen P, et al. (2008) Protective action of bacterial melanin against DNA damage in full UV spectrums by a sensitive plasmid-based noncellular system. *J Biochem Biophys Methods* 70: 1151-1155.
- Baker JA (1938) Light As A Factor In The Production Of Pigment By Certain Bacteria. *J Bacteriol* 35: 625-631.
- Batra PP, Rilling HC (1964) On The Mechanism Of Photoinduced Carotenoid Synthesis: Aspects Of The Photoinductive Reaction. *Arch Biochem Biophys* 107: 485-492.
- Eisenstadt J, Hall GS (1995) Microbiology and classification of mycobacteria. *Clin Dermatol* 13: 197-206.
- Brown-Elliott BA, Wallace RJ Jr (2002) Clinical and taxonomic status of pathogenic nonpigmented or late-pigmenting rapidly growing mycobacteria. *Clin Microbiol Rev* 15: 716-746.
- Rilling HC (1962) Photoinduction of carotenoid synthesis of a *Mycobacterium* sp. *Biochim Biophys Acta* 60: 548-556.
- Saviola B, Felton J (2011) Acidochromogenicity is a common characteristic in nontuberculous mycobacteria. *BMC Res Notes* 4: 466.
- Ichiyama S, Shimokata K, Tsukamura M (1988) Relationship between mycobacterial species and their carotenoid pigments. *Microbiol Immunol* 32: 473-479.
- Brown BA, Springer VA, Steingrube RW, Wilson GE, Pfyffer MJ, et al. (1999) *Mycobacterium wolinski* sp. nov. and *Mycobacterium goodii* sp. nov. Two New Rapidly Growing Species Related to *Mycobacterium smegmatis* and Associated with Human Wound Infections: a Cooperative Study from the International Working Group on Mycobacterial Taxonomy. *Int J Syst Bacteriol* 49: 1493-1511.
- Stevenson K, Hughes VM, de Juan L, Inglis NF, Wright F, et al. (2002) Molecular characterization of pigmented and nonpigmented isolates of *Mycobacterium avium* subsp. *paratuberculosis*. *J Clin Microbiol* 40: 1798-1804.
- Stormer RS, Falkinham JO 3rd (1989) Differences in antimicrobial susceptibility of pigmented and unpigmented colonial variants of *Mycobacterium avium*. *J Clin Microbiol* 27: 2459-2465.
- Gao L, Groger R, Cox JS, Beverley SM, Lawson EH, et al. (2003) Transposon Mutagenesis of *Mycobacterium marinum* Identifies a Locus Linking Pigmentation and Intracellular Survival. *Infect. And Immun.* 71: 922-929.
- Ramakrishnan L, Tran HT, Federspiel NA, Falkow S (1997) A crtB homolog essential for photochromogenicity in *Mycobacterium marinum*: isolation, characterization, and gene disruption via homologous recombination. *J Bacteriol* 179: 5862-5868.
- Provvedi R, Kocincova D, Dona V, Euphrasie D, Daffe D, et al. (2008) SigF controls Carotenoid Pigment Production and Affects Transformation Efficiency and Hydrogen Peroxide Sensitivity in *Mycobacterium smegmatis*. *J. Bact.* 190: 7859-7863.
- Gebhard S, Hümpel A, McLellan AD, Cook GM (2008) The alternative sigma factor SigF of *Mycobacterium smegmatis* is required for survival of heat shock, acidic pH and oxidative stress. *Microbiology* 154: 2786-2795.
- Wu H, Hu X, Xiao J, Zhang J, Tao J, et al. (2012) [Sigma factor F regulates *Mycobacterium smegmatis* hydrogen peroxide resistance]. *Wei Sheng Wu Xue Bao* 52: 1352-1359.
- Geiman DE, Kaushal D, Ko C, Tyagi S, Manabe YC, et al. (2004) Attenuation of late-stage disease in mice infected by the *Mycobacterium tuberculosis* mutant lacking the SigF alternate sigma factor and identification of SigF-dependent genes by microarray analysis. *Infection and Immunity.* 72: 1733-1745.
- Fisher MA, Plikaytis BB, Scinnick TM (2002) Microarray of the *Mycobacterium tuberculosis* Transcriptional Response to the Acidic Conditions Found in Phagosomes. *J. of Bact* 184: 4025-4032.
- Rohde KH, Abramovitch RB, Russell DG (2007) *Mycobacterium tuberculosis* invasion of macrophages: linking bacterial gene expression to environmental cues. *Cell Host Microbe* 2: 352-364.
- MacMicking JD, Taylor GA, McKinney JD (2003) Immune control of tuberculosis by IFN-gamma-inducible LRG-47. *Science* 302: 654-659.
- Mayer RL (1944) A Yellow Pigment Formed from p-Aminobenzoic Acid by *Mycobacterium tuberculosis* var. *hominis*. *J Bacteriol* 48: 337-345.
- Mayer RL, Crane C (1950) A yellow pigment formed from para-aminobenzoic acid and para-aminosalicylic acid by various strains of *Mycobacterium*. *Nature* 165: 37.
- Cunningham AF, Ashton PR, Spreadbury CL, Lammas DA, Craddock R, et al. (2004) Tubercle bacilli generate a novel cell wall-associated pigment after long-term anaerobic culture. *FEMS Microbiol Lett* 235: 191-198.