

Pilot Test of Elemental Fingerprint in Otolith Core as a Natural Biological Tag to Discriminate among *Gymnocypris przewalskii* Stocks in Lake Qinghai in Qinghai-Tibet Plateau, China

Zhang JY^{1*}, Liu XH¹, Zhao YL¹, Wei FL² and Li CZ^{2*}

¹State Key Laboratory of Freshwater Ecology and Biotechnology, Institute of Hydrobiology, Chinese Academy of Sciences, China

²State Key Laboratory of Plateau Ecology and Agriculture College of Ecological and Environmental Engineering, Qinghai University, China

Abstract

Naked or scaleless carp, *Gymnocypris przewalskii* as reported by Kessler in 1876 is the unique economic fish species and play an important ecological function to maintain the health ecosystem of Qinghai Lake watershed. The natural resource of naked carp, however, has sharply decreased during the past decades. Understanding the stock structure of this endangered fish is prerequisite to make a scientific management strategy to restore its stock resources. Otolith fingerprint has been widely applied to identify and discriminate fish stocks. In the present work, a pilot test was performed to evaluate the possibility of applying trace element fingerprint of otolith core as a natural tag to discriminate among stocks of two inflowing river and releasing artificially hatched and nursed individuals of naked carp. Results showed that otolith core elemental fingerprinting can successfully discriminate the three analyzed naked carp groups which will provide a robust tool for conservation and future development of aquaculture industry of this economic fish in Qinghai-Tibet plateau.

Keywords: *Gymnocypris przewalskii*; Otolith fingerprint; Stock discrimination; Qinghai-Tibet plateau

Introduction

Lake Qinghai is the largest inland saltwater lake, locating in the Qinghai-Tibet Plateau of northwestern China at an elevation of 3200. Extreme conditions, including high pH, high salinity and harsh climate make fish species very few. Naked or scaleless carp, *Gymnocypris przewalskii* was reported by Kessler in 1876 (Cyprinidae: Schizothoracinae) is the unique endemic species which is of most economically and ecologically importance in the Lake Qinghai ecosystem [1,2]. The naked carp resource, however, has sharply decreased from about 200,000 tons in the 1960s to less than 2,600 tons in the 2002 [3] for global warming and anthropogenic interruption, including overfishing, reduction of hatchery habitat by construction of weirs, irrigation diversions and inflowing rivers running dry and ecological deterioration [2]. *G. przewalskii* has become an endangered fish species and been listed in China's Red list since 2004 [4]. The depressed status of naked carp not only seriously threatens the ecological integrity of symbiotic relationship of fish and birds in Lake Qinghai, but also affects the uptake of fish protein for local residents in Qinghai province. To protect and restore the resources of naked carp, 5 consecutive forbidden fishing periods of 10 years since 1982 and stock enhancement and releasing since 2002 were implemented by the local government [3,5]. Recent resource survey by hydroacoustic survey showed the overall fish biomass of naked carp has restored to about 30,000 tons, but far less than original resource [3,6]. Further measures should be taken to support the sustainable increasing of naked carp population. Full understanding the stock structure and dynamics of existed naked carp in watershed of Qinghai Lake is prerequisite to make a holistic scientific and reasonable fishery management strategy.

G. przewalskii is an anadromous species which seasonally migrates to inflowing rivers for reproduction from April to July and then return to the lake for growth [7]. Previous studies showed that there were significant genetic differences among naturally produced stocks in different tributaries of Lake Qinghai and suggested that different groups should be treated as separate management units [5,8]. As an alternative and synergistic tool with genetic approaches, otolith

fingerprint has been widely applied to identify and discriminate fish stocks which is based on the assumption of variation in otolith chemical composition as the proxy of water environment experienced by fish stocks [9,10]. Previous report of remarkable differences among water chemical characteristics of the adjoining rivers of Lake Qinghai is the basis of applying otolith microchemistry to discriminate among stocks of naked carp [11]. So, one of purpose of the present work is to test the possibility of applying the elemental composition in the otolith core of naked carp to discriminate among stocks with different natal and nursery sites. Additionally, assessment of the actual contribution of enhancement and releasing activities to the resource restoration of naked carp has not yet been conducted, although this program has been carried out more than 10 years. Taking into account the different water environment between artificially hatchery and nursery and wild water environment, it is possible, we thought, to apply the elemental signatures of otolith core as a natural biological tag to discriminate the artificially releasing fish from naturally produced fish.

As a pilot study, the trace elemental signatures of otolith core of two naturally produced stocks and the artificially spawn and nursed stock were analyzed by laser ablation inductively coupled plasma mass spectrometry (LA-ICPMS) to test this possibility here.

***Corresponding authors:** JY Zhang, State Key Laboratory of Freshwater Ecology and Biotechnology, Institute of Hydrobiology, Chinese Academy of Sciences, 430072 Wuhan, China, Tel: 86-27-68780007; Fax: 86-27-687800720; E-mail: zhangjy@ihb.ac.cn

*CZ Li, State Key Laboratory of Plateau Ecology and Agriculture College of Ecological and Environmental Engineering, Qinghai University, 810016 Xining, China, Tel: 86-971- 5313741; Fax: 86-971- 5313741; E-mail: licz05@gmail.com

Received May 13, 2016; **Accepted** September 19, 2016; **Published** September 21, 2016

Citation: Zhang JY, Liu XH, Zhao YL, Wei FL, Li CZ (2016) Pilot Test of Elemental Fingerprint in Otolith Core as a Natural Biological Tag to Discriminate among *Gymnocypris przewalskii* Stocks in Lake Qinghai in Qinghai-Tibet Plateau, China. J Aquac Res Development 7: 447. doi: 10.4172/2155-9546.1000447

Copyright: © 2016 Zhang JY, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Material and Methods

Sample collection

Ten samples of naturally hatched naked carp from Quanji River (QJ) and Shaliu River (SL), two adjacent tributaries of Lake Qinghai were respectively collected with cast net at the peak of the hatching season on June 1 of 2014. Ten artificially hatched and nursed fish fingerlings (AP) which were hatched in a hatchery station at the estuary of SL in May of 2013 and transported to the earth pond for nursery in the Rescue Center of Naked carp in Xining city, the province capital of Qinghai for rearing were also captured by trawl net on July 1, 2015. The geographic co-ordinates of sampling site in QJ and SL can be seen in Figure 1, similar with those of O'Bryan et al. [8]. All artificially fertilized eggs of naked carp obtained from stocks of different inflowing rivers were transported to the artificially hatchery station locating at the estuary of SL within 3 h after insemination for hatching. The Rescue Center of Naked carp is about 135 km away from Lake Qinghai (please refer the Figure 1 for the geographical information). All sampled fish were frozen immediately after measuring the body weight (BW) and body length (BL) and transported to the laboratory of Institute of Hydrobiology, Chinese Academy of Sciences in Wuhan city in central China for elemental analysis.

Otolith preparation and elemental analysis

The otoliths extraction was performed according to the previous report [12,13]. Briefly, the sagittal otoliths were removed with acid-washed (0.5% Optima HNO₃, Fisher Inc.) non-metallic forceps, cleaned by sonicating in distilled Milli Q water for 10 min and then oven-dried overnight at 35°C after defrosting, followed by preserving in dry, acid-washed polyethylene boxes until elemental analysis. The longest and shortest axes of the decontaminated otolith were measured to the nearest 1 µm under light microscopy (Olympus BX 35) and their weight to the nearest 0.1 mg using an electrical balance (Sartorius BS-124 S). The left otolith of each pair was used for elemental analysis by LA-ICPMS while the right for age determination based on methods of Chen et al. [14]. All fish were sampled after getting the approval of local and national authorities and all procedures were carried out in accordance with animal care regulations issued by Institute of Hydrobiology, Chinese Academy of Sciences. For LA-ICPMS analysis, concentrations of otolith core elements (including Li, Na, Mg, Al, K, Ca, Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn, Sr, Cd, Ba) were determined on an NW213esi laser ablation system (New Wave Research) coupled with an Agilent 7500ce ICPMS (Agilent Technologies, Tokyo). The laser was operated at a wavelength of 213nm with a pulse rate of 10 Hz and energy density of 9.68Jcm⁻². Data were collected by the default Agilent 7,500s ICPMS software and each element presented

as a ratio to Ca, corrected for mass bias using calibration standards with known element to Ca ratio. To determine the limits of detection (LOD), blank ablations, only consisting of measuring Ar and He gases were done before and after each analysis session for approximately 120 s and calculated from three times the standard deviation of this session. Relative standard deviations (%RSD) based on replicated measurements of the calibration standard, were calculated to reflect the level of precision achieved for each element.

The location of the otolith cores for laser ablating was followed by Dou et al. [15] and ablation spot size was 80 µm and a scan speed 30 µm·s⁻¹. Solid glass standard material from the National Institute of Standards and Technology (NIST 612) was analyzed after every 5 randomized samples besides at the beginning and the end of each analysis session to avoid the possible instrument drift. Data were collected by the default Agilent 7,500s ICPMS software and the concentration of each element was standardized to element: Ca.

Data analysis

For K and Na were strongly physically regulated, they were excluded for subsequent statistical analysis. All detected elemental concentration ratios above the LOD were firstly examined for within-group normality and then ln-transformed if required and also homogeneity of variances among groups. One-way ANOVA was applied to analyze the univariate differences in concentration ratios of each element among three stocks. The overall differences in the elemental fingerprint among all three groups were tested by MANOVA. Stepwise discriminate function analysis (DFA) was used to test the possibility to discriminate among the naked carp stocks by multi-elements variation in otolith core and identify the element: Ca ratio which contributes the most to fingerprint differences among stocks by plotting the first two discriminate function axes against each other. Statistical analyses were performed on SPSS 22.0 for windows (SPSS Inc.) at a significance level of p<0.05.

Results

During otolith preparation, some otolith were destroyed and excluded for elemental analysis. The analyzed sample number of SL, QJ and AP was 7, 6 and 7, respectively. The information of naked carp of three stocks for fingerprint analysis of otolith core can be seen in Table 1. The average BL and BW is 182.4 mm and 64.4 g, 189.3 mm, 67.1 g and 77.1 mm and 6.2 g for SL, QJ and AP stock, respectively. Beside Al, Ti, V, Cr and Cd, the concentration of all analyzed element were significant higher than the corresponding LOD. The analytical accuracy for the NIST standard averaged across all samples was high for the remained elements with %RSD ranged from 7.86 (⁷Li), 3.92 (²⁴Mg), 5.03 (⁵⁵Mn), 7.84 (⁵⁷Fe), 4.91 (⁵⁹Co), 9.30 (⁶⁰Ni), 4.76 (³Cu), 6.80 (⁶⁶Zn), 5.09 (⁸⁸Sr), 6.47 (¹³⁸Ba). The applied element: Ca ratios for statistical analysis fit well the assumptions of within-groups normality (Kolmogorov-Smirnov's test, p>0.05) and homogeneity of variances among groups (Levene's test, p>0.05). One-way ANOVA showed that Li: Ca, Mn: Ca, Sr: Ca and Ba: Ca differed significantly among SL, QJ and AP samples (Table 2). MANOVA also showed significant differences in multi-element fingerprints between all three sample groups (p<0.01). DFA can effectively discriminate SL, QJ and AP sample groups with classification accuracy rate of 100% (Figure 2) and Li: Ca and Ba: Ca contributed most to group separation by F-to-remove analysis.

Discussion

Otolith chemistry has been widely used to address a variety of fishery ecological and conservation issues, including tracking movements in individual fish, discriminating stocks and assessing connectivity




Figure 1: Map of the Qinghai Lake watershed, showing the sampling sites, QL (Quanji River), SL (Shaliu River) and AP (Artificially nursed ponds in Xining city).

No.	Fish samples				Otolith			Sample site
	BL(mm)	BW(g)	Sex	Age	W(mg)	D _{max} (μ m)	D _{min} (μ m)	
1	185	61.2	♀	6	6.8	3583	2890	SL
2	196	87.7	♂	9	8.1	3597	2931	SL
3	176	55.6	♂	6	6.7	3579	2878	SL
4	188	73.1	♀	8	7.8	3588	2899	SL
5	193	61.5	♀	8	8.1	3596	2906	SL
6	164	52.6	♂	7	6.7	3519	2769	SL
7	175	59.0	♀	6	6.9	3579	2896	SL
8	192	64.5	♀	9	9.7	3613	2987	QJ
9	192	68.3	♂	9	9.8	3672	2991	QJ
10	190	66.7	♀	7	9.8	3671	2979	QJ
11	182	60.7	♂	7	9.5	3598	2889	QJ
12	200	84.5	♂	9	9.9	3710	3010	QJ
13	180	58.1	♀	6	9.3	3561	2933	QJ
14	79	4.7	-	1	0.88	1855	1432	AP
15	90	9.5	-	1	0.99	1896	1513	AP
16	64	3.2	-	1	0.82	1784	1416	AP
17	83	7.6	-	1	0.87	1897	1469	AP
18	100	12.2	-	1	0.86	1923	1537	AP
19	72	4.3	-	1	0.79	1826	1398	AP
20	52	1.7	-	1	0.62	1769	1313	AP


Table 1: Information of sampled naked carp from two inflowing rivers (Shaliu river, SL and Quanji River, QJ) and an artificially bred freshwater earth pond (AP) for element analysis using LA-ICPMS (BL: Body Length; BW: Body Weight; W: Otolith Weight; D_{max}: Major Axis Diameter; D_{min}: Minor Axis Diameter).

Element: Ca (mmol mol ⁻¹)		Sum of Squares	df	MS	F _{1,20}	P
Li: Ca	Between Groups	0.07	2	0.04	34.233	0
	Within Groups	2.195	2	1.098		
Mg: Ca	Between Groups	0	2	0	5.844	0.012
	Within Groups	0	17	0		
	Total	0	19			
Fe: Ca	Between Groups	0.001	2	0	0.246	0.785
	Within Groups	0.027	17	0.002		
	Total	0.028	19			
Co: Ca	Between Groups	0	2	0	0.413	0.668
	Within Groups	0	17	0		
	Total	0	19			
Ni: Ca	Between Groups	0	2	0	0.187	0.831
	Within Groups	0.001	17	0		
	Total	0.001	19			
Cu: Ca	Between Groups	0	2	0	1.195	0.327
	Within Groups	0	17	0		
	Total	0	19			
Zn: Ca	Between Groups	0.001	2	0.001	0.859	0.441
	Within Groups	0.011	17	0.001		
	Total	0.012	19			
Sr: Ca	Between Groups	0.032	2	0.016	5.725	0.013
	Within Groups	0.048	17	0.003		
	Total	0.08	19			
Ba: Ca	Between Groups	0	2	0	7.522	0.005
	Within Groups	0	17	0		
	Total	0	19			

Table 2: Analysis of variance (ANOVA) results of element: Ca concentration ratios in otolith core regions of naked carp collected from Shaliu River (SL), Quanji River (QJ) and artificially pond (AP).

between nursery grounds and adults stocks [9,10]. However, no application of otolith fingerprint was reported as the natural biological tag to discriminate stocks of naked carp, an endangered unique fish in Qinghai-Tibet Plateau, although EMPA (electron probe microanalysis) was used to explore its migration history [12,13]. Naked carp is a migratory fish between inflowing river for reproduction and Lake

Qinghai for grow-out [7]. But whether naked carp are characterized by natal homing and rate of exchangeability of hatching populations of different inflowing rivers remains unknown. Understanding the spatial structure of fish stocks and the connectivity between them has been increasingly considered a crucial requirement for sustainable fisheries management [15,16]. So, the development of an effective tag to



discriminate among populations of naked carp of different spawn and nursery sites is of very importance for sustainable increase of naked carp resource. Previous reports based on genetic markers have proven that some degree of genetic divergence existed among naked carp populations of different tributaries of Lake Qinghai [5,8]. But, straying rate of different populations of naked carp is not easy to determine by solely depending on population genetic markers [8]. Differences of water chemistry features among different spawn and nursery grounds provide a consolidate basis for applying the trace element fingerprint of otolith core to discriminate among stocks of naked carp. In the present work, we firstly test this possibility. The results showed that fingerprint of trace elements deposited in the otolith core can significantly discriminate SL and QJ stocks with 100% of classification accuracy rate, although we don't know if the differences of water chemistry of these two neighbor rivers exclusively contribute to this classification for many biotic and abiotic factors affect the deposition of trace elements in the otolith.

Additionally, the contribution of the applied stock enhancement and releasing to the increase of naked carp population remains unknown, although recent investigation showed that it was significantly increased after taking measures to prevent and restore them in the past 30 years [3,6]. And, it is also unknown if the releasing individuals can sexually mature to develop the stable reproduction population to support restoration of naked carp. No economical and practical tag to discriminate releasing individuals from naturally produced individuals seriously prevents from answering the above questions. Furthermore, the local government of Qinghai province is planning to develop the aquaculture industry of naked carp in the future by taking into account the big market demand of this fish. By that time, a robust tool to discriminate between wild and farmed naked carp is prerequisite to prevent against illegal capture of naked carp in the market. Obviously, analysis of genetic spatial structure cannot differentiate wild from farmed naked carp [8,16]. So, development of an appropriate tag is of an important priority.

Although water from SL was drew directly to the artificial

hatching station of naked carp in Gangcha country for artificially reproduction, increasing water temperature and application of some disinfectants (including povidone iodine, ferrous sulfate and potassium permanaganate) are general protocols to assure high hatching rate which possibly make the water chemistry differ from the natural SL water [17]. Our previous analysis also showed that some trace elements in artificially hatchery and all natural spawn grounds of naked carp are significantly different (unpublished data). DFA analysis in the present work could significantly discriminate AP sample from SL and QJ with 100% of success rate, but the centroid of AP group was closer to SL group than QJ group which represented water in artificially hatchery introducing from SL River rather than QJ River to some extent.

Conclusion

In conclusion, this pilot study shows that elemental signatures of core region of otolith can represents a robust tool to discriminate among different stocks of naked carp in Lake Qinghai catchment which will improve us to get more insight into the stock structure of naked carp. Also, the ability of this too to discriminate the farmed from the naturally produced individuals will be useful to assess the contribution of stock enhancement and releasing to resource restoration of naked carp and support the development of aquaculture of naked carp in the future.

Acknowledgements

The authors would like to thank Prof. Yang J and Dr. Jiang T of Freshwater Fishery Center of Chinese Academy of Fishery Sciences in Wuxi City for their help in element fingerprint analysis. The Science and Technology Department of Qinghai Province provide fund to support this work by 2013-Z-715.

References

1. Walker KF, Dunn IG, Edwards D, Petr T, Yang HZ (1996) A fishery in a changing lake environment: the naked carp *Gymnocypris przewalskii* (Kessler) (Cyprinidae: Schizothoracinae) in Qinghai HU, China. *Int J Salt Lake Res* 4: 169-222.
2. Xiong F, Chen DQ, Duan XB (2010) Threatened fishes of the world: *Gymnocypris przewalskii* (Kessler, 1876) (Cyprinidae: Schizothoracinae). *Environ Biol Fish* 87: 351-352.
3. Chen D, Zhang X, Tan K, Wang K, Qiao Y, et al. (2009) Hydroacoustic study of spatial and temporal distribution of *Gymnocypris przewalskii* (Kessler, 1876) in Qinghai Lake, China. *Environ Biol Fish* 84: 231-239.
4. Wang S, Xie Y (2004) China species red list, Red list. Higher Education Press, Beijing.
5. Wang ZX, Shi JQ, Wei FL, Qi DL, Yuan S, et al. (2015) AFLP genetic diversity of scaleless carp (*Gymnocypris przewalskii*) in different freshwater tributaries of Qinghai Lake. *Geno Appl Biol* 34: 82-99.
6. Wang CY, Wei QW, Sun LY, Qi HF, Zhang H, et al. (2013) Distributions and abundance of *Gymnocypris przewalskii* (Kessler, 1876) in Qinghai Lake, China: an approach based on hydroacoustic sampling. *J Appl Ichthyol* 29: 1473-1476.
7. Wood CM, Du J, Rogers J, Brauner CJ, Richards JG, et al. (2007) Przewalski's naked carp (*Gymnocypris przewalskii*): an endangered species taking a metabolic holiday in Qinghai Lake, China. *Physiol Biochem Zool* 80: 59-77.
8. O'Bryan DM, Xie Z, Wang Y, Du J, Brauner CJ, et al. (2010) Phylogeography and conservation genetics of Lake Qinghai scaleless carp (*Gymnocypris przewalskii*). *J Fish Biol* 77: 2072-2092.
9. Campana SE (1999) Chemistry and composition of fish otoliths: pathways, mechanisms and applications. *Mar Ecol Prog Ser* 188: 263-297.
10. Tanner SE, Reis-Santos P, Cabral HN (2016) Otolith chemistry in stock delineation: A brief overview, current challenges and future prospects. *Fish Res* 173: 206-213.
11. Zhang F, Jin ZD, Hu G (2009) Seasonally chemical weathering and CO₂ consumption flux of Lake Qinghai river system in the northeastern Tibetan Plateau. *Environ Earth Sci* 59: 297-313.

12. Zhou L, Jin ZD, Li FC (2012) Mineralogy of the otoliths of naked carp *Gymnocypris przewalskii* (Kessler) from Lake Qinghai and its Sr/Ca potential implications for migratory pattern. Sci China Earth Sci 55: 983-990.
13. Liu HB, Jiang T, Tan XC, Yang J (2012) Preliminary investigation on otolith microchemistry of naked carp (*Gymnocypris przewalskii*) in Lake Qinghai, China. Environ Biol Fish 95: 455-461.
14. Xiong F, Chen DQ, Liu SP, Duan XB, Shi JQ (2006) Annuli characteristics of the different ageing materials of *Gymnocypris przewalskii przewalskii* (Kessler). Acta Hydrobiol Sinica, 30: 185-191.
15. Dou SZ, Amano Y, Yu X, Cao L, Shirai K, et al. (2012) Elemental signature in otolith nuclei for stock discrimination of anadromous tapertail anchovy (*Coilia nasus*) using laser ablation ICPMS. Environ Biol Fish 95: 431-443.
16. Zite A, Sturm M, Waidbacher H, Prohaska T (2010) Discrimination of wild and hatchery trout by natural chronological patterns of elements and isotopes in otolith using LA-ICP-MS. Fisheries Manag Ecol 17: 435-445.
17. Shi JQ, Qi HF, Yang JX, He WH, Hao GC (2000) Study on reproductive biology of *Gymnocypris przewalskii*. Qinghai Sci Tec7: 12-15.

Citation: Zhang JY, Liu XH, Zhao YL, Wei FL, Li CZ (2016) Pilot Test of Elemental Fingerprint in Otolith Core as a Natural Biological Tag to Discriminate among *Gymnocypris przewalskii* Stocks in Lake Qinghai in Qinghai-Tibet Plateau, China. J Aquac Res Development 7: 447. doi: [10.4172/2155-9546.1000447](https://doi.org/10.4172/2155-9546.1000447)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>