

## Plantar Erythema Nodosum Associated with Crohn's Disease

Sumeyra Acar

Department of Gastroenterology, Istanbul University, Istanbul Medical Faculty, Istanbul, Turkey

\*Corresponding author: Sumeyra Acar, Department of Gastroenterology, Istanbul University, Istanbul Medical Faculty, Istanbul, Turkey, Tel: 90 212 440 00; E-mail: acar.sumeyra@gmail.com

Rec date: Mar 6, 2015, Acc date: Mar 7, 2015, Pub date: Mar 16, 2015

Copyright: © 2015 Acar S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

### Medical Image

A 22-year-old woman who previously diagnosed Crohn's disease was admitted with a 1-month history of diarrhea. She reported that during the preceding 3 months, she had stopped taking her drugs for Crohn's disease. She was not able to tread on her left foot. On examination, there was an erythematous, non-purpuric, painful, slightly nodular and, 3-4 cm diameter lesion (Figure 1). Plantar erythema nodosum was diagnosed associated with Crohn's disease and steroid treatment was started. Resolution of the lesion was noted at follow-up 2 weeks later. Plantar erythema nodosum is a very rare entity and typically occurs insidiously in children with good health. There was only one plantar erythema nodosum case associated with Crohn's disease in the literature (Figures 1 and 2).



**Figure 1:** Plantar erythema nodosum associated with Crohn's disease.



**Figure 2:** Plantar erythema nodosum associated with Crohn's disease.