

Plasmids for Amoxicillin and Ciprofloxacin Resistance in *Escherichia coli* Isolate Causing Urinary Tract Infection

Sabrina Suhani, Auditi Purkaystha, Musammat Kulsuma Begum, Md. Jahidul Islam and Abul Kalam Azad*

Department of Genetic Engineering & Biotechnology, Shahjalal University of Science and Technology, Sylhet-3114, Bangladesh

*Corresponding author: Abul Kalam Azad, Shahjalal University of Science and Technology, Sylhet, Bangladesh, Tel: +880 1715 160921; E-mail: dakazad-btc@sust.edu

Received date: June 07, 2017; Accepted date: June 30, 2017; Published date: July 5, 2017

Copyright: ©2017 Suhani S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: *Escherichia coli* causes urinary tract infection (UTI), one of the most frequent bacterial infections in humans, shows resistance to various commonly prescribed antibiotics. This is termed as multi drug resistance, which is increasing among *E. coli*. The multi-drug resistant *E. coli* isolates harbor multiple plasmids that might be responsible for antibiotic resistance. The resistance can be transferred among bacteria through horizontal transfer of the genetic materials.

Objectives: We undertook this study 1) to find a link between antibiotic resistance and plasmid of *E. coli* from UTI patients of Sylhet region of Bangladesh 2) to analyze the spread of the plasmid mediated antibiotic resistant gene between *E. coli* isolates.

Methods: Conjugation experiment was carried out in Luria Broth with *E. coli* DH5 α as recipient. For transformation, competent cells were prepared using *E. coli* DH5 α . Plasmid isolation was done by mini alkaline lysis method and plasmid was extracted from agarose gel and transferred to the competent cell.

Results: After conjugation, the donor *E. coli* isolate that showed resistance to amoxicillin (AMX), ciprofloxacin (CIP) and ceftriaxone (CTR) transferred its AMX and CIP resistance to the recipient *E. coli* DH5 α , that was previously sensitive to all antibiotics. After transformation, the recipient *E. coli* DH5 α became resistant to CIP and AMX, while the donor showed resistance to gentamycin (CN), ceftriaxone (CTR), amoxicillin (AMX), ciprofloxacin (CIP), cefixime CFM, and cotrimoxazole (COT). Plasmid extracted from the transformant revealed that, three genes (3 kb, 5 kb and 20 kb) have been transferred from the donor to recipient.

Conclusion: Plasmids responsible for amoxicillin and ciprofloxacin were transferred.

Keywords: Antimicrobial resistance; Horizontal plasmid transfer; MDR; UTI

Introduction

Urinary tract infection (UTI), usually caused by *Escherichia coli*, is the most common bacterial infection in humans. The risk factors for UTIs include sexual activity, gender, genetics and presence of urinary catheters [1,2]. Females are more prone to UTIs than male because their urethra is much shorter and closer to the anus. Females further lack the bacteriostatic properties of prostatic secretions [1].

E. coli causing UTI belongs to several subgroups that are selected by some factors, which enhance extra intestinal survival. These factors include structural features such as fimbriae or pili for adherence, flagella for motility and chemical adhesion [3]. The type-1 pili of uropathogenic *E. coli* are known to be associated with increased severity of UTI by binding to mannose containing glycoprotein receptors on facet cells lining the bladder or vaginal epithelial cells [4]. Other virulence factors which confer the ability to fecal *E. coli* to colonize the vaginal mucosa and cause symptomatic UTI have also been identified [3]. However, the ability of *E. coli* to cause UTIs is increasing, while the ease of treating these infections due to resistance to first generation antibiotics such as cotrimoxazole, ampicillin and

nitrofurantoin is becoming progressively more elusive. The greater concern is the recent increase in resistance to second or even third generation antibiotics such as ciprofloxacin, levofloxacin and ceftriaxone [2,5,6].

From studies of the following few decades, it has been clear that multidrug resistance phenomena is due to chromosomal genetic elements as well as existing plasmids and their association with other genetic mobile elements. In Japan, the role of plasmids in antibiotic resistance was first recognized by Watanabe and Fukasawa in 1961 [7]. Their report shows that transfer of a plasmid, which is known as resistance transfer factor or R-factor, that carries the resistance genes, is a single step process. Further reports showed that plasmids were carriers of not only multi-drug resistance genes, but also genes associated to contribute to the virulence of the host bacteria [8,9].

Antibiotic resistant bacterial infection has been identified as one of the greatest epidemic threats to human health by the World Health Organization. Within this epidemic, a group of pathogens has been individualized and collectively named ESKAPE (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterobacter*). These are opportunistic pathogens; the majority of them cause hospital-acquired infection that is resistant to antibiotic treatment. Plasmids play a

central role in the dissemination and acquisition of the resistant determinants in these bacteria [10].

The increasing ability of *E. coli* to cause UTIs and the difficulty encountered in treating these infections due to multidrug resistance necessitates updating the knowledge of their drug resistance in a given environment. Horizontal gene transfer between bacterial cells contributes to bacterial adaptation to various environments and, in the long term, to bacterial evolution [11]. In human environments, it results in the undesirable spread of pathogenic, antibiotic resistance, or artificially engineered genes. Three mechanisms of horizontal gene transfer in bacteria are generally known: conjugation, transduction and transformation. Conjugation and transduction involve specific apparatus for DNA transfer from donor cells to recipient cells; they are conjugative pili and phage capsids, respectively. However, transformation is mainly performed by the recipient cells that express genetic competence to take up extracellular free DNA [12]. Competence for transformation can be induced naturally and artificially but not all bacterial species develop natural competence.

E. coli is not assumed to be naturally transformable; it develops high genetic competence only under artificial conditions, e.g. exposure to high Ca^{2+} concentrations [13]. However, several reports have shown that *E. coli* can express modest genetic competence in certain conditions that can arise in its environment and make it subjected to horizontal transformation [11,12,14]. We herein report that inheritance of antibiotic resistance occurs due to transconjugation or transformation of plasmids. It has been further discussed that a particular plasmid is responsible for resistance against a specific antibiotic.

Materials and Methods

E. coli isolates and their maintenance

The multi-drug resistant *E. coli* isolates were obtained from urine samples of UTI patients in North East region of Bangladesh [2]. *E. coli* isolates were identified based on morphological, cultural and biochemical characteristics as described by Cappuccino et al. [15] and Bergey's manual of systematic bacteriology [16]. The reference strain, *E. coli* DH5 α , was collected from department of microbiology, University of Dhaka, Bangladesh. The pure culture of *E. coli* isolates and *E. coli* DH5 α was routinely sub-cultured on nutrient agar. The pure culture was preserved at -30°C with 20% glycerol.

The plasmid transfer in *E. coli* DH5 α through conjugation

A single colony of multidrug resistant *E. coli* isolate (donor) and that of antibiotic sensitive *E. coli* DH5 α (recipient) were inoculated into Luria broth (LB) separately and incubated in a shaker at 37°C for overnight. Five hundred micro litre of donor and recipient cultures was added to 4 ml fresh LB and incubated overnight at 37°C without shaking [17]. A 10-fold serial dilution was then done with LB and 100 μ l from each dilution was plated into sorbitol-MacConkey agar (SMAC medium) [18] containing selected antibiotics. SMAC medium was used as a differential color forming medium for detection of sorbitol fermenting (donor) isolate with characteristic red to colorless colonies and non-sorbitol-fermenting (recipient) isolate with characteristic colorless colonies. In SMAC, the lactose of MacConkey agar was replaced by sorbitol. After conjugation, the recipient *E. coli* DH5 α transformed with plasmid produced pink colonies on the antibiotic selection plate.

Extraction and purification of plasmids

Plasmids were extracted by mini alkaline lysis method [19] from 16 h cultured cells of *E. coli*. Plasmids were then subjected to electrophoresis onto 0.7% agarose gel using standard protocol. The gels stained with ethidium bromide were visualized on an ultraviolet transilluminator (UVP, High Performance transilluminator; USA) and recorded with a canon camera (PowerShot A3200 IS). A particular plasmid band was excised from the gel and purified with the FavorPrep™ plasmid DNA extraction mini kit (Favorgen Biotech Corp., Taiwan).

Competent cell preparation, plasmid transformation and antibiotic susceptibility

Competent cells of *E. coli* DH5 α were prepared as described by Sambrook et al. [19]. For transformation, 50 μ l of plasmid DNA was added to a tube containing 0.2 ml of competent cells. The mixture was placed on ice for 20 min, and then exposed to heat shock at 42°C for 1 min and kept immediately in ice for 10 min. A 800 μ l LB broth was then added to transformation mixture and incubated at 37°C for 60 min. About 100 μ l from transformation mixture was spread on nutrient agar plates containing the appropriate antibiotic. An aliquot (100 μ l of 5-10 fold diluted) of competent cells spread on nutrient agar containing the same antibiotics was used as a control. All plates were incubated at 37°C for 48 h. The transformant colonies were scored and fresh cultured several times on plates containing the desired antibiotics. The transformant was investigated for the presence of target plasmid. The susceptibility of transformant and non-transformant *E. coli* DH5 α to different antibiotics was investigated by the modified Kirby-Bauer disc diffusion method [20].

Results

Inherence of antibiotic resistance in *E. coli* DH5 α through conjugation

E. coli isolate resistant to amoxicillin (AMX), ceftriaxone (CTR) and ciprofloxacin (CIP) and *E. coli* DH5 α sensitive to these antibiotics were used as donor and recipient, respectively (Figure 1A and B).




Figure 1: Antibiogram of *E. coli* isolate and *E. coli* DH5 α . *E. coli* isolate was resistance to AMX, CTR and CIP (A), whereas *E. coli* DH5 α was sensitive to all antibiotics investigated (B).

SMAC media was used for selecting the transconjugants. The donor *E. coli* isolate produced red to pink colonies (Figure 2A) due to acid production by fermentation of sorbitol in the media. The recipient *E. coli* DH5 α cannot ferment sorbitol, and use peptone instead [17]. This forms ammonia, which raises the pH of the agar and lead to the formation of white/colorless colonies (Figure 2B).




Figure 2: Cultural characteristics of *E. coli* isolate (A) and *E. coli* DH5 α (B) on SMAC.

Following conjugation, less-whitish colonies of *E. coli* DH5 α were observed on SMAC media in the presence of AMX and CIP (Figure 3A and B) but not in the presence of CTR (Figure 3C). Growth of *E. coli* DH5 α in the presence of AMX and CIP indicated that resistance to these antibiotics was inherited to them due to transfer of plasmid form the donor through conjugation.



Figure 3: Growth of the donor, recipient and transconjugant on SMAC media containing AMX (A), CIP (B) and CTR (C).

Plasmid DNA transformation makes *E. coli* DH5 α resistant to AMX and CIP

After plasmid DNA transformation, the transformant *E. coli* DH5 α was screened in the presence of AMX and CIP separately. It was

observed that some transformants could grow in the presence of these two antibiotics (Figure 4A-D).




Figure 4: Selection of transformants in nutrient agar containing AMX (A) Growth of *E. coli* DH5 α without AMX (positive control), (B) transformant growing in the media containing AMX, (C) no growth of *E. coli* DH5 α in the presence of AMX (negative control), (D) growth of donor in the presence of AMX.

After selection in the media containing antibiotic, the transformants were subjected to antibiogram of some antibiotics to determine which antibiotic resistant gene had been transferred to transformant *E. coli* DH5 α . For this, eight antibiotic disks were used to compare the

susceptibility of the transformants (Figure 5B) to that of *E. coli* DH5 α (Figure 5C) and multidrug resistant *E. coli* isolate from which plasmid was extracted (Figure 5A).




Figure 5: Antibiogram of *E. coli* isolate (A), transformant (B) and *E. coli* DH5α (C).

It was observed that the MDR *E. coli* isolates were resistant to six antibiotics and *E. coli* DH5α was sensitive to all antibiotics. The transformant *E. coli* DH5α inherited resistance to two antibiotics, CIP and AMX. This result indicated that CIP and AMX resistance genes might have been transferred to *E. coli* DH5α.

Plasmid profiling of transconjugates and transformants

Plasmids were isolated from the donor or MDR *E. coli* isolate, transconjugates, transformants and recipient *E. coli* DH5α and their plasmid profiling were compared. It was observed that the donor *E. coli* isolate harbored multiple plasmids of different sizes, whereas, the recipient *E. coli* DH5α had no plasmid (Figure 6).




Figure 6: Electrogram of plasmids from donor (D, lanes 2 and 4), transconjugants (TC, lane 3), transformant (T, lane 5), recipient (R, lane 6); DNA ladder (L) is shown in lane 1.

However, AMX and CIP resistant transconjugates and transformants harbored plasmids of about 3 kb, 5 kb and 20 kb, which were present in the donor or MDR *E. coli* isolate. This result indicated that the recipient or the competent cells obtained these plasmids during conjugation and transformation.

Discussion

The emergence, proliferation, aggregation, and management of antimicrobial-resistant pathogenic bacteria have become a worldwide health concern. When the resistance associated with the presence of transposable DNA elements, such as plasmids, it may be difficult to eliminate resistance, which tends to increase over time [21]. Our previous study depicted that *E. coli* isolates showing MDR were found

to harbor multiple plasmids of different sizes and left an assumption that the MDR genes might be in the plasmid or chromosomal DNA [2]. It has been reported that Enterobacteriaceae isolates might be potential carrier of antibiotic resistant genes that could be transferred between bacterial strains of the same or different species of UTI pathogens [22-24]. In the present study, we investigated the plasmid associated antibiotic resistance in *E. coli* isolates causing UTI.

The present study showed that plasmids responsible for AMX and CIP resistance were transferred to the recipient *E. coli* DH5α. Many studies have shown that the dissemination of resistant genes mediated by plasmids may occur by conjugation [25-28].

Genetic transformation is an important mechanism of the horizontal exchange of genes. Natural genetic transformation is a gene transfer process where the bacteria can pick up the naked DNA from their environment and the DNA may come from a variety of sources. The most frequent source is remnants from dead bacterial cells. The exogenous DNA bind to specific cell surface receptors, then the DNA is transported across the membrane and one strand of the DNA is digested away. As a consequence, the single stranded DNA enters the cell. Recombination occurs between the incorporated DNA and homologous host DNA and the new DNA replaces a strand of the host DNA. This new DNA may contain noble and different genes compared to the host DNA, including antibiotic resistance gene [29]. An investigation carried out in the UK revealed that Enterobacteriaceae isolates from 43 hospitals during 1990-1991 showed antibiotic resistance due to plasmid DNA by transformation [30]. Another study reported transformation process responsible for chloramphenicol, tetracycline and sulfonamide resistance transferred with the CTX-M-1 enzyme as co-transformant [31].

The present study showed that three plasmids, eg. 3 kb, 5 kb and 20 kb were transferred to the recipient cell, and thereby, the recipient *E. coli* DH5α became resistant to CIP and AMX. However, the non transformant recipient cells remained sensitive to these antibiotics. These three or any of these plasmids might carry CIP and AMX resistance gene. This report reveals the spread of the plasmid mediated antibiotic resistant genes between *E. coli* isolates of UTI patient and *E. coli* DH5α. Nevertheless, further research is needed to determine the specific gene in these plasmids associated with CIP and AMX resistance.

Conclusion

Our study revealed that, following conjugation and transformation, amoxicillin and ciprofloxacin resistance was inherited by *E. coli* DH5α that was previously sensitive to all antibiotics, from the MRD *E. coli* isolates obtained from UTI patients. This finding indicates that, plasmids responsible for antibiotic resistance can be transferred among bacteria through horizontal gene transfer. Careful use of antimicrobial agents, developing standard treatment guidelines and increasing awareness among people can ensure proper health care and reduce the spread of antimicrobial resistance among bacteria.

References

1. Nicolle LE (2008) Uncomplicated urinary tract infection in adults including uncomplicated pyelonephritis. Urol Clin North Am 35(1): 1-12.
2. Rabbee MF, Begum MK, Islam MJ, Chowdhury PA, Chowdhury OA, et al. (2016) Multidrug Resistance Phenotype and Plasmid Profiling of *Escherichia coli* Isolates Causing Urinary Tract Infections in North East Part of Bangladesh. Br Microbiol Res J 15(6): 1-11.

3. Emo L, Kerenyi M, Nagy G (2003) Virulence factors of uropathogenic *Escherichia coli*. Int J Antimicrob Agents 22: 29-33.
4. Venegas MF, Navas EL, Gaffney RA, Duncan JL, Anderson BE, et al. (1995) Binding of type 1-piliated *Escherichia coli* to vaginal mucus. Infection and immunity 63(2): 416-422.
5. Karlowsky JA, Hoban DJ, Decorby MR, Laing NM, Zhanel GG (2006) Fluoroquinolone-resistant urinary isolates of *Escherichia coli* from outpatients are frequently multidrug resistant: results from the North American Urinary Tract Infection Collaborative Alliance-Quinolone Resistance study. Antimicrob Agents Chemother 50(6): 2251-2254.
6. Hooton TM (2000) Pathogenesis of urinary tract infections: an update. J Antimicrob Chemother. 46(suppl 1): 1-7.
7. Watanabe T, Fukasawa T (1961) Episome-Mediated Transfer of Drug Resistance In Enterobacteriaceae I: Transfer of Resistance Factors by Conjugation. J Bacteriol 81(5): 669.
8. Watanabe T (1963) Infective heredity of multiple drug resistance in bacteria. Bacteriol Rev 27(1): 87.
9. Ramirez MS, Traglia GM, Lin DL, Tran T, Tolmasey ME (2014) Plasmid-mediated antibiotic resistance and virulence in gram-negatives: the *Klebsiella pneumoniae* paradigm. Microbiol Spectr 2(5): 1.
10. Boucher HW, Talbot GH, Benjamin DK, Bradley J, Guidos RJ, et al. (2013) 10 × 20 progress—development of new drugs active against gram-negative bacilli: an update from the Infectious Diseases Society of America. Clin Infect Dis 56(12): 1685-1694.
11. Thomas CM, Nielsen KM (2005) Mechanisms of, and barriers to, horizontal gene transfer between bacteria. Nat Rev Microbiol 3(9): 711-721.
12. Keese P (2008) Risks from GMOs due to horizontal gene transfer. Environ Biosafety Res 7(3): 123-149.
13. Méjean V, Claverys JP (1993) DNA processing during entry in transformation of *Streptococcus pneumoniae*. J Biol Chem 268(8): 5594-5599.
14. Lorenz MG, Wackernagel W (1994) Bacterial gene transfer by natural genetic transformation in the environment. Microbiol Rev 58(3): 563-602.
15. Cappuccino J, Sherman N (2001) Microbiology: A Laboratory Manual.
16. Whitman W, Goodfellow M, Kämpfer P, Busse HJ, Trujillo M, et al. (2012) The Actinobacteria Bergey's Manual of Systematic Bacteriology: Volume 5 Springer Science & Business Media.
17. Wang A, Yang Y, Lu Q, Wang Y, Chen Y, et al. (2008) Presence of *qnr* gene in *Escherichia coli* and *Klebsiella pneumoniae* resistant to ciprofloxacin isolated from pediatric patients in China. BMC Infect Dis 8(1): 68.
18. March SB, Ratnam SA (1986) Sorbitol-MacConkey medium for detection of *Escherichia coli* O157: H7 associated with hemorrhagic colitis. J Clin Microbiol 23(5): 869-872.
19. Sambrook J, Russell D (2001) Molecular Cloning: A Laboratory manual. (3 732 volume set), Cold Spring Harbor Laboratory Press, 2344.
20. Bauer AW, Kirby WM, Sherris JC, Turck M (1966) Antibiotic susceptibility testing by a standardized single disk method. Am J Clin Pathol 45(4): 493.
21. Kazemnia A, Ahmadi M, Dilmaghani M (2014) Antibiotic resistance pattern of different *Escherichia coli* phylogenetic groups isolated from human urinary tract infection and avian colibacillosis. Iran Biomed J 18(4): 219.
22. Österblad M, Hakanen A, Manninen R, Leistevuo T, Peltonen R, et al. (2000) A between-species comparison of antimicrobial resistance in enterobacteria in fecal flora. Antimicrob Agents Chemother 44(6): 1479-1484.
23. Lawrence JG (2005) Horizontal and vertical gene transfer: the life history of pathogens, in Concepts in Bacterial Virulence. Contrib Microbiol 12: 255-271.
24. Tenover FC (2006) Mechanisms of antimicrobial resistance in bacteria. The American journal of medicine. Am J Med 119(6): S3-S10.
25. Aluyi H, Akortha E (2002) Transfer of ampicillin resistance gene (*amp^r*) from some enteric bacteria of diarrhoeal origin to *Escherichia coli* (UBE201). J Med Lab Sci 11(2): 39-41.
26. Díaz MA, Cooper RK, Cloeckeaert A, Siebeling RJ (2006) Plasmid-mediated high-level gentamicin resistance among enteric bacteria isolated from pet turtles in Louisiana. Appl Environ Microbiol 72(1): 306-312.
27. Ferguson GC, Heinemann JA, Kennedy MA (2002) Gene transfer between *Salmonella enterica* serovar Typhimurium inside epithelial cells. J Bacteriol 184(8): 2235-2242.
28. Shoemaker NB, Vlamakis H, Hayes K, Salyers AA (2001) Evidence for Extensive Resistance Gene Transfer among *Bacteroides* spp. and among *Bacteroides* and Other Genera in the Human Colon. Appl Environ Microbiol 67(2): 561-568.
29. Navarro F (2010) Acquisition and horizontal diffusion of β -lactam resistance among clinically relevant microorganisms. Int Microbiol 9(2): 79-81.
30. Piddock LJ, Walters RN, Jin YF, Turner HL, Gascoyne-Binzi DM, et al. (1997) Prevalence and mechanism of resistance to third-generation cephalosporins in clinically relevant isolates of Enterobacteriaceae from 43 hospitals in the UK, 1990-1991. J Antimicrob Chemother 39(2): 177-187.
31. Arpin C, Dubois V, Coulange L, André C, Fischer I, et al. (2003) Extended-spectrum β -lactamase-producing Enterobacteriaceae in community and private health care centers. Antimicrob Agents Chemother 47(11): 3506-3514.