

Post-Industrial Dumps as a “Unique” Experiment for Ecologists

Skubala Piotr

Department of Ecology, University of Silesia, Bankowa 9, 40-007 Katowice, Poland

Corresponding author: Piotr S, Department of Ecology, University of Silesia, Bankowa 9, 40-007 Katowice, Poland, Tel: 48 32 359-11-48; Email: piotr.skubala@us.edu.pl

Received date: March 11, 2017; **Accepted date:** March 17, 2017; **Published date:** March 21, 2017

Copyright: © 2017 Piotr S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Piotr S (2017) Post-Industrial Dumps as a “Unique” Experiment for Ecologists. J Pollut Eff Cont 5: e115. doi:[10.4172/2375-4397.1000e115](https://doi.org/10.4172/2375-4397.1000e115)

Editorial

Post-industrial wastelands are still constantly increasing in area; some of them may be dangerous to human health. These facts provide the major stimulus for the reclamation and restoration of large devastated areas. The reclamation of dumps, especially of toxic dumps, is a challenge and is a severe problem of today. For this purpose, a knowledge and understanding of soil development – mainly induced by soil organisms – is an important prerequisite for any form of soil management on dumps [1]. From the viewpoint of reclamation, a practice very poorly developed at this time, knowledge of soil fauna development is required as an indicator for artificial cultivation techniques [2]. The restoration of disturbed ecosystems has recently received increasing attention from ecologists and land-use managers. This is due, in a large part, to more stringent government regulations, which mandate extensive reclamation procedures followed by biological monitoring efforts [3]. It seems obvious that dumps should at first be managed biologically. In this way the negative influence of dumps on the natural environment may be limited, and the beauty of the landscape can be improved. Nevertheless, vegetation cover on dumps may develop naturally. Hitherto, reclamation measures taken on dumps have appeared unsatisfactory in many cases despite the high expenditure of money and effort. Spontaneous succession of plants is more desirable [4, 5], but unfortunately the spontaneous process of colonization and succession is a long-term process.

Much of the research on land restoration has been devoted to the reestablishment of vegetation and vertebrates. Soil fauna has been treated marginally in many restoration practices and studies, whereas we can obtain useful results for derelict land management, acquiring knowledge of all processes during succession on wastelands. Studies on recolonization of arthropods on wastelands are few [6-10], whereas arthropods can drastically influence revegetation efforts via herbivory, seed predation, litter decomposition, pollination, and soil aeration [11]. More empirical studies of these processes on different kinds of dumps are urgently needed to better understand the factors affecting soil fauna development in a post-industrial landscape [12].

The growing area of dumps is a real disaster for humankind, although from the viewpoint of soil ecology dumps are not “waste land”. They offer a tremendous experimental field in which to study the colonization by animals and the development of their communities in this hostile environment. A new habitat for many new inhabitants is created as a result of spoil heap construction. A dump is a “land” lacking plants and animals, initially with a complete absence of soil. Then for a long period the “soil” that is present lacks stratification, has insufficient organic matter, few nutrients and an inadequate water content. Additionally, what makes post-industrial dumps an excellent polygon on which to test ecological hypotheses is the simplified and

variable relationships between living components. Furthermore, they are usually not under threat by prospectors. The variability of spoil heaps and the range of environmental conditions established on post-industrial dumps result in a heterogeneous environment for soil biota [13, 14]. Furthermore, different reclamation technologies used on dumps also create varied conditions. So ecologists get an experiment with many differentiated environmental variables

References

1. Wanner M, Dunger W, Schulz H-J, Voigtländer K (1998) Primary immigration of soil organisms on coal mined areas in Eastern Germany, in: Soil Zoological Problems in Central Europe, Pižl V, Tajovský K, Editors, České Budějovice, 267-275.
2. Dunger W, Wanner M, Hauser H, Hohberg K, Schulz H-J, et al. (2001) Development of soil fauna at mine sites during 46 years after afforestation. *Pedobiologia* 45: 243-271.
3. Parmenter RR, MacMahon JA (1987) Early Successional Patterns of Arthropod Recolonization on Reclaimed Strip Mines in Southwestern Wyoming: The Ground-dwelling Beetle Fauna (Coleoptera). *Environ Entomol* 16: 168-177.
4. Skubala P (2006) Do we really need land reclamation on dumps? (Oribatid fauna case studies). in: *Advances in Polish Acarology*, Gabryś G, Ignatowicz S, Editors. Wyd. SGGW, Warszawa, 367-374.
5. Weidemann G, Koehler H, Schriefer Th (1982) Recultivation: a problem of stabilization during ecosystem development, in: *Urban Ecology*, Bornkamm R, Lee JA, Seaward MRD. Blackwell Oxford, 305-313.
6. Majer JD (1985) Recolonization by ants of rehabilitated mineral sand mines on North Stradbroke Island, Queensland, with particular reference to seed removal. *Aust J Ecol* 10: 31-48.
7. Neumann U (1971) Die Sukzession der Bodenfauna (Carabidae [Coleoptera], Diplopoda und Isopoda) in den forstlich rekultivierten Gebieten des Rheinischen Braunkohlenreviers. *Pedobiologia* 11: 193-226.
8. Skubala P (1999) Colonization of a dolomitic dump by oribatid mites (Acarina, Oribatida). *Pedobiologia* 43: 145-159.
9. Skubala P, Rola K, Osyczka P (2016) Oribatid communities and heavy metal bioaccumulation in selected species associated with lichens in a strongly metal contaminated habitat. *Environ Sci Pollut Res* 23: 8861-8871.
10. Usher MB (1979) Natural communities of plants and animals in disused quarries. *J Environ Managem* 8: 223-236.
11. Majer JD (1989) *Animals in primary succession. The role of fauna in reclaimed lands.* Cambridge University Press, New York.
12. Skubala P (1995) Moss mites (Acarina: Oribatida) on industrial dumps of different ages. *Pedobiologia* 39: 170-184.
13. Dunger W (1991) Zur primärsukzession humiphager tiergruppen auf bergbauflächen. *Zool Jahrb Syst* 118: 423-447.
14. Skubala P, Rola K, Osyczka P, Kafel A (2014) Oribatid Mite Communities on Lichens in Heavily Contaminated Post-Smelting Dumps. *Arch Environ Contam Toxicol* 67: 578-592.