

PPAR Gamma at the Crossroads of Health and Disease: A Masterchef in Metabolic Homeostasis

Souhad El Akoum^{1,2*}

¹CHUM Research Center, 900 rue St-Denis, Montréal (Qc) H2X 0A9, Canada

²Département d'Immunologie et de Microbiologie, Faculté de Médecine, Université de Montréal, Canada

*Corresponding author: Souhad El Akoum, Département d'Immunologie et de Microbiologie, Faculté de Médecine, Université de Montréal, 2900 boulevard Edouard-Montpetit, Montréal (QC) H3T 1J4, Canada, Tel: 1-514-890-8000; E-mail: souhad.el.akoum@umontreal.ca

Received date: Jan 28, 2014, Accepted date: Mar 27, 2014, Published date: Mar 31, 2014

Copyright: © 2014 El Akoum S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The peroxisome proliferator-activated receptor gamma (PPAR γ) is a ligand-activated transcription factor involved conferring the role of an active gland to the adipose tissue. This nuclear receptor is actively involved, mainly through its regulation to the physiology and the endocrine activity of the adipose tissue, in the regulation of a variety of processes governing the metabolic homeostasis. PPAR γ , activated by a wide variety of fatty acids molecule or their metabolite, governs metabolic processes implicated in glucose and lipid metabolism and adipose mass control by modulating the expression of a large number of target genes. Furthermore, PPAR γ is a molecular target for antidiabetic thiazolidinedione molecules that selectively bind this nuclear receptor to improve systemic insulin sensitivity and glucose tolerance. Accordingly, the specific position of PPAR γ in systemic metabolic control is resumed in its pivotal role in the regulation of glucose and lipid homeostasis, lipid storage and adipogenesis. Here, we present an overview of the involvement of PPAR γ in metabolic control leading to health improvement.

The emphasis is on adipose tissue mass regulation by PPAR γ and its implication in glucose homeostasis and cardiovascular modulation.

Keywords: Metabolic homeostasis; Adipogenesis; Cardiovascular modulation

Review

Since their identification in 1990s [1], the peroxisome proliferator-activated receptors (PPARs) are soliciting considerable interest and our knowledge about their physiological roles is evolving. Involved in the regulation of the metabolism of lipids and lipoproteins, glucose homeostasis, cell proliferation and cell differentiation, PPARs are a pharmacological target for the treatment of metabolic disorders such as diabetes or dyslipidemia.

The discovery of PPARs started when our view towards adipose tissue changed [2]; long believed to be a simple energy storage tissue, adipose tissue is nowadays considered an endocrine gland itself. The development of fibrate family to treat hyperlipidemia in 1962 [3,4] led to the discovery in the 1990s of the first PPARs member the PPAR α [5]. This finding thus stimulated worldwide research to elucidate PPARs family's role in the systemic metabolism control [6].


Peroxisome Proliferation Response Elements (PPRE) were then described in the promoter of microsomal and peroxisomal genes known to be upregulated during proliferation of peroxisome as CYP4A1, CYP4A6 and acyl CoA oxidase [7-9]. International interest in the study of PPARs took on even greater significance after the identification and cloning of three subtypes of PPAR [10,11], named α , β/δ and γ , each being encoded by a specific gene to play key roles in metabolic homeostasis [12-15].

PPAR γ a ligand-activated nuclear receptor

PPAR γ was first identified as a mediator of the activity of the oral antidiabetic thiazolidinedione (TZD) family and recognized as a major regulator of glucose homeostasis and adipogenesis [16,17]. PPAR γ belongs to the nuclear receptors superfamily of ligand-inducible transcription factors [13,18]. Often of lipid nature, ligands activate PPAR γ forcing it to bind to PPRE of the promoter region of specific target genes involved in adipogenesis, lipid metabolism, inflammation and metabolic homeostasis. Similarly to typical nuclear receptors domain structure, PPAR γ primary structure is composed of approximately 500 amino acids, and their structure is represented by a sequence of six areas (Figure 1). The N-terminal domain (A/B domain) is of length and primary structure variable from one receptor to another. It contains the ligand-independent transactivation segment (AF-1, Activation Function-1) that binds co-activators.

The C domain contains the DNA Binding Domain (DBD) characterized by a double folding of the protein chain held by two zinc atoms interacting with four cystein residues. The DNA binding occurs on consensus sequences called hormone response element located before the target gene near the promoter. This domain is highly conserved between nuclear receptors sequences allowing the three members of PPARs family to bind to the same PPRE DNA sequence. The D domain is a hinge region involved in binding of the chaperone protein to the receptor and in the DNA binding. The E domain liaises mediators on the C-terminal portion of the receptor. This Ligand-Binding Domain (LBD) also provides receptor dimerization and comprises a second pattern of AF-2 transactivation. This later protein sequences differs between the three subtypes of PPARs members leading to three pharmacologically distinct forms of nuclear receptors

appointed α , β/δ and γ . The F domain, meanwhile, is a variable sequence that constitutes the C-terminal portion of the protein sequence of each receptor.



Although these domains are all potential targets mediating its signaling cascade, PPAR γ transcriptional activity is initiated by endogenous and exogenous ligands. These later induce chaperone proteins dissociation from the nuclear receptor that represses its activity and conformational changes allowing PPAR γ heterodimerisation with the retinoid X receptors (RXR, vitamin A and 9-cis-retinoic acid receptor) [5] (Figure 1). The complex, thus enabled, will bind to the typical PPSE sequences located in the promoter regions of many genes involved in adipogenesis, adipokines secretion and glucose and lipid homeostasis (Table 1) whose expression is thus stimulated [19].

Adipogenesis	
FABP4	Fatty Acid Binding Protein 4, also known as adipocyte fatty acid binding protein P2 (aP2)
Pref-1	Preadipocyte factor 1
UCP1	UnCoupling Protein 1 (key determinant of brown adipocytes)
PLIN1/2/4	Perilipin-1, 2 & 4, also known as lipid droplet-associated protein
C/EBP α	CCAAT/enhancer binding protein (C/EBP), alpha
STAT 1	Signal Transducers and Activators of Transcription-1
STAT5A/B	

CDKN1A	Cyclin-Dependent Kinase inhibitor 1A
Cidec	Cell death-inducing DFFA-like effector c
Nr1d1	Nuclear receptor subfamily 1, group D, member 1
Adipokine secretion	
ADPN	Adiponectin, also known as adipoQ
FGF1/21	Fibroblast Growth Factor 1 & 21
Ob (Lep)	Leptin
RETN	Resistin also known as adipose tissue-specific secretory factor (ADSF)
APLN	Apelin
ACS	Acylation Stimulating Protein
FIAF	Fasting Induced Adipose Factor
OMN	Omentin
Rbp-4	Retinol Binding Protein-4
Serpina12	Vaspin
PBEF1	Visfatin also called pre-B cell enhancing factor (PBEF) or nicotinamide phosphoribosyltransferase (Nampt)
RARRES2	Chemerin, also known as retinoic acid receptor responder protein 2 (RARRES2)
SERPINE1	Plasminogen activator inhibitor-1 (PAI-1)
ANGPTL2	Angiotensin-like Protein 2
Lipid homeostasis	
Lpl	Lipoprotein lipase
Gyk	Glycerol kinase, key player of glycolysis/glycogenesis process
ACS	Acetyl-CoA synthetase
Pnpla2	Patatin-like phospholipase domain containing 2, involved in the triglyceride hydrolysis
Dbi	Diazepam binding inhibitor, involved in lipid metabolism and dislocation of β -carbolines and benzodiazepines
ACACA	Acetyl-CoA carboxylase- α
ELOVL4	Elongation of very long-chain fatty acids-like 4
LXRA	Liver X receptor α
ME1	Malic enzyme 1; involved in acetyl-CoA is transport
SCD1	Stearoyl-CoA desaturase 1, (Δ 9)-desaturase
APOA2	Apolipoprotein A-II
APOE	Apolipoprotein E
CD36	Leukocyte differentiation antigen 36, also known as fatty acid translocase,

	FAT
LDLR	LDL receptor
LIPC	Hepatic triglyceride lipase (HTGL)
LRP1	LDL receptor-related protein 1
LPL	Lipoprotein lipase
OLR1	Oxidized LDL (oxLDL) receptor, also known as the endothelial oxLDL receptor: LOX-1
FATP1/2	Fatty acid transport protein 1 & 2
Glucose homeostasis	
Glut-4	Glucose transporter type 4
PI3K	Phosphatidylinositol-4,5-bisphosphate 3-kinase
CAP	Catabolite Activator Protein
IRS-1/2	Insulin receptor substrate 1 & 2
Sorbs1	Sorbin and SH3 domain containing 1
Aqp7	Aquaporin 7
G6PC	Glucose-6-phosphatase
GPD1	Glycerol-3-phosphate dehydrogenase 1
GCK	Glucokinase
PEPCK 1	Phosphoenolpyruvate carboxykinase 1
PDK4	Pyruvate dehydrogenase kinase 4
ACAT1	Acetyl-CoA acetyltransferase
Other factor type	
EBF1	Early B cell factor 1, Transcription factor essential for the maintenance of B cell identity
GSTA2	Glutathione S-transferase alpha 2
BR1	Bradykin receptor type1
NFkb	Nuclear factor-kappa B

Table 1: PPAR γ target genes

A wide range of compounds could be identified as PPAR γ ligands [20]. Several studies have focused their research on the elucidation of the link between the lipid nature and their capacity to activate PPAR γ [20,21]. Dietary fats and oils are major source of PPAR γ activators especially polyunsaturated fatty acids (PUFA) that activate it in micromolar concentration [3,22]. Other natural ligands derived from arachidonic degradation such as 15-deoxy- Δ 12,14-prostaglandin J2 acid show high affinity to PPAR γ [23]. Finally, the FFA oxidized phospholipids derived from oxidized LDL (9- and 13-HODE) can also activate PPAR γ . Depending on its ligand nature, PPAR γ activation can be differently modulated [24-30]. All these potential natural ligands are summarized in the Table 2.

Unsaturated Fatty Acids	
-------------------------	--

ω 3-PUFA	
α -Linolenic acid	[175]
γ -Linolenic acid	[176]
Eicosapentaenoic acid (EPA)	[177]
Docohexaenoic acid (DHA)	[178]
4-Hydroxy docosahexaenoic acid (4-HDHA)	[178]
4-Oxodocosahexaenoic acid (4-oxo-DHA)	[178]
ω 6-PUFA	
Linoleic acid	[179]
Nitrolinoleic acid	[179]
Conjugated linoleic acid isomers (CLA)	[180]
9/10-NO ₂ -linoleic acid	[181]
12-NO ₂ -linoleic acid	[181]
13-NO ₂ -linoleic acid	[181]
Arachidonic acid	[182]
ω 9-MUFA	
Palmitoleic acid	[183]
Oleic acid	[184]
Eicosanoids	
9-Hydroxyoctadecadienoic acid (9-HODE)	[179]
13-Hydroxyoctadecadienoic acid (13-HODE)	[179]
15-Deoxy- Δ 12, 14-PGJ2	[185]
Other	
Azelaoyl phosphatidylcholine (component of the lipid pool within oxLDL)	[186]
Isoflavones:	
Genistein	[187]
Daidzein	[188,189]
Equol	[188,189]
Biochanin A	[189]
Flavonoids:	
Psi-baptigenin	[190]
Hesperidin	
Quercetin (from dill, bay leaves, and oregano)	[191]
2'-Hydroxy chalcone (cinnamon in polymeric form)	[191]
Rosmarinic acid (marjoram)	[191]

Table 2: PPAR γ natural ligands

There is also well-known synthetic ligand of PPAR γ thiazolidinedione family (TZD) for treatment of type 2 diabetes and insulin resistance [16,17,31]. Other active ingredients such as non-steroidal anti-inflammatory drugs may also be PPAR agonists.

Tissue distribution of PPAR γ

The tissue distribution and the expression level of PPARs differ for each isoform [32,33]. PPAR γ is mainly found in adipose tissue and the gastrointestinal tract. The PPAR γ gene encodes two protein sub-types, γ 1 and γ 2, arising from differential splicing of exon B [34-36]. PPAR γ 2 possess 28 amino acids more than its counterpart γ 1 at its N-terminus protein sequence in mice and 30 in humans. This additional peptide sequence confers transcriptional activity 10 times higher for PPAR γ 2 compared to PPAR γ 1 subtype [34]. Whereas PPAR γ 1 is expressed in muscle cells, hepatocytes, monocytes and others, PPAR γ 2 is, for its part, the specific form of PPAR γ in the adipose tissue [37,38]. Recently, two new subtypes of PPAR γ were identified in human: PPAR γ 3 and PPAR γ 4 from two different promoters. PPAR γ 4 expression seems to be restricted to adipose tissue [39] while mRNA PPAR γ 3 was detected in white adipose tissue, large intestine and macrophages [37,40].

PPAR γ is also expressed in cells of the vascular wall, monocytes and macrophages [38,41-43]. Further, PPAR γ is present in the atherosclerotic plaque at the sub-endothelial area in the lipid core and of atherosclerotic lesions where they co-localize with specific markers of macrophages, smooth muscle cells and foam cells [44-46].

Adipose tissue and PPAR γ

PPAR α and PPAR δ/β appear to have limited adipogenic effect on the adipose tissue. The PPAR α is mainly expressed in the brown adipose tissue controlling β -oxidation for heat production [47-50], while the PPAR δ/β is found in preadipocytes controlling the expression of genes involved in their proliferation but seems to be slightly implicated in the adipogenesis process [51].

In adipose tissue, the predominant PPAR isoform controlling its differentiation is PPAR γ particularly PPAR γ 2 subtype [35,52-54] (Figure 1). PPAR γ is involved in both processes of adipogenesis, lipid metabolism and the secretion of several hormones called adipokine in adipose tissue. Thus, PPAR γ confers the endocrine functions of the mature adipocyte.

PPAR γ and adipose mass control

PPAR γ is the main nuclear receptor implicated in the adipose mass control triggering the recruitment of new preadipocytes and steer their differentiation into mature adipocytes controlling thus the adipose tissue homeostasis [17,55]. The expression of this nuclear receptor is highly important for the embryogenic development and a decrease in its activity lead to a lipodystrophy in human [39,56-59].

Adipogenesis refers to the process of differentiation of progenitor cells, called preadipocytes, into mature adipocytes in which the gene expression, the cell morphology and the sensitivity to exogenous hormones and factors change. During differentiation, the expression of various genes encoding proteins involved in lipid uptake and metabolism, such as aP2, Pref-1, phosphoenolpyruvate carboxykinase (PEPCK) and lipoprotein lipase (LPL) (Table 1), are induced through the activation of PPAR γ [54,60,61]. PPAR γ is the most important factor implicated in the formation of mature adipocytes and its

overexpression in non-adipocytes cells is sufficient to induce their transformation to adipocytes [52,53]. Furthermore, an increased level of circulating fatty acids in the body is believed to raise PPAR γ activity that will lead to increased adipose tissue mass and obesity development. If so, it explains the fact that TZD treatment, through its activation of PPAR γ , increase body weight gain. However, it has been shown that selective activation of PPAR γ in the adipose tissue is sufficient to prevent diabetes in HFD-fed mice without any change in their body mass [62]. These data demonstrate that PPAR γ adipose tissue activation is essential to improve insulin sensitization but not responsible of the nuclear receptor activation side effect such as weight gain. Moreover, recent studies showed that PPAR γ activation in the brain, rather than in the adipose tissue, is directly linked to weight gain [63,64]. Thus, the development of a treatment that could selectively activate adipose tissue PPAR γ seems to be a highly interesting alternative of a TZD treatment.

In parallel with its adipogenic activity, PPAR γ seems to induce apoptosis in adipocytes in a process of regeneration and cells turnover of the adipose tissue [55,65]. Thus, PPAR γ regulates the adipose tissue mass by enabling and recruiting new adipocytes more sensitive to insulin and lipid storage and disabling and clearing mature adipocytes with saturated lipid vacuoles and less sensitive to insulin.

PPAR γ and endocrine function of the adipose tissue

PPAR γ is not only a key factor controlling adipogenesis and adipose tissue mass control but also serves as the master regulator of metabolic genes in this tissue (Figure 2). This activity conferred the definition of an active endocrine gland to the adipose tissue allowing it to secrete a wide range of bioactive substances called adipokines [2,66,67] actively implicated in the regulation of glucose and lipid homeostasis [17,31,53]. By governing adipokines production through ligand systemic availability in the body (natural ligand, TZD, etc.), PPAR γ improves insulin sensitivity both at adipocyte, muscular and hepatic levels by stimulation of adipogenesis, increasing muscle glucose and FFA consumption, inhibiting the hepatic glycolysis and decreasing the release of FFA in the blood [31,68,69]. The most important determinant of amount and nature of adipokine secreted by the adipose tissue is the nature of the ligand that stimulates PPAR γ activity, the number of adipocytes contained in the adipose tissue and their size [24,70,71].

The first hormone to be identified as adipokine was leptin discovered in 1994 [10]. Leptin is an adipokine secreted exclusively by mature adipocytes and its plasma level is positively correlated with body fat mass [70,72]. During a meal, increased systemic FFA level are positively correlated with PPAR γ activity that leads to increased leptin secretion [73]. Leptin seems to be secreted by mature adipocytes as a negative retrocontrol on PPAR γ activity [74-76] to limit the adipose tissue over-expansion. However, this adipokine has a central role in glucose homeostasis acting on several organs. Leptin act on the sympathetic nervous system to regulate satiety [77,78], inhibit insulin secretion from pancreatic β -cells [79,80] and decrease insulin receptor sensitivity in the peripheral cells to limit glucose uptake and lipid overload [81,82]. Leptin limits also adipose tissue expansion by increasing TNF α production [83], also known to decrease insulin systemic sensitivity [84-86], to inhibit adipogenesis and to decrease the fat storage [86-89]. On the other hand, leptin activates the 5'-AMP-activated protein kinase (AMPK) in target tissues [90,91]. This kinase stimulates the oxidation of FFA by inhibiting the activity of acetylcoenzyme A carboxylase governing the production of the

enzyme malonyl-CoA responsible for lipogenesis [92]. As a result, leptin prevents fat accumulation in peripheral tissues and prevent lipotoxicity. Thus, increased caloric intake will cause PPAR γ activity to be higher and leptin secretion to elevate, leading to a leptin systemic resistance [93,94] that will be developed into systemic insulin resistance and diabetes.




Figure 2: PPAR γ has multiple direct and indirect, through adipokine secretion, effects on glucose and lipid homeostasis as well as on several physiologic organ function.

PPAR γ management of metabolic homeostasis passes also through the regulation of the adiponectin secretion, an insulin-sensitizing adipokine [95,96] (Figure 2). In contrast to leptin, plasma concentrations decrease with weight gain and obesity and hypo-adiponectinemia is correlated with insulin-resistant diabetes [70,97]. This counterbalance between leptin and adiponectin is not fully elucidated, but a possible action of leptin on the adiponectin expression level in order to limit adipose tissue expansion could be the answer as seen with PPAR γ activity leptin-control. This mechanism could be plausible in the perspective of the main leptin action working on reduction of systemic lipid overload. In fact, adiponectin work on improving insulin sensitivity and controls FFA oxidation through the activation of AMPK [98,99] leading to a decrease in muscles and liver lipid accumulation and an improvement of the signal transduction of insulin. Adiponectin also increases the translocation of GLUT4 transporters from the cytoplasm to the plasma membrane to facilitate the glucose uptake [100]. Adiponectin also improves blood glucose by inhibiting the expression of mRNA coding for hepatic G6Pase and PEPCK, which has the effect of reducing the production of glucose [98,101]. Finally, adiponectin abolish or reprehend the leptin-induced TNF α secretion improving thus the insulin sensitivity [102]. Thus, by increasing adiponectin level [103,104], PPAR γ work in concert with insulin receptors to improve systemic glycemia by improving insulin sensitivity and glucose tolerance. However, a possible role of increased leptin level in the blood, correlated with obesity, could be the main cause of decreased glyemic parameters correlated with decreased adiponectin levels and insulin resistance. Furthermore, resistin, a relatively new adipokine that is gaining in importance in research for its implication in glucose homeostasis [105], directly impact adiponectin action. Its secretion profile is also governed by PPAR γ activity [106,107] and its circulating levels are increased with obesity inducing insulin resistance and an impaired glucose tolerance

[108-110]. The implication of resistin in diabetogenesis remains very controversial and several studies showed that resistin couldn't be implicated in metabolic diseases [111,112]. These contradictory data could be due to the fact that resistin is very weakly expressed in adipose tissue in humans and the small amounts found in adipose tissue are originate from macrophages [113,114]. More investigations should be pursued on this adipokine.

The list of the PPAR γ regulated adipokine is expanding day after day and regrouping more family subtype [71]. Among them the exclusive adipose tissue adipokine vaspin [115,116], the insulin-like adipokine visfatin [117,118] and the insulin and TNF α -stimulated adipokine apelin [119,120] are recent adipokine secreted during PPAR γ activation. These adipokines, like adiponectin, are increased in weight gain to counteract the development of insulin resistance and improve glucose intolerance, but decreases with the progression of diabetes and obesity. Further, the lipogenic adipokine ASP (Acylation Stimulating Protein) [121], the Fasting Induced Adipose Factor (FIAF) [122] are all adipokines that their genes contain a PPRE sequence and their expression is therefore controlled by PPAR γ activity. These adipokines play key role in glucose homeostasis by improving systemic insulin sensitivity and lipid metabolism by stimulating hepatic FFA uptake and inhibiting LPL.

PPAR γ -mediating adipokine secretion activity depends on different factors including ligand nature, adipocyte status and systemic metabolic changes. The whole system is settled to control metabolic homeostasis by stimulation of insulin sensitizer adipokines secretion and suppression of other diabetogenic factors. However, the maintaining of harmful exogenous factors supply will unbalance the system towards a decrease of PPAR γ activity that becomes dangerous for systemic homeostasis, to limit unlimited adipose tissue expansion and glucose storage; all together will lead to metabolic disease development.

PPAR γ and cardiovascular alterations

Besides regulating numerous metabolic pathways, PPAR γ also governs cardiovascular processes linked to their homeostasis (Figures 2 and 3). The expression of PPAR γ has been shown in many cardiovascular cell types including monocytes and macrophages [38], smooth muscle cells [123] and endothelial cells [124]. The first clue towards cardio-protective effects of PPAR γ came from observation of the cardiomyocyte-specific PPAR γ knock out mice that exhibit a cardiac hypertrophy [125]. The cardioprotective effects of PPAR γ in these mice are likely explained by the decrease of the glucose tolerance in the cardiomyocyte leading them to consume FFA as a source of energy but more susceptible to induce ROS secretion. These later are known to affect endothelial function and increase cardiac inflammation [126,127]. Various evidences have shown that PPAR γ exerts an inhibitory effect on NF- κ B [128,129], a nuclear factor involved in the transcription of many genes encoding inflammatory proteins. As a result, NF- κ B is retained in a nonactive form leading to suppression of its transcriptional activity for inflammatory factors such as type 1 receptor of bradykinin [130] whose expression is increased in inflammation and in diabetes. The anti-inflammatory effects of PPAR γ could also be explained by the inhibitory effect of this nuclear receptor on signal transducer and activator of transcription (Stat) and Activator protein-1 (AP-1) [129,131]. Here also, PPAR γ act to inhibit the transcriptional activity of these inflammatory factors leading to a decreased impact of these molecules on systemic hypertension induction [129]. Furthermore, PPAR γ action on

proliferation potential of vascular smooth muscle cells prevents hypertension events through adipokine secretions [24,123,132,133].




Figure 3: In obesity-linked type 2 diabetes, characterized by increased lipid and glucose circulating levels in the body, adipose tissue secretions profile actively contribute to the development of diabetic cardiomyopathy. In addition to the negative influence of glucose and lipid disturbed homeostasis on cardiac function, modulation of the adipose tissue activity, illustrated by an increase PPAR γ activity, by the hyperlipidemia and hyperglycaemia directly influence the cardiac physiology through its adipokine secretion profile modulation.

It has been well established that TZD-PPAR γ activation decreases the production of cytokines (TNF α , IL-1, IL-6, IL-18, CRP, etc.) by the macrophages [134], a potential anti-inflammatory effect. The glitazones could induce the transformation of macrophages into foam cells in the atherosclerotic lesions, but the results are somewhat contradictory depending on the chosen experimental system [135,136]. They inhibit the proliferation and migration of vascular smooth cells [123,137]. Finally, in endothelial cells in culture, TZD induce the expression of Plasminogen Activator Inhibitor -1 (PAI-1, an inhibitor of fibrinolysis) [138], whereas in diabetic patients treated with troglitazone reduces the concentration of PAI-1 circulating [139]. Such contradictory experimental observations led to several questions to determine if glitazones treatment is benefic or not for cardiovascular system.

Even if it has shown overt improvement in fasting glucose and insulin sensitivity, TZD treatment has worsened cardiac parameters in the normal, diabetic and transgenic rodent models [125,127,140,141]. PPAR γ pharmacological activation also increased incidence of congestive heart failure and induce fluid retention as reported in the RECORD European clinical trial [142,143]. These data led to the withdrawal from the US and European markets of Rosiglitazone in 2010 and to the emission of black box warning on the other member of the TZD family.

Known as an agonist of PPAR γ , TZD could induce FFA scavenger receptors expression in vascular cells wall leading to the development of foam cells and trigger thus atherosclerotic formation [24,135]. PPAR γ activation also increases TNF α secretion that plays an autocrine role on the adipose tissue to inhibit adipogenesis and decrease fat storage [65]. However, TNF α exert its effects on other organs like heart and vascular cells leading to insulin resistance [144], which also leads to an increased oxidative stress and cell dysfunctions.

Furthermore, TNF α down regulates the PPAR α activity in cardiomyocytes [145] leading to a worsen status for these cells.

TNF α upregulate apelin expression levels [119], an adipokine known for its cardiotoxic effects [146] that could lead to a heart failure. The apelin also controls blood pressure and heart activity due to its direct action on the cardiovascular system and its action on the autonomic nervous system [147-149] and an overexpression of this hormone could lead to several heart complications.

Other adipokines catalyze the effect of PPAR γ activity on the cardiovascular system. Among them leptin and adiponectin are the most studied adipokines for this issue. Leptin is known to regulate the hypothalamic-pituitary-adrenal axis responsible for blood pressure regulation [77]. In subjects suffering of metabolic diseases this control is disturbed if not missing. Thus, high leptin concentration leads to diastolic dysfunction associated with higher cardiac sympathetic nervous system activity and increased left ventricle mass [150]. This dysfunction with a reduction in cardiac compliance is thus associated with left ventricle dilatation and an increased left ventricle mass in obesity-linked diabetic mice [70,151]. The leptin receptor Ob-R belongs to the cytokine receptor family class I [152,153] that include interleukins and growth hormone receptors. This suggests other possible biological effects of leptin such as inflammation associated with its related cytokine nature [154-156]. In addition, leptin stimulates the synthesis of ET-1 [157,158], NOS [159], ROS production [158] and expression of MCP-1 [160] that have a direct impact on the increasing of the oxidative stress in endothelial cells. All these factors could lead to the atherogenesis process.

Further, leptin has angiogenic activity and promotes migration and proliferation of vascular smooth muscle cells [24,161]. This effect is important in the physiological process of the expansion of adipose tissue that requires a good blood and oxygen supply. Leptin also promotes FFA oxidation, glucose uptake, platelet aggregation and accumulation of cholesterol in macrophages involved in atherogenesis effect [90,156,162,163].

In parallel, decreased level of adiponectin in obesity and diabetes is correlated with hypertension, presence of coronary heart disease and diabetic cardiomyopathy [70,151,164,165]. Its protective properties against atherosclerosis pass through its inhibitory effects on the expression of adhesion molecules on endothelial cells limiting the recruitment of monocytes to the vascular wall, and by its anti-inflammatory properties which inhibit the production of TNF α and macrophages activity [102,166-169]. This adipokine also inhibits smooth muscle cells proliferation by inhibiting the proliferative effects of PDGF [24,132].

Finally, cathepsins are adipokines with protease activity of the papain family actively involved in protein metabolism [170-173]. This family includes several members, including cathepsins S, K and L. They are secreted by adipose tissue in parallel with food intake and leptin secretion. They directly affect adipocyte differentiation and remodeling of the endothelial cells. This action of matrix remodeling (Degradation of collagen, elastin, fibronectin, etc.) is essential for adipogenesis and adipose tissue expansion; their secretion are thus increased in obese person and possibly implicated in several cardiovascular alterations [174].

All together, these data suggest that the modulation of the expression of PPAR γ receptors in the diabetic or TZD treated mice might be a pharmacological model of two situations: a permanent blocking of these receptors as a response of the organism for obesity to

prevent diabetes or desensitization of this nuclear receptor due to continuous stimulation with its ligand during treatment.

Conclusion

This year, we are celebrating the 20th anniversary of the discovery of the first adipokine that gave us a novel definition for adipose tissue: leptin. Since then it has become clear that adipose tissue is a source of a wide range of bioactive molecules called adipokine leading the regulation of systemic metabolism. The secretion of these hormones is henceforth controlled by what we can call nowadays the masterchef of metabolic homeostasis: PPAR γ .

Being modulated by a considerable variety of endogenous and synthetic ligand, PPAR γ is at the present time considered as a crucial metabolic sensor modulating numerous gene expression implicated in body homeostasis. Although PPAR γ has mostly been connected with glycemic modulation, it is now evident that its effects are much more extensive and cover adipogenesis, cardiometabolic control and lipid catabolism. One of the major challenges lying ahead remains to better understand the molecular mechanism underlying its modulated activity related to the ligand nature, to improve our knowledge of its specificity to the chosen ligand in each therapeutic treatment.

References

1. Issemann I, Green S (1990) Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 347: 645-650.
2. Siiteri PK (1987) Adipose tissue as a source of hormones. *Am J Clin Nutr* 45: 277-282.
3. Kliewer SA, Sundseth SS, Jones SA, Brown PJ, Wisely GB, et al. (1997) Fatty acids and eicosanoids regulate gene expression through direct interactions with peroxisome proliferator-activated receptors alpha and gamma. *Proc Natl Acad Sci U S A* 94: 4318-4323.
4. Frick MH, Elo O, Haapa K, Heinonen OP, Heinsalmi P, et al. (1987) Helsinki Heart Study: primary-prevention trial with gemfibrozil in middle-aged men with dyslipidemia. Safety of treatment, changes in risk factors, and incidence of coronary heart disease. *N Engl J Med* 317: 1237-1245.
5. Issemann I, Prince RA, Tugwood JD, Green S (1993) The peroxisome proliferator-activated receptor:retinoid X receptor heterodimer is activated by fatty acids and fibrate hypolipidaemic drugs. *J Mol Endocrinol* 11: 37-47.
6. Tyagi S, Gupta P, Saini AS, Kaushal C, Sharma S (2011) The peroxisome proliferatorK activated receptor: A family of nuclear receptors role in various diseases. *J Adv Pharm Technol Res* 2: 236-240.
7. Aldridge TC, Tugwood JD, Green S (1995) Identification and characterization of DNA elements implicated in the regulation of CYP4A1 transcription. *Biochem J* 306: 473-479.
8. Muerhoff AS, Griffin KJ, Johnson EF (1992) The peroxisome proliferator-activated receptor mediates the induction of CYP4A6, a cytochrome P450 fatty acid omega-hydroxylase, by clofibrilic acid. *J Biol Chem* 267: 19051-19053.
9. Tugwood JD, Issemann I, Anderson RG, Bundell KR, McPheat WL, et al. (1992) The mouse peroxisome proliferator activated receptor recognizes a response element in the 5' flanking sequence of the rat acyl CoA oxidase gene. *EMBO J* 11: 433-439.
10. Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, et al. (1994) Positional cloning of the mouse obese gene and its human homologue. *Nature* 372: 425-432.
11. Zhu Y, Alvares K, Huang Q, Rao MS, Reddy JK (1993) Cloning of a new member of the peroxisome proliferator-activated receptor gene family from mouse liver. *J Biol Chem* 268: 26817-26820.
12. Aranda A, Pascual A (2001) Nuclear hormone receptors and gene expression. *Physiol Rev* 81: 1269-1304.
13. Desvergne B, Wahli W (1999) Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocr Rev* 20: 649-688.
14. Ferré P (2004) The biology of peroxisome proliferator-activated receptors: relationship with lipid metabolism and insulin sensitivity. *Diabetes* 53 Suppl 1: S43-50.
15. Laudet V (1997) Evolution of the nuclear receptor superfamily: early diversification from an ancestral orphan receptor. *J Mol Endocrinol* 19: 207-226.
16. Lehmann JM, Moore LB, Smith-Oliver TA, Wilkison WO, Willson TM, et al. (1995) An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor gamma (PPAR gamma). *J Biol Chem* 270: 12953-12956.
17. Spiegelman BM (1998) PPAR-gamma: adipogenic regulator and thiazolidinedione receptor. *Diabetes* 47: 507-514.
18. Sundvold H, Lien S (2001) Identification of a novel peroxisome proliferator-activated receptor (PPAR) gamma promoter in man and transactivation by the nuclear receptor RORalpha1. *Biochem Biophys Res Commun* 287: 383-390.
19. Dreyer C, Keller H, Mahfoudi A, Laudet V, Krey G, et al. (1993) Positive regulation of the peroxisomal beta-oxidation pathway by fatty acids through activation of peroxisome proliferator-activated receptors (PPAR). *Biol Cell* 77: 67-76.
20. Houseknecht KL, Cole BM, Steele PJ (2002) Peroxisome proliferator-activated receptor gamma (PPARgamma) and its ligands: a review. *Domest Anim Endocrinol* 22: 1-23.
21. Schoonjans K, Peinado-Onsurbe J, Lefebvre AM, Heyman RA, Briggs M, et al. (1996) PPARalpha and PPARgamma activators direct a distinct tissue-specific transcriptional response via a PPRE in the lipoprotein lipase gene. *EMBO J* 15: 5336-5348.
22. Xu HE, Lambert MH, Montana VG, Parks DJ, Blanchard SG, et al. (1999) Molecular recognition of fatty acids by peroxisome proliferator-activated receptors. *Mol Cell* 3: 397-403.
23. Willermain F, Dulku S, Gonzalez NS, Blero D, Driessens G, et al. (2006) 15KDeoxyK12,14Kprostaglandin J2 inhibits interferon gamma induced MHC class II but not class I expression on ARPE cells through a PPAR gamma independent mechanism. *Prostaglandins Other Lipid Mediat* 80: 136-143.
24. El Akoum S, Cloutier I, Tanguay JF (2012) Vascular smooth muscle cell alterations triggered by mice adipocytes: role of high-fat diet. *J Atheroscler Thromb* 19: 1128-1141.
25. Fan YY, Ramos KS, Chapkin RS (2001) Dietary gamma-linolenic acid suppresses aortic smooth muscle cell proliferation and modifies atherosclerotic lesions in apolipoprotein E knockout mice. *J Nutr* 131: 1675-1681.
26. Harding AH, Day NE, Khaw KT, Bingham S, Luben R, et al. (2004) Dietary fat and the risk of clinical type 2 diabetes: the European prospective investigation of Cancer-Norfolk study. *Am J Epidemiol* 159: 73-82.
27. Herberg L, Döppen W, Major E, Gries FA (1974) Dietary-induced hypertrophic--hyperplastic obesity in mice. *J Lipid Res* 15: 580-585.
28. Kubota N, Terauchi Y, Miki H, Tamemoto H, Yamauchi T, et al. (1999) PPAR gamma mediates high-fat diet-induced adipocyte hypertrophy and insulin resistance. *Mol Cell* 4: 597-609.
29. Lee YI, Wang MY, Kakuma T, Wang ZW, Babcock E, et al. (2001) Liporegulation in diet-induced obesity. The antisteatotic role of hyperleptinemia. *J Biol Chem* 276: 5629-5635.
30. Manco M, Calvani M, Mingrone G (2004) Effects of dietary fatty acids on insulin sensitivity and secretion. *Diabetes Obes Metab* 6: 402-413.
31. Wilmsen HM, Ciaraldi TP, Carter L, Reehman N, Mudaliar SR, et al. (2003) Thiazolidinediones upregulate impaired fatty acid uptake in skeletal muscle of type 2 diabetic subjects. *Am J Physiol Endocrinol Metab* 285: E354-362.

32. Braissant O, Foufelle F, Scotto C, Dauça M, Wahli W (1996) Differential expression of peroxisome proliferator-activated receptors (PPARs): tissue distribution of PPAR-alpha, -beta, and -gamma in the adult rat. *Endocrinology* 137: 354-366.
33. Mukherjee R, Jow L, Noonan D, McDonnell DP (1994) Human and rat peroxisome proliferator activated receptors (PPARs) demonstrate similar tissue distribution but different responsiveness to PPAR activators. *J Steroid Biochem Mol Biol* 51: 157-166.
34. Werman A, Hollenberg A, Solanes G, Bjorbaek C, Vidal-Puig AJ, et al. (1997) Ligand-independent activation domain in the N terminus of peroxisome proliferator-activated receptor gamma (PPARgamma). Differential activity of PPARgamma1 and -2 isoforms and influence of insulin. *J Biol Chem* 272: 20230-20235.
35. Medina-Gomez G, Gray SL, Yetukuri L, Shimomura K, Virtue S, et al. (2007) PPAR gamma 2 prevents lipotoxicity by controlling adipose tissue expandability and peripheral lipid metabolism. *PLoS Genet* 3: e64.
36. Zhu Y, Qi C, Korenberg JR, Chen XN, Noya D, et al. (1995) Structural organization of mouse peroxisome proliferator-activated receptor gamma (mPPAR gamma) gene: alternative promoter use and different splicing yield two mPPAR gamma isoforms. *Proc Natl Acad Sci USA* 92: 7921-7925.
37. Bortolotto JW, Margis R, Ferreira AC, Padoin AV, Mottin CC, et al. (2007) Adipose tissue distribution and quantification of PPARbeta/delta and PPARgamma1K3 mRNAs: discordant gene expression in subcutaneous, retroperitoneal and visceral adipose tissue of morbidly obese patients. *Obes Surg* 17: 934-9340.
38. Tontonoz P, Nagy L, Alvarez JG, Thomazy VA, Evans RM (1998) PPARgamma promotes monocyte/macrophage differentiation and uptake of oxidized LDL. *Cell* 93: 241-252.
39. Al-Shali K, Cao H, Knoers N, Hermus AR, Tack CJ, et al. (2004) A single-base mutation in the peroxisome proliferator-activated receptor gamma4 promoter associated with altered in vitro expression and partial lipodystrophy. *J Clin Endocrinol Metab* 89: 5655-5660.
40. Fajas L, Fruchart JC, Auwerx J (1998) PPARgamma3 mRNA: a distinct PPARgamma mRNA subtype transcribed from an independent promoter. *FEBS Lett* 438: 55-60.
41. Jiang C, Ting AT, Seed B (1998) PPAR-gamma agonists inhibit production of monocyte inflammatory cytokines. *Nature* 391: 82-86.
42. Marx N, Sukhova GK, Collins T, Libby P, Plutzky J (1999) PPARalpha activators inhibit cytokine-induced vascular cell adhesion molecule-1 expression in human endothelial cells. *Circulation* 99: 3125-3131.
43. Muraio K, Imachi H, Momoi A, Sayo Y, Hosokawa H, et al. (1999) Thiazolidinedione inhibits the production of monocyte chemoattractant protein-1 in cytokine-treated human vascular endothelial cells. *FEBS Lett* 454: 27-30.
44. Ricote M, Huang J, Fajas L, Li A, Welch J, et al. (1998) Expression of the peroxisome proliferator-activated receptor gamma (PPARgamma) in human atherosclerosis and regulation in macrophages by colony stimulating factors and oxidized low density lipoprotein. *Proc Natl Acad Sci U S A* 95: 7614-7619.
45. Francis GA, Annicotte JS, Auwerx J (2003) PPAR agonists in the treatment of atherosclerosis. *Curr Opin Pharmacol* 3: 186-191.
46. Chinetti G, Griglio S, Antonucci M, Torra IP, Delerive P, et al. (1998) Activation of proliferator-activated receptors alpha and gamma induces apoptosis of human monocyte-derived macrophages. *J Biol Chem* 273: 25573-25580.
47. Goto T, Lee JY, Teraminami A, Kim YI, Hirai S, et al. (2011) Activation of peroxisome proliferator-activated receptor-alpha stimulates both differentiation and fatty acid oxidation in adipocytes. *J Lipid Res* 52: 873-884.
48. Lee JY, Hashizaki H, Goto T, Sakamoto T, Takahashi N, et al. (2011) Activation of peroxisome proliferator-activated receptor- α enhances fatty acid oxidation in human adipocytes. *Biochem Biophys Res Commun* 407: 818-822.
49. Mottillo EP, Bloch AE, Leff T, Granneman JG (2012) Lipolytic products activate peroxisome proliferator-activated receptor (PPAR) α and β in brown adipocytes to match fatty acid oxidation with supply. *J Biol Chem* 287: 25038-25048.
50. Valmaseda A, Carmona MC, Barberá MJ, Viñas O, Mampel T, et al. (1999) Opposite regulation of PPAR-alpha and -gamma gene expression by both their ligands and retinoic acid in brown adipocytes. *Mol Cell Endocrinol* 154: 101-109.
51. Hansen JB, Zhang H, Rasmussen TH, Petersen RK, Flindt EN, et al. (2001) Peroxisome proliferator-activated receptor delta (PPARdelta)-mediated regulation of preadipocyte proliferation and gene expression is dependent on cAMP signaling. *J Biol Chem* 276: 3175-3182.
52. Tontonoz P, Hu E, Graves RA, Budavari AI, Spiegelman BM (1994) mPPAR gamma 2: tissue-specific regulator of an adipocyte enhancer. *Genes Dev* 8: 1224-1234.
53. Tontonoz P, Hu E, Spiegelman BM (1994) Stimulation of adipogenesis in fibroblasts by PPAR gamma 2, a lipid-activated transcription factor. *Cell* 79: 1147-1156.
54. Tontonoz P, Hu E, Devine J, Beale EG, Spiegelman BM (1995) PPAR gamma 2 regulates adipose expression of the phosphoenolpyruvate carboxykinase gene. *Mol Cell Biol* 15: 351-357.
55. Xiao Y, Yuan T, Yao W, Liao K (2010) 3T3-L1 adipocyte apoptosis induced by thiazolidinediones is peroxisome proliferator-activated receptor-gamma-dependent and mediated by the caspase-3-dependent apoptotic pathway. *FEBS J* 277: 687-696.
56. Agarwal AK, Garg A (2002) A novel heterozygous mutation in peroxisome proliferator-activated receptor-gamma gene in a patient with familial partial lipodystrophy. *J Clin Endocrinol Metab* 87: 408-411.
57. Lütke A, Buettner J, Wu W, Muchir A, Schroeter A, et al. (2007) Peroxisome proliferator-activated receptor-gamma C190S mutation causes partial lipodystrophy. *J Clin Endocrinol Metab* 92: 2248-2255.
58. Barroso I, Gurnell M, Crowley VE, Agostini M, Schwabe JW, et al. (1999) Dominant negative mutations in human PPARgamma associated with severe insulin resistance, diabetes mellitus and hypertension. *Nature* 402: 880-883.
59. Savage DB, Tan GD, Acerini CL, Jebb SA, Agostini M, et al. (2003) Human metabolic syndrome resulting from dominant-negative mutations in the nuclear receptor peroxisome proliferator-activated receptor-gamma. *Diabetes* 52: 910-917.
60. Rosen ED, Sarraf P, Troy AE, Bradwin G, Moore K, et al. (1999) PPAR gamma is required for the differentiation of adipose tissue in vivo and in vitro. *Mol Cell* 4: 611-617.
61. Barak Y, Nelson MC, Ong ES, Jones YZ, Ruiz-Lozano P, et al. (1999) PPAR gamma is required for placental, cardiac, and adipose tissue development. *Mol Cell* 4: 585-595.
62. Sugii S, Olson P, Sears DD, Saberi M, Atkins AR, et al. (2009) PPARgamma activation in adipocytes is sufficient for systemic insulin sensitization. *Proc Natl Acad Sci U S A* 106: 22504-22509.
63. Ryan KK, Li B, Grayson BE, Matter EK, Woods SC, et al. (2011) A role for central nervous system PPAR- β in the regulation of energy balance. *Nat Med* 17: 623-626.
64. Lu M, Sarruf DA, Talukdar S, Sharma S, Li P, et al. (2011) Brain PPAR- β promotes obesity and is required for the insulin-sensitizing effect of thiazolidinediones. *Nat Med* 17: 618-622.
65. Keuper M, Wernstedt Asterholm I, Scherer PE, Westhoff MA, Möller P, et al. (2013) TRAIL (TNF-related apoptosis-inducing ligand) regulates adipocyte metabolism by caspase-mediated cleavage of PPARgamma. *Cell Death Dis* 4: e474.
66. Trayhurn P, Bing C, Wood IS (2006) Adipose tissue and adipokines--energy regulation from the human perspective. *J Nutr* 136: 1935S-1939S.
67. Rabe K, Lehrke M, Parhofer KG, Broedl UC (2008) Adipokines and insulin resistance. *Mol Med* 14: 741-751.
68. Basciano H, Miller AE, Naples M, Baker C, Kohen R, et al. (2009) Metabolic effects of dietary cholesterol in an animal model of insulin resistance and hepatic steatosis. *Am J Physiol Endocrinol Metab* 297: E462-473.

69. Ahmadian M, Suh JM, Hah N, Liddle C, Atkins AR, et al. (2013) PPAR δ signaling and metabolism: the good, the bad and the future. *Nat Med* 19: 557-566.
70. El Akoum S, Lamontagne V, Cloutier I, Tanguay JF (2011) Nature of fatty acids in high fat diets differentially delineates obesity-linked metabolic syndrome components in male and female C57BL/6J mice. *Diabetol Metab Syndr* 3: 34.
71. Mirzaei K, Hossein-Nezhad A, Keshavarz SA, Koohdani F, SaboorYaraghi AA, et al. (2013) Crosstalk between circulating peroxisome proliferator-activated receptor gamma, adipokines and metabolic syndrome in obese subjects. *Diabetology & metabolic syndrome* 5: 79.
72. Frederich RC, Hamann A, Anderson S, Löllmann B, Lowell BB, et al. (1995) Leptin levels reflect body lipid content in mice: evidence for diet-induced resistance to leptin action. *Nat Med* 1: 1311-1314.
73. Hollenberg AN, Susulic VS, Madura JP, Zhang B, Moller DE, et al. (1997) Functional antagonism between CCAAT/Enhancer binding protein-alpha and peroxisome proliferator-activated receptor-gamma on the leptin promoter. *J Biol Chem* 272: 5283-5290.
74. Cabrero A, Cubero M, Llaverias G, Alegret M, Sánchez R, et al. (2005) Leptin down-regulates peroxisome proliferator-activated receptor gamma (PPAR-gamma) mRNA levels in primary human monocyte-derived macrophages. *Mol Cell Biochem* 275: 173-179.
75. Wang L, Shao YY, Ballock RT (2012) Leptin Antagonizes Peroxisome Proliferator-Activated Receptor- δ Signaling in Growth Plate Chondrocytes. *PPAR Res* 2012: 756198.
76. Zhou Y, Jia X, Qin J, Lu C, Zhu H, et al. (2010) Leptin inhibits PPARgamma gene expression in hepatic stellate cells in the mouse model of liver damage. *Mol Cell Endocrinol* 323: 193-200.
77. Correia ML, Morgan DA, Sivitz WI, Mark AL, Haynes WG (2001) Leptin acts in the central nervous system to produce dose-dependent changes in arterial pressure. *Hypertension* 37: 936-942.
78. McGarry JD (1995) Appetite control: Does leptin lighten the problem of obesity? *Curr Biol* 5: 1342-1344.
79. Seufert J (2004) Leptin effects on pancreatic beta-cell gene expression and function. *Diabetes* 53 Suppl 1: S152-158.
80. Seufert J, Kieffer TJ, Leech CA, Holz GG, Moritz W, et al. (1999) Leptin suppression of insulin secretion and gene expression in human pancreatic islets: implications for the development of adipogenic diabetes mellitus. *J Clin Endocrinol Metab* 84: 670-676.
81. Benomar Y, Wetzler S, Larue-Achagiotis C, Djiane J, Tomé D, et al. (2005) In vivo leptin infusion impairs insulin and leptin signalling in liver and hypothalamus. *Mol Cell Endocrinol* 242: 59-66.
82. Unger RH (2005) Hyperleptinemia: protecting the heart from lipid overload. *Hypertension* 45: 1031-1034.
83. Shen J, Sakaida I, Uchida K, Terai S, Okita K (2005) Leptin enhances TNF-alpha production via p38 and JNK MAPK in LPS-stimulated Kupffer cells. *Life Sci* 77: 1502-1515.
84. Nieto-Vazquez I, Fernández-Veledo S, Krämer DK, Vila-Bedmar R, Garcia-Guerra L, et al. (2008) Insulin resistance associated to obesity: the link TNF-alpha. *Arch Physiol Biochem* 114: 183-194.
85. Liu LS, Spelleken M, Röhrig K, Hauner H, Eckel J (1998) Tumor necrosis factor-alpha acutely inhibits insulin signaling in human adipocytes: implication of the p80 tumor necrosis factor receptor. *Diabetes* 47: 515-522.
86. Qi C, Pekala PH (2000) Tumor necrosis factor-alpha-induced insulin resistance in adipocytes. *Proc Soc Exp Biol Med* 223: 128-135.
87. Cawthorn WP, Heyd F, Hegyi K, Sethi JK (2007) Tumour necrosis factor-alpha inhibits adipogenesis via a beta-catenin/TCF4(TCF7L2)-dependent pathway. *Cell Death Differ* 14: 1361-1373.
88. Cawood TJ, Moriarty P, O'Farrelly C, O'Shea D (2006) The effects of tumour necrosis factor-alpha and interleukin1 on an in vitro model of thyroid-associated ophthalmopathy; contrasting effects on adipogenesis. *Eur J Endocrinol* 155: 395-403.
89. Warne JP (2003) Tumour necrosis factor alpha: a key regulator of adipose tissue mass. *J Endocrinol* 177: 351-355.
90. Minokoshi Y, Kim YB, Peroni OD, Fryer LG, Müller C, et al. (2002) Leptin stimulates fatty-acid oxidation by activating AMP-activated protein kinase. *Nature* 415: 339-343.
91. Suzuki A, Okamoto S, Lee S, Saito K, Shiuchi T, et al. (2004) Leptin stimulates fatty acid oxidation and peroxisome proliferator-activated receptor alpha gene expression in mouse C2C12 myoblasts by changing the subcellular localization of the alpha2 form of AMPK activated protein kinase. *Mol Cell Biol* 27: 4317K27.
92. Park SH, Gammon SR, Knippers JD, Paulsen SR, Rubink DS, et al. (2002) Phosphorylation-activity relationships of AMPK and acetyl-CoA carboxylase in muscle. *J Appl Physiol* (1985) 92: 2475-2482.
93. Banks WA, Coon AB, Robinson SM, Moinuddin A, Shultz JM, et al. (2004) Triglycerides induce leptin resistance at the blood-brain barrier. *Diabetes* 53: 1253-1260.
94. Steinberg GR, Parolin ML, Heigenhauser GJ, Dyck DJ (2002) Leptin increases FA oxidation in lean but not obese human skeletal muscle: evidence of peripheral leptin resistance. *Am J Physiol Endocrinol Metab* 283: E187-192.
95. Maeda K, Okubo K, Shimomura I, Funahashi T, Matsuzawa Y, et al. (1996) cDNA cloning and expression of a novel adipose specific collagen-like factor, apM1 (AdiPose Most abundant Gene transcript 1). *Biochem Biophys Res Commun* 221: 286-289.
96. Astapova O, Leff T (2012) Adiponectin and PPAR δ : cooperative and interdependent actions of two key regulators of metabolism. *Vitam Horm* 90: 143-162.
97. Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, et al. (1999) Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun* 257: 79-83.
98. Yamauchi T, Kamon J, Minokoshi Y, Ito Y, Waki H, et al. (2002) Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat Med* 8: 1288-1295.
99. Fu Y, Luo N, Klein RL, Garvey WT (2005) Adiponectin promotes adipocyte differentiation, insulin sensitivity, and lipid accumulation. *J Lipid Res* 46: 1369-1379.
100. Ceddia RB, Somwar R, Maida A, Fang X, Bikopoulos G, et al. (2005) Globular adiponectin increases GLUT4 translocation and glucose uptake but reduces glycogen synthesis in rat skeletal muscle cells. *Diabetologia* 48: 132-139.
101. Bulló M, Salas-Salvadó J, García-Lorda P (2005) Adiponectin expression and adipose tissue lipolytic activity in lean and obese women. *Obes Surg* 15: 382-386.
102. Zhao T, Hou M, Xia M, Wang Q, Zhu H, et al. (2005) Globular adiponectin decreases leptin-induced tumor necrosis factor-alpha expression by murine macrophages: involvement of cAMP-PKA and MAPK pathways. *Cell Immunol* 238: 19-30.
103. Lakota K, Wei J, Carns M, Hinchcliff M, Lee J, et al. (2012) Levels of adiponectin, a marker for PPAR-gamma activity, correlate with skin fibrosis in systemic sclerosis: potential utility as biomarker? *Arthritis Res Ther* 14: R102.
104. Yilmaz MI, Sonmez A, Caglar K, Gok DE, Eyileten T, et al. (2004) Peroxisome proliferator-activated receptor gamma (PPAR-gamma) agonist increases plasma adiponectin levels in type 2 diabetic patients with proteinuria. *Endocrine* 25: 207-214.
105. Steppan CM1, Bailey ST, Bhat S, Brown EJ, Banerjee RR, et al. (2001) The hormone resistin links obesity to diabetes. *Nature* 409: 307-312.
106. Zhou L, Yu X, Meng Q, Li H, Niu C, et al. (2013) Resistin reduces mitochondria and induces hepatic steatosis in mice by the protein kinase C/protein kinase G/p65/PPAR gamma coactivator 1 alpha pathway. *Hepatology* 57: 1384-1393.
107. Patel L, Buckels AC, Kinghorn IJ, Murdock PR, Holbrook JD, et al. (2003) Resistin is expressed in human macrophages and directly regulated by PPAR gamma activators. *Biochem Biophys Res Commun* 300: 472-476.

108. Adegate E (2004) An update on the biology and physiology of resistin. *Cell Mol Life Sci* 61: 2485-2496.
109. Junkin KA, Dyck DJ, Mullen KL, Chabowski A, Thrush AB (2009) Resistin acutely impairs insulin-stimulated glucose transport in rodent muscle in the presence, but not absence, of palmitate. *Am J Physiol Regul Integr Comp Physiol* 296: R944-951.
110. McTernan PG, Fisher FM, Valsamakis G, Chetty R, Harte A, et al. (2003) Resistin and type 2 diabetes: regulation of resistin expression by insulin and rosiglitazone and the effects of recombinant resistin on lipid and glucose metabolism in human differentiated adipocytes. *J Clin Endocrinol Metab* 88: 6098-6106.
111. Nagaev I, Smith U (2001) Insulin resistance and type 2 diabetes are not related to resistin expression in human fat cells or skeletal muscle. *Biochem Biophys Res Commun* 285: 561-564.
112. Utschneider KM, Carr DB, Tong J, Wallace TM, Hull RL, et al. (2005) Resistin is not associated with insulin sensitivity or the metabolic syndrome in humans. *Diabetologia* 48: 2330-2333.
113. Lee TS, Lin CY, Tsai JY, Wu YL, Su KH, et al. (2009) Resistin increases lipid accumulation by affecting class A scavenger receptor, CD36 and ATP-binding cassette transporter-A1 in macrophages. *Life Sci* 84: 97-104.
114. Xu W, Yu L, Zhou W, Luo M (2006) Resistin increases lipid accumulation and CD36 expression in human macrophages. *Biochem Biophys Res Commun* 351: 376-382.
115. Li Q, Chen R, Moriya J, Yamakawa J, Sumino H, et al. (2008) A novel adipocytokine, visceral adipose tissue-derived serine protease inhibitor (vaspin), and obesity. *J Int Med Res* 36: 625-629.
116. Youn BS, Klötting N, Kratzsch J, Lee N, Park JW, et al. (2008) Serum vaspin concentrations in human obesity and type 2 diabetes. *Diabetes* 57: 372-377.
117. Fukuhara A, Matsuda M, Nishizawa M, Segawa K, Tanaka M, et al. (2005) Visfatin: a protein secreted by visceral fat that mimics the effects of insulin. *Science* 307: 426-430.
118. Klötting N, Klötting I (2005) Visfatin: gene expression in isolated adipocytes and sequence analysis in obese WOKW rats compared with lean control rats. *Biochem Biophys Res Commun* 332: 1070-1072.
119. Daviaud D, Boucher J, Gesta S, Dray C, Guigne C, et al. (2006) TNF α up-regulates apelin expression in human and mouse adipose tissue. *FASEB J* 20: 1528-1530.
120. Yue P, Jin H, Aillaud M, Deng AC, Azuma J, et al. (2010) Apelin is necessary for the maintenance of insulin sensitivity. *Am J Physiol Endocrinol Metab* 298: E59-67.
121. Cianflone K, Maslowska M, Sniderman AD (1999) Acylation stimulating protein (ASP), an adipocyte autocrine: new directions. *Semin Cell Dev Biol* 10: 31-41.
122. Kersten S, Mandard S, Tan NS, Escher P, Metzger D, et al. (2000) Characterization of the fasting-induced adipose factor FIAF, a novel peroxisome proliferator-activated receptor target gene. *J Biol Chem* 275: 28488-28493.
123. Law RE, Meehan WP, Xi XP, Graf K, Wuthrich DA, et al. (1996) Troglitazone inhibits vascular smooth muscle cell growth and intimal hyperplasia. *J Clin Invest* 98: 1897-1905.
124. Delerive P, Martin-Nizard F, Chinetti G, Trottein F, Fruchart JC, et al. (1999) Peroxisome proliferator-activated receptor activators inhibit thrombin-induced endothelin-1 production in human vascular endothelial cells by inhibiting the activator protein-1 signaling pathway. *Circ Res* 85: 394-402.
125. Duan SZ, Ivashchenko CY, Russell MW, Milstone DS, Mortensen RM (2005) Cardiomyocyte-specific knockout and agonist of peroxisome proliferator-activated receptor-gamma both induce cardiac hypertrophy in mice. *Circ Res* 97: 372-379.
126. Zhang L, Zalewski A, Liu Y, Mazurek T, Cowan S, et al. (2003) Diabetes-induced oxidative stress and low-grade inflammation in porcine coronary arteries. *Circulation* 108: 472-478.
127. Son NH, Park TS, Yamashita H, Yokoyama M, Huggins LA, et al. (2007) Cardiomyocyte expression of PPARgamma leads to cardiac dysfunction in mice. *J Clin Invest* 117: 2791-2801.
128. Desreumaux P, Philippe D (2003) [Dual function of NF-kappa B system during inflammation]. *Gastroenterol Clin Biol* 27: 1052-1053.
129. Green DE, Sutliff RL, Hart CM (2011) Is peroxisome proliferator-activated receptor gamma (PPAR γ) a therapeutic target for the treatment of pulmonary hypertension? *Pulm Circ* 1: 33-47.
130. Maurer M, Bader M, Bas M, Bossi F, Cicardi M, et al. (2011) New topics in bradykinin research. *Allergy* 66: 1397-1406.
131. Pascual G, Fong AL, Ogawa S, Gamliel A, Li AC, et al. (2005) A SUMOylation-dependent pathway mediates transrepression of inflammatory response genes by PPAR-gamma. *Nature* 437: 759-763.
132. Fuerst M, Taylor CG, Wright B, Tworek L, Zahradka P (2012) Inhibition of smooth muscle cell proliferation by adiponectin requires proteolytic conversion to its globular form. *J Endocrinol* 215: 107-117.
133. Meredith D, Panchatcharam M, Miriyala S, Tsai YS, Morris AJ, et al. (2009) Dominant-negative loss of PPARgamma function enhances smooth muscle cell proliferation, migration, and vascular remodeling. *Arterioscler Thromb Vasc Biol* 29: 465-471.
134. Kim HJ, Kang ES, Kim DJ, Kim SH, Ahn CW, et al. (2007) Effects of rosiglitazone and metformin on inflammatory markers and adipokines: decrease in interleukin-18 is an independent factor for the improvement of homeostasis model assessment-beta in type 2 diabetes mellitus. *Clin Endocrinol (Oxf)* 66: 282-289.
135. Lim HJ, Lee S, Lee KS, Park JH, Jang Y, et al. (2006) PPARgamma activation induces CD36 expression and stimulates foam cell like changes in rVSMCs. *Prostaglandins Other Lipid Mediat* 80: 165-174.
136. Little PJ, Osman N, de Dios ST, Cemerlang N, Ballinger M, et al. (2007) Anti-proliferative activity of oral anti-hyperglycemic agents on human vascular smooth muscle cells: thiazolidinediones (glitazones) have enhanced activity under high glucose conditions. *Cardiovasc Diabetol* 6: 33.
137. Gu MX, Liu XC, Jiang L (2013) [Effect of peroxisome proliferator-activated receptor-gamma on proliferation of airway smooth muscle cells in mice with asthma]. *Zhongguo Dang Dai Er Ke Za Zhi* 15: 583-587.
138. Marx N, Bourcier T, Sukhova GK, Libby P, Plutzky J (1999) PPARgamma activation in human endothelial cells increases plasminogen activator inhibitor type-1 expression: PPARgamma as a potential mediator in vascular disease. *Arterioscler Thromb Vasc Biol* 19: 546-551.
139. Koh H, Akiyama Y, Yui K (2006) Effect of troglitazone on serum gamma-glutamyltransferase activity and plasma t-PA/PAI-1 complex in type 2 diabetic patients. *Arzneimittelforschung* 56: 288-294.
140. Frantz S, Hu K, Widder J, Bayer B, Witzel CC, et al. (2004) Peroxisome proliferator-activated-receptor agonism and left ventricular remodeling in mice with chronic myocardial infarction. *Br J Pharmacol* 141: 9-14.
141. Geng DF, Wu W, Jin DM, Wang JF, Wu YM (2006) Effect of peroxisome proliferator-activated receptor gamma ligand. Rosiglitazone on left ventricular remodeling in rats with myocardial infarction. *Int J Cardiol* 113: 86-91.
142. Ehrenstein V, Hernandez RK, Ulrichsen SP, Rungby J, Lash TL, et al. (2013) Rosiglitazone use and post-discontinuation glycaemic control in two European countries, 2000-2010. *BMJ Open* 3: e003424.
143. Komajda M, McMurray JJ, Beck-Nielsen H, Gomis R, Hanefeld M, et al. (2010) Heart failure events with rosiglitazone in type 2 diabetes: data from the RECORD clinical trial. *Eur Heart J* 31: 824-831.
144. Peraldi P, Hotamisligil GS, Buurman WA, White MF, Spiegelman BM (1996) Tumor necrosis factor (TNF)-alpha inhibits insulin signaling through stimulation of the p55 TNF receptor and activation of sphingomyelinase. *J Biol Chem* 271: 13018-13022.
145. Beier K, Völkl A, Fahimi HD (1997) TNF-alpha downregulates the peroxisome proliferator activated receptor-alpha and the mRNAs encoding peroxisomal proteins in rat liver. *FEBS Lett* 412: 385-387.

146. Ashley EA, Powers J, Chen M, Kundu R, Finsterbach T, et al. (2005) The endogenous peptide apelin potently improves cardiac contractility and reduces cardiac loading in vivo. *Cardiovasc Res* 65: 73-82.
147. O'Carroll AM, Selby TL, Palkovits M, Lolait SJ (2000) Distribution of mRNA encoding B78/apj, the rat homologue of the human APJ receptor, and its endogenous ligand apelin in brain and peripheral tissues. *Biochim Biophys Acta* 1492: 72-80.
148. Sunter D, Hewson AK, Dickson SL (2003) Intracerebroventricular injection of apelin-13 reduces food intake in the rat. *Neurosci Lett* 353: 1-4.
149. Zhong JC, Yu XY, Huang Y, Yung LM, Lau CW, et al. (2007) Apelin modulates aortic vascular tone via endothelial nitric oxide synthase phosphorylation pathway in diabetic mice. *Cardiovasc Res* 74: 388-395.
150. Grassi G, Seravalle G, Quarti-Trevano F, Dell'Oro R, Arenare F, et al. (2009) Sympathetic and baroreflex cardiovascular control in hypertension-related left ventricular dysfunction. *Hypertension* 53: 205-209.
151. Stakos DA, Papaioannou HI, Angelidou I, Mantadakis E, Paraskakis E, et al. (2014) Plasma leptin and adiponectin concentrations correlate with cardiometabolic risk and systemic inflammation in healthy, non-obese children. *J Pediatr Endocrinol Metab* 27: 221-228.
152. Nakashima K, Narazaki M, Taga T (1997) Overlapping and distinct signals through leptin receptor (OB-R) and a closely related cytokine signal transducer, gp130. *FEBS Lett* 401: 49-52.
153. Tartaglia LA, Dembski M, Weng X, Deng N, Culpepper J, et al. (1995) Identification and expression cloning of a leptin receptor, OB-R. *Cell* 83: 1263-1271.
154. Wendremaire M, Mourtialon P, Goirand F, Lirussi F, Barrichon M, et al. (2013) Effects of leptin on lipopolysaccharide-induced remodeling in an in vitro model of human myometrial inflammation. *Biol Reprod* 88: 45.
155. Maya-Monteiro CM, Bozza PT (2008) Leptin and mTOR: partners in metabolism and inflammation. *Cell Cycle* 7: 1713-1717.
156. Martin SS, Qasim A, Reilly MP (2008) Leptin resistance: a possible interface of inflammation and metabolism in obesity-related cardiovascular disease. *J Am Coll Cardiol* 52: 1201-1210.
157. Quehenberger P, Exner M, Sunder-Plassmann R, Ruzicka K, Bieglmayer C, et al. (2002) Leptin induces endothelin-1 in endothelial cells in vitro. *Circ Res* 90: 711-718.
158. Xu FP, Chen MS, Wang YZ, Yi Q, Lin SB, et al. (2004) Leptin induces hypertrophy via endothelin-1-reactive oxygen species pathway in cultured neonatal rat cardiomyocytes. *Circulation* 110: 1269-1275.
159. Otero M, Lago R, Gómez R, Lago F, Gomez-Reino JJ, et al. (2007) Phosphatidylinositol 3-kinase, MEK-1 and p38 mediate leptin/interferon-gamma synergistic NOS type II induction in chondrocytes. *Life Sci* 81: 1452-1460.
160. Yamagishi SI, Edelstein D, Du XL, Kaneda Y, Guzmán M, et al. (2001) Leptin induces mitochondrial superoxide production and monocyte chemoattractant protein-1 expression in aortic endothelial cells by increasing fatty acid oxidation via protein kinase A. *J Biol Chem* 276: 25096-25100.
161. Oda A, Taniguchi T, Yokoyama M (2001) Leptin stimulates rat aortic smooth muscle cell proliferation and migration. *Kobe J Med Sci* 47: 141-150.
162. Konstantinides S, Schäfer K, Koschnick S, Loskutoff DJ (2001) Leptin-dependent platelet aggregation and arterial thrombosis suggests a mechanism for atherothrombotic disease in obesity. *J Clin Invest* 108: 1533-1540.
163. Maya-Monteiro CM, Almeida PE, D'Avila H, Martins AS, Rezende AP, et al. (2008) Leptin induces macrophage lipid body formation by a phosphatidylinositol 3-kinase- and mammalian target of rapamycin-dependent mechanism. *J Biol Chem* 283: 2203-2210.
164. Adamczak M, Wiecek A, Funahashi T, Chudek J, Kokot F, et al. (2003) Decreased plasma adiponectin concentration in patients with essential hypertension. *Am J Hypertens* 16: 72-75.
165. Antoniadou C, Antonopoulos AS, Tousoulis D, Stefanadis C (2009) Adiponectin: from obesity to cardiovascular disease. *Obes Rev* 10: 269-279.
166. Jain SK, Croad JL, Velusamy T, Rains JL, Bull R (2010) Chromium dinicocysteinate supplementation can lower blood glucose, CRP, MCP-1, ICAM-1, creatinine, apparently mediated by elevated blood vitamin C and adiponectin and inhibition of NFkappaB, Akt, and Glut-2 in livers of Zucker diabetic fatty rats. *Mol Nutr Food Res* 54: 1371-1380.
167. Okamoto Y, Kihara S, Ouchi N, Nishida M, Arita Y, et al. (2002) Adiponectin reduces atherosclerosis in apolipoprotein E-deficient mice. *Circulation* 106: 2767-2770.
168. Ouchi N, Kihara S, Arita Y, Nishida M, Matsuyama A, et al. (2001) Adipocyte-derived plasma protein, adiponectin, suppresses lipid accumulation and class A scavenger receptor expression in human monocyte-derived macrophages. *Circulation* 103: 1057-1063.
169. Uji Y, Yamamoto H, Tsuchihashi H, Maeda K, Funahashi T, et al. (2009) Adiponectin deficiency is associated with severe polymicrobial sepsis, high inflammatory cytokine levels, and high mortality. *Surgery* 145: 550-557.
170. Han J, Luo T, Gu Y, Li G, Jia W, et al. (2009) Cathepsin K regulates adipocyte differentiation: possible involvement of type I collagen degradation. *Endocr J* 56: 55-63.
171. Taleb S, Canello R, Clément K, Lacasa D (2006) Cathepsin s promotes human preadipocyte differentiation: possible involvement of fibronectin degradation. *Endocrinology* 147: 4950-4959.
172. Taleb S, Lacasa D, Bastard JP, Poitou C, Canello R, et al. (2005) Cathepsin S, a novel biomarker of adiposity: relevance to atherogenesis. *FASEB J* 19: 1540-1542.
173. Yang M, Zhang Y, Pan J, Sun J, Liu J, et al. (2007) Cathepsin L activity controls adipogenesis and glucose tolerance. *Nat Cell Biol* 9: 970-977.
174. Xiao Y, Junfeng H, Tianhong L, Lu W, Shulin C, et al. (2006) Cathepsin K in adipocyte differentiation and its potential role in the pathogenesis of obesity. *J Clin Endocrinol Metab* 91: 4520-4527.
175. Yang L, Yuan J, Liu L, Shi C, Wang L, et al. (2013) \pm -linolenic acid inhibits human renal cell carcinoma cell proliferation through PPAR- β activation and COX-2 inhibition. *Oncol Lett* 6: 197-202.
176. Jiang WG, Redfern A, Bryce RP, Mansel RE (2000) Peroxisome proliferator activated receptor-gamma (PPAR-gamma) mediates the action of gamma linolenic acid in breast cancer cells. *Prostaglandins Leukot Essent Fatty Acids* 62: 119-127.
177. Li H, Ruan XZ, Powis SH, Fernando R, Mon WY, et al. (2005) EPA and DHA reduce LPS-induced inflammation responses in HK-2 cells: evidence for a PPAR-gamma-dependent mechanism. *Kidney Int* 67: 867-874.
178. Itoh T, Yamamoto K (2008) Peroxisome proliferator activated receptor gamma and oxidized docosahexaenoic acids as new class of ligand. *Naunyn Schmiedebergs Arch Pharmacol* 377: 541-547.
179. Schopfer FJ, Lin Y, Baker PR, Cui T, Garcia-Barrio M, et al. (2005) Nitrolinoleic acid: an endogenous peroxisome proliferator-activated receptor gamma ligand. *Proc Natl Acad Sci U S A* 102: 2340-2345.
180. Cimini A, Cristiano L, Colafarina S, Benedetti E, Di Loreto S, et al. (2005) PPARgamma-dependent effects of conjugated linoleic acid on the human glioblastoma cell line (ADF). *Int J Cancer* 117: 923-933.
181. Alexander RL, Wright MW, Gorczynski MJ, Smitherman PK, Akiyama TE, et al. (2009) Differential potencies of naturally occurring regioisomers of nitrolinoleic acid in PPARgamma activation. *Biochemistry* 48: 492-498.
182. Alaoui-El-Azher M, Wu Y, Havet N, Israël A, Lilenbaum A, et al. (2002) Arachidonic acid differentially affects basal and lipopolysaccharide-induced sPLA(2)-IIA expression in alveolar macrophages through NF-kappaB and PPAR-gamma-dependent pathways. *Mol Pharmacol* 61: 786-794.
183. Kurotani K, Sato M, Ejima Y, Nanri A, Yi S, et al. (2012) High levels of stearic acid, palmitoleic acid, and dihomo- β -linolenic acid and low levels of linoleic acid in serum cholesterol ester are associated with high insulin resistance. *Nutr Res* 32: 669-675.

184. Rabkin SW, Lodhia P (2009) Stearic acid-induced cardiac lipotoxicity is independent of cellular lipid and is mitigated by the fatty acids oleic and capric acid but not by the PPAR agonist troglitazone. *Exp Physiol* 94: 877-887.
185. Forman BM, Tontonoz P, Chen J, Brun RP, Spiegelman BM, et al. (1995) 15-Deoxy-delta 12, 14-prostaglandin J2 is a ligand for the adipocyte determination factor PPAR gamma. *Cell* 83: 803-812.
186. Davies SS, Pontsler AV, Marathe GK, Harrison KA, Murphy RC, et al. (2001) Oxidized alkyl phospholipids are specific, high affinity peroxisome proliferator-activated receptor gamma ligands and agonists. *J Biol Chem* 276: 16015-16023.
187. Dang ZC, Audinot V, Papapoulos SE, Boutin JA, Löwik CW (2003) Peroxisome proliferator-activated receptor gamma (PPARGamma) as a molecular target for the soy phytoestrogen genistein. *J Biol Chem* 278: 962-967.
188. Cho KW, Lee OH, Banz WJ, Moustaid-Moussa N, Shay NF, et al. (2010) Daidzein and the daidzein metabolite, equol, enhance adipocyte differentiation and PPARgamma transcriptional activity. *J Nutr Biochem* 21: 841-847.
189. Shen P, Liu MH, Ng TY, Chan YH, Yong EL (2006) Differential effects of isoflavones, from *Astragalus membranaceus* and *Pueraria thomsonii*, on the activation of PPARalpha, PPARgamma, and adipocyte differentiation in vitro. *J Nutr* 136: 899-905.
190. Salam NK, Huang TH, Kota BP, Kim MS, Li Y, et al. (2008) Novel PPAR-gamma agonists identified from a natural product library: a virtual screening, induced-fit docking and biological assay study. *Chem Biol Drug Des* 71: 57-70.
191. Jungbauer A, Medjakovic S (2012) Anti-inflammatory properties of culinary herbs and spices that ameliorate the effects of metabolic syndrome. *Maturitas* 71: 227-239.