

Prevalence and Associated Factors of Sexual Violence among High School Female Students in Dilla Town, Gedeo Zone SNNPR, Ethiopia

Desalegn Tarekegn¹, Balcha Berhanu² and Yigrem Ali^{3*}

¹Department of Midwifery, college of health sciences and Medicine, Dilla University, Dilla Ethiopia

²Department of Nursing and Midwifery, college of health sciences and Medicine, Addis Ababa University, Addis Ababa, Ethiopia

³Department of Psychiatry, college of health sciences and Medicine, Dilla University, Dilla Ethiopia

*Corresponding author: Yigrem Ali, Department of Psychiatry, College of Health Sciences and Medicine, Dilla University, Dilla Ethiopia, Tel: 0910461378; E-mail: alyigrem@gmail.com

Received date: May 26, 2017; Accepted date: July 12, 2017; Published date: July 24, 2017

Copyright: © 2017 Tarekegn D et al., This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Violence against girls takes many forms including rape, sexual harassment, intimidation, teasing and threats. It affects all girls, regardless of age, race, class, caste or location. Poverty, war and long journeys to school put girls at additional risk. The causes are rooted in male-dominated cultures which belittle or condone violence against girls and women. Therefore; this study aimed to assess the prevalence and associated factors of sexual violence against women among high school students in Dilla town.

Methodology: Institutional based Cross-sectional study was conducted high school female students at Dilla town, Gedeo zone. Also total sample size of this study was 280 participants. The data was collected by using self-administered questionnaires. Data was coded and entered in to EPI data software and was exported to SPSS version 20. Crude and adjusted OR were analyzed using logistic regression and the level of significance of association was determined at P- value<0.05.

Results: The Findings of this study indicated that the prevalence of sexual violence was 13.2%. The minimum age of the study subject was 14 years and the maximum age was 21 years. From the respondents who encountered sexual violence, majority of the respondents are victimized by unknown person and much older than their age. Using chat, alcohol drinking and using drugs or substances was the main associated factors of sexual violence.

Conclusion: The study indicates that female high school students in Dilla town are at high risk of sexual violence. Unwelcome touching of body part, facing sexual attempt, having boyfriend, and a habit of taking alcohol and chat are associated with life time experience of sexual violence.

Keywords: Prevalence; Sexual violence; Female students; Ethiopia

Introduction

Sexual violence is any sexual act, attempt to obtain a sexual act, unwanted sexual comments or advances, or acts to traffic, or otherwise directed, against a person's sexuality using coercion, by any person regardless of their relationship to the victim, in any setting, including but not limited to home and work [1].

The term violence against women includes forms of violence such as sexual partner violence and non-sexual partner violence [2]. It was found that 0.3% to 12% of women respondents reported being forced to have sex or to perform a sexual act that they did not want to by non-partners since the age of 15 years [3]. In rural Tanzania, 47% of ever-partnered women have ever experienced physical violence by an intimate partner, while 31% have ever experienced sexual violence. In rural Ethiopia, 49% of ever-partnered women have ever experienced physical violence by an intimate partner, rising to 59% ever experiencing sexual violence [4].

Violence against girls takes many forms including rape, sexual harassment, intimidation, teasing and threats [5,6]. Rape now carries

the additional danger of HIV infection. Girls are the first to be taken out of school as families are affected by HIV/AIDS [6].

Women are significantly more likely than men to be perpetrated of dating violence during young adulthood (38% vs. 19%). Based on the World Health Organization's report, the rate of violence against women is as following, Eastern Mediterranean countries (37%), and African countries (36.3%) American countries (29.8%), European countries (25.4%), Western Pacific (24.6%) and other countries with high salaries (23.2%) [2,7].

Level of education, economic freedom of women, men communication with their wives, the male dominated society, the physically stronger nature of men, presence of alcohol or drug use, unsatisfactory income level, and witnessing domestic violence to their mothers during childhood [8,9].

Sexual violence is an extreme restriction of the sexual and reproductive autonomy of women [10]. Although it is largely hidden by the victims, it is estimated that 12 million people around the world face sexual violence every year. Young people are the most frequent victims of sexual violence; it is generally thought that 12% to 25% of girls and 8% to 10% of boys under 18 years of age will suffer sexual violence [11]. Therefore, assessing and showing the significance of

violence against girls at school institutions might be important to enforce policy makers and different stakeholders to integrate this problem.

Methods

Study design, area and period

Institution based cross-sectional was conducted on high schools students in SNNP regional state, Dilla town from April to March 2015 G.C. Dilla town is the center of the Gedeo zone. The town is located 359 kilo meter distance far from Addis Ababa in the southern part of Ethiopia. Dilla has an estimated total population of 209,000, of whom 107,000 were males and 102,000 were females. There are three public high schools and three private high schools in the town in 2015. Trade is the main source of income of the residents.

Sampling size determination and procedure

It was determined by Level of significance (0.05), Power (0.50) with $z=95\%$ confidence interval and by considering assumptions of (p =proportion of prevalence) sexual violence among high school students in Ethiopia was 25% from the study done in Harare region [7], Then by adding 10% of non respondents and adjusting of the sample size since ($n < 10,000$), thus total sample size for this study is 280. A series of sampling methods was used to achieve actual study subjects. The stratified sampling proportional to size allocation method used based on their grade (9th-12th). Finally a simple random sampling technique was applied to select actual participants by using their alphabetical registration numbers/roster as sampling frame. Finally the study subject was selected using proportional allocation in each class.

Data collection and analyses procedures

Data collection instruments and collection technique: Pretested and structured self-administered questionnaires was prepared in English and then translated into Amharic and back to English to ensure its consistence of translation. The questionnaire was developed by adapting other questionnaires previous similar studies materials and review of relevant literature. With three days training, three diploma holder facilitated and guided the data collection process.

Data quality control issues: Data quality control issues was insured by conducting the pretest prior to the actual study period on 5% of the students from my total sample study subjects on the other high school students in Yirgachefe town. It was also insured by giving training for the facilitators as well as strict and regular supervision at the period of data collection.

Data processing and analysis procedures

First the data was checked for completeness and consistency. Then it was coded and entered in the computer using EPI data software. Then, data was analyzed by using SPSS version 21.0. Descriptive statistic was used to explain the study participants in relation to study variables. A Bivariate analysis was performed to determine the effect each of factors on the outcome variable. Only factors with p value < 0.2 on Bivariate analyses were kept for multivariate analyses and a p value of < 0.05 on multivariate analyses was considered as statistically significant.

Ethical consideration

Ethical clearance or approval was obtained from school of graduate studies of Addis Ababa University and then by Institutional reviews board of the University. Permission was obtained from schools directors. Informed consent would obtain from each participant. Privacy and confidentiality of the information was kept secretly by excluding names for identification in the questionnaire. The study variable would have the right to refuse and withdrawal from this study at any time. Individuals must understand the nature of the study and possible implications. The rights of individuals must be protected. Coercion and deception must be avoided at all.

Results

Socio-demographic characteristics

A total of 280 female students participated in the study with 97% response rate. Based on the sample size determined, among the total respondent students 104 (38.1%) from 9th grade, 72 (26.4%) from 10th grade, 55 (20.1%) from 11th grade and 42(15.4%) from 12th grade have participated in the study (Table 1).

The minimum age of the respondents was 14 years and the maximum was 21years and the mean age of the respondent's was 16 years. Among the total respondents, the predominant religion was protestant 116 (42.5%) and also the predominant ethnic group was gedeo 114 (41.8%) of the respondents. Majority 266 (97.4%) of the respondent were never married. 263(96.3%) of the respondent usual place of residence was in Dilla town. Regarding current living condition, 219 (80.2%) were reported to live together with both parents and 5 (1.8%) of the respondent were living alone (Table 1).

Variables	Frequency	Percent (%)
Age		
<16 years	91	33.3
17-19 years	149	54.6
>19 years	33	12.1
Religion		
Orthodox	106	38.8
Muslim	33	12.1
Catholic	11	4
Protestant	116	42.5
Other	7	2.6
Ethnicity		
Amhara	39	14.3
Oromo	38	13.9
Tigre	16	5.9
Gurage	36	13.2
Gedeo	114	41.8
other	30	11

Total	273	100
Living with		
With my parents	219	80.2
With my boy friend	13	4.8
With my husband	6	2.2
With relative	30	11
Alone	5	1.8
total	273	100
Marital status		
Yes	7	2.6
No	266	97.4
Education		
Grade 9 th	104	38.1
Grade 10 th	72	26.4
Grade 11 th	55	20.1
Grade 12 th	42	15.4
Residence		
Rural	10	3.7
Urban	263	96.3

Table 1: Frequency distribution of socio demographic distribution of female high school students under study in Dilla town Gedeo zone SNNP, Ethiopia 2015 G.C.

Family history of the respondents

From 273 of the respondents 115(42.1%) of their father educational level were certificate and above. But majorities of the respondents father educational level was only grade 12th completed, from this parents 33(12.1%) was illiterate. Majority 116(42.6%) of the respondents mother educational level was up to 8th grade completed, 64(23.4%) had certificate and above and 43(15.8%) of the respondents mother was they cannot read and write (Table 2).

From 273 of the respondents majority 201 (73.6%) of the students both parents had live together, 28 (10.3%) of the respondents parent were divorced, 34 (12.5%) of the respondents only mother alive 10 (3.7%) of the respondents were father only alive. More than half 147 (53.8%) of the respondents family income was between 1000 and 2500birr in month and 86 (31.5%) of the respondents family income were less than one thousands birr in month.

However from those study participants majority 186 (68.1%) of the respondents had get enough pocket money from their family, 87 (31.9%) of the respondents had not get enough pocket money from their family.

Regarding's to family relationship 41 (15%) of the respondents watch their mother beaten by her husband and majority 232 (85%) of the respondent's mother had not beaten by her husband (Table 2).

Variables	Frequency	Percent (%)
Father education level		
Illiterate	33	12.1
1-8 grade	70	29.4
9-12 grade	45	16.5
Diploma and above	115	42.1
Mother education level		
Illiterate	43	15.8
1-8 grade	116	42.6
9-12 grade	50	18.5
Diploma and above	64	23.4
Mother and father live together		
Yes	201	73.6
Divorced	28	10.3
Mother only alive	34	12.5
Father only alive	10	3.7
Family monthly income in birr		
<100 birr	86	31.5
1000-2500 birr	147	53.8
>2500 birr	40	14.7
Receiving enough money		
Yes	186	68.1
No	87	31.9
Mother being beaten by her husband?		
Yes	41	15
No	232	85

Table 2: frequency distribution of parent or family history of female high school students in dilla town Gedeo zone SNNP, Ethiopia 2015 G.C.

Factors associated with sexual violence against female students

In my study alcohol drinking, chat chewing and use illicit drug is the main factors that exposed students to sexual violence. From 273 study participants 198 (72.5%) of the respondents were not drink alcohol, 75 (27.5%) of the respondents drinking alcohol. From the respondents who had drinking alcohol majority 29 (38.7%) of them drink alcohol at least once or twice a week, 23 (30.7%) drink occasionally less than once in a month, 21 (28%) of the respondents drink alcohol two up to three times in a month and 2 (2.7%) of the respondents drink alcohol every day (Table 3).

Chat chewing was one of the factors associated with sexual violence in this study, from 273 study participants 34 (12.5) of the respondents

use chat and majority 239 (87.5%) of the respondents were not use chat. From the respondents who was using chat 2 (2.7%) of the respondents using chat daily, 17 (50%) of the respondents uses chat at least once or twice a week, 10 (29.4%) of the respondents uses chat one up to three times in a month and 5 (14.7%) of the respondents use chat occasionally less than once in a month (Table 3).

The other associated factor was drug or substance use, from 273 study participants 21 (7.7%) of the respondents had use drug or substance, majority 252 (92.3%) of the respondents had not uses drug or substances. From the respondents who had uses drug or substances, 1 (4.8%) of the respondents uses drug or substances daily, 2 (9.5%) of the respondents uses drug or substances at least once or twice a week, the majority 12 (57.1%) of the respondents uses drugs or substances one up to three times in a month and 6 (28.6%) of the respondents uses drug or substances occasionally less than once in a month (Table 3).

Variables	Frequency	Percent (%)
Drink alcohol		
Yes	75	27.5
no	198	72.5
Time drink alcohol		
Every day	2	2.7
Once or twice a week	29	38.7
1-3 times a month	21	28
Occasionally , less than once a month	23	30.7
Chewing chat		
Yes	34	12.5
No	239	87.5
Time of chat chewing		
Every day	2	5.9
Once or twice a week	17	50
1-3 times a month	10	29.4
Occasionally , less than once a month	5	14.7
Drugs or substances use		
Yes	21	7.7
No	252	92.3
Time of drug/substance use		
Every day	1	4.8
Once or twice a week	2	9.5
1-3 times a month	12	57.1
Occasionally , less than once a month	6	28.6

Table 3: frequency distribution offactors associated with sexual violence among female high school students in Dilla town Gedeo zone SNNP, Ethiopia, 2015 G.C.

Sexual history of the respondent

From the 273 total study participant majority 219 (80.2%) of the respondents had no regular boyfriend currently, 54 (19.8%) of the respondents had regular boyfriend currently. In addition from 273 study participants 64 (23.4%) of the respondents had sexual relation, and the rest 209 (76.6%) of the respondents had no sexual relation (Table 4). From those who had sexual relationship the highest number 21 (33.8%) of the respondents condition of start sexual intercourse was by peer pressure, the second highest number 18 (27.7%) of the respondents condition of starting sexual relation was by personal interest, 11 (16.9%), 5 (7.7%), 5 (7.7%) and 4 (6.2%) by marriage, for passing examination/for mark, by forcing against me, or financial purpose was the reason or condition of starting sexual relation respectively. In addition from those respondents who had sexual relation most 44 (67.7%) of them had sexual relation before the age of 17, 20 (30.8%) of the respondents had sexual relation between the age of 17 and 19, only 1 (1.5%) had sexual relation above the age of 19. there for almost all of the respondents who had sexual relation they start sexual relation before the 19 years (Table 4).

Concerning's to sexual harassment from the total study participants 61 (22.3%) of the respondent had faced unwanted sexual act like talking unnecessary words and the majority 212 (77.7%) of the respondents had not faced unwanted sexual act. Moreover from total study participants 65 (23.8%) of the respondents had facing unwanted kiss, 208 (76.2%) of the respondents had not facing unwanted kiss. On the other hand from 273 study participants 50 (18.3%) of the respondents had facing unwelcome touching of body parts likes breast, hair and majority 223 (81.7%) of the respondents had not facing unwelcome touching of body parts (Table 4).

Regarding's to forced sexual attempt from 273 study participants 46 (16.8%) of the respondents had encountered forced sexual attempt but they escaped from that sexual violence by different means; 12 (36.4%) of the respondents escaped by shouting resulting to arriving somebody, 8 (24.1%) of the respondents were escaped from that sexual attempt by hiding them self from that person/attackers, 6 (18.2%) of the respondents were escaping from that sexual attempt by taking a defensive measure and fighting against the attacker, 7 (21.2%) of the respondents was escaped by giving false promise word from sexual attempt and the majority 227 (83.2%) of the respondents were had not facing forced sexual attempt (Table 4).

Surprisingly from 273 study participants 36 (13.2%) of the respondents had encountered sexual violence, most of the victim students 16 (50%) of the respondents was victimized in hotel, 6 (18.8%) of the respondents was victimized in their home, 6 (18.8%) of the respondents was victimized in school and the rest 4 (12.5%) of the respondents was victimized in perpetrators home and also 237 (86.8%) of the respondents were had not encountered forced sexual intercourse, in addition to that from the respondents who was encountered sexual violence majority 31 (86%) of the respondents violence is made by frightened by using different sharp material (Table 4).

Concerning to the time of sexual violence from those who encountered sexual violence more than half 19 (59.4%) of the respondents were victimized in day time and 13 (40.6%) of the respondents were victimized in night time. In addition from the respondents who had sexual violence majority 17 (50%) of the respondents had faced two times in their life time.

Moreover most 9 (28.1%) of the students encountered sexual violence by unknown person and majority 22 (68.8%) of the students sexual violence was done by older men compared the female students age (Table 4).

Variables	Frequency	Percent (%)
Have regular boyfriend currently		
Yes	54	19.8
No	219	80.2
Had a sexual intercourse		
Yes	64	23.4
No	209	76.6
Conditions of start sexual intercourse		
In a marriage	11	16.9
Personal desire	18	27.7
Peer pressure	21	33.8
For financial purpose	4	6.2
For passing examination	5	7.7
By forcing against you	5	7.7
total	64	100
Age at first sexual intercourse		
<17years	44	67.7
17-19 years	20	30.8
>19 years	1	1.5
Unwanted sexual act		
Yes	61	22.3
No	212	77.7
Faced unwanted kiss		
Yes	65	23.8
no	208	76.2
Unwelcome touching of body part		
Yes	50	18.3
no	223	81.7
Forced sexual attempt		
Yes	46	16.8
no	227	83.2
Forced sex		
Yes	36	13.2
no	237	86.8
Forced sex that made by frightened and degrading		

Yes	31	11.4
no	241	88.6
Place of forced sex encountered		
In my home	6	18.8
In his home	4	12.5
In school	6	18.8
In hotel	16	50
Time of forced sex encountered		
Day	19	59.4
Night	13	40.6
Age of Perpetuator		
Same age	4	12.5
Older than me	22	68.8
Much older than me	6	18.8
Numbers of forced sex encountered in life time		
One times	14	41.2
Two times	17	50
Three times	1	2.9
More than four times	2	5.9
Numbers of boyfriend experienced		
One times	13	39.4
Two times	15	45.5
Three times	4	12.5
More than four times	1	3

Table 4: Frequency distribution of sexual history of female high school students among high school students in Dilla town Gedeo zone SNNP, Ethiopia 2015 G.C.

Factors associated with sexual violence

The likelihood of experiencing sexual violence in their life time was higher among those students who uses drugs or substances than students who didn't have using drugs or substances, (OR=5.103, 95% CI, 1.174, 22.188), (Table 5).

In this study using drugs or substances had significant association with sexual violence (p=0.030) (Table 5).

Among the female students, the odds of life time experiencing sexual violence was higher among those students who had using chats than those who had not using chat (OR=15.099, 95% CI, 3.735, 61.038), (Table 5).

Using chat was one of the major significant variables that related with life time sexual violence. Therefore using chat was the main associated factors of sexual violence (p-0.000) (Table 5).

The likelihood of experiencing sexual violence in their life time was higher among those students who had a habit of taking alcohol than those who didn't have a habit of alcohol intake, (OR=1.579, 95% CI; 0.450, 5.541) (Table 5) and the chance of experiencing life time rape was higher among those who were having a low income family than those who had medium and higher income of family (AOR=0.912, 95% CI=0.227, 3.667) (Table 5), From the respondents who had not getting enough amount of pocket money was more exposed than those who had getting enough amount of money (AOR= 1.303, 95% CI, 0.430, 3.953) (Table 5).

Variables	Sexual violence		AOR	95% CI	p-value
	Yes	No			
Family monthly income in birr					
<1000	19	67	0.912	(0.227,3.667)	0.896
1000-2500	13	134			
>2500	4	36			
Receiving enough money					
Yes	30	156	1.303	(0.430, 3.953)	0.64
No	6	81			
Alcohol drinking					
Yes	26	49	1.579	(0.450, 5.541)	0.476
No	10	188			
Using chat					
Yes	24	10	15.099	(0.735, 61.038)	0
No	12	227			
Using drugs or substances					
Yes	16	5	5.103	(1.174, 22.188)	0.03
No	20	232			

Table 5: Multivariate analysis of sexual violence and factors associated with sexual violence of female high school students in dilla town.

Sexual factors associated with sexual violence

According to the result of this study from the respondents who had regular boyfriend currently (AOR=1.009, 95% CI, 0.212, 4.806) was highly exposed to sexual violence than those who had no regular boyfriend currently.

In addition in this study from the respondents who had sexual intercourse was significant association with sexual violence (p=0.010) and the odds of the respondents (AOR=5.760, 95% CI -1.517, 21.871) this value indicates that from the students who had sexual intercourse has around six times risk of exposure to sexual violence than those who had not sexual intercourse.

As indicated in (Table 6) the odds of female students who had encountered unwanted sexual act was (AOR 1.860, 95% CI, 0.319, 10.856) this result indicates that the students who had face unwanted sexual act was two time exposure risk than those who had not face unwanted sexual act, From the respondents who had encountered


unwelcome touching of their body part had sixteen times risk to sexual violence than those who had not facing unwelcome touching of body part and in addition to that unwelcome touching of body part was significant association with sexual violence(p=0.004).

Moreover in this study facing sexual attempt was significant association with sexual violence with a p-value of (p=0.009) and the odds of the respondents who had facing forced sexual attempt was (AOR=9.426, 95% CI(1.757, 50.568) this indicates that nine times risk than those who had not facing forced sexual attempt.

Variables	Sexual violence		AOR	95% CI	p-value
	Yes	No			
Have regular boyfriend currently					
Yes	6	48	1.009	(0.212, 4.806)	0.991
No	30	189			
Have sexual intercourse					
Yes	26	38	5.76	(1.517, 21.871)	0.01
No	10	199			
Face Unwanted sexual act					
Yes	31	30	1.86	(0.319, 10.856)	0.491
No	5	207			
Encountered Unwanted kiss					
Yes	32	33	1.115	(0.134, 9.255)	0.92
No	4	204			
Unwelcome touching of body part					
Yes	33	17	16.512	(2.411, 113.103)	0.004
No	3	220			
Faced Forced sexual attempt					
Yes	32	14	9.426	(1.757, 50.568)	0.009
No	4	223			
Her mother beaten by her father					
Yes	4	37	1.443	(0.309, 6.745)	0.641
No	32	200			

Table 6: Multivariate analysis of sexual violence and independent variable of female high school students in dilla town.

Given below in the (Figure 1) shows that the mechanisms of female students how was escaped from sexual violence attempt, majority of 36.36% of the respondents escaping by shouting, 24.24%, 18.18%, and 21.21% of the respondents was escaped by hiding, by fighting and by giving a false hope respectively.



Discussion

In my study the prevalence of sexual violence its associated factors had been assessed and the life time prevalence rate of sexual violence among female high school students was 13.2% when compared to study's conducted in Harare shows that Sexual violence among in school adolescents were 25% (12, 7) and a study conducted in Bahir Dar town among private college female students revealed that the prevalence of sexual violence among female college students was 37.3% [12,13]. And also in this study had a lower prevalence rate than the study of Harare and Bahir Dar town. In addition to that a cross sectional study conducted in Iran indicates that the prevalence of sexual violence against women was found to be around 63.8%.

Regarding's to factors associated with sexual violence in this study using chat is one of the important contributing factors for sexual violence of school students in dilla town ($p=0.000$) (Table 5) and in the other hand using drugs or substances like shisha had significant association with sexual violence with ($p=0.030$). When I compared with other past study on sexual violence, a study conducted in Harare revealed that students used addictive drugs like alcohol (18%), chat and shish (7.2%) as very important contributing factors for sexual violence's [12,7]. Also this study finding supports my study findings.

Furthermore a study done in eastern Ethiopia with a multivariate analyses revealed that high-rejection sensitivity, having multiple sexual partners, the frequent watching of pornography, and use of alcohol or other soft drugs (chat or shisha) are factors associated with higher levels of sexual violence victimization. On the other hand a study conducted in china indicates that, students who had sexual intercourse, drug use (odds ratios [OR, 2.44], attempted suicide (OR, 2.30), physical abuse (OR, 1.74), binge drinking (OR, 1.62), verbal abuse (OR, 1.29), experience of being drunk (OR, 0.68), and smoking of cigarettes (OR, 0.52) were related to a history of sexual coercion [19].

In addition a study in Brazil indicates that the prevalence of violence against women was 27.5%. The associated factors for sexual violence were low level of schooling (OR=2.34), a history of domestic violence

(OR=2.21) the woman being mentally disturbed (OR=2.35), and the partner's consumption of alcohol (OR=1.77) [16]. This finding is also having some similarity with my study explored.

Findings of this study revealed that family income was one of the associated factors of sexual violence among female high school students (AOR=0.912, 95% CI, 0.227, 3.667) those students who had low amount of income of family was had more experienced sexual violence than those who had medium and higher income family. A study conducted in Nigeria indicates that most of violence was associated with economical reason; the prevalence of violence against women (VAW) six months preceding the survey was 52.5%. Sexual violence was the commonest type (41.9%) of violence experienced, followed by economic (37.7%), physical violence (35.7%) and psychological (31.9%) [19].

According to this findings from the respondents who had sexual intercourse was significant association with students exposure to sexual violence (AOR=5.670, 95% CI 1.517, 21.871, $p=0.010$) five times more likely exposure to sexual violence than not had sexual intercourse. The same study conducted in Nigeria indicates that sexual violence was significantly more experienced (AOR 2.23; 95% CI, 1.15-4.36) by older female sex workers (FSWs) than their younger counterparts, by permanent brothel residents (AOR 2.08; 95% CI 1.22-3.55) and among those who had been in the sex industry for more than five years [17-19]. Also this findings helps to conclude that females who had on sexual relation more likely vulnerable to sexual violence than those who had not sexual relation.

The result of these findings indicates that from the respondents who had unwelcome touching of body part was more likely vulnerable to sexual violence than those who had not unwelcome touching of body part. Other study conducted in Harare indicates that females dressing style, their act, peer pressure, revenge and males emotionality identified as contributing factors for sexual violence [12,7].

According to this study from the respondents who had boyfriend/partner was more likely vulnerable to sexual violence than those who had not have boyfriends or partners (AOR=1.009, 95% CI 0.212, 4.806, $p=0.991$) may the violence made by their partner. Also this findings had supported by a study conducted by WHO multi countries study against women the prevalence of partner violence nearly one half (49%) of ever-partnered women experienced physical violence by a partner at some point in their lives, and 29% during the past 12 months. 59% of ever-partnered women experienced sexual violence at some point, and 44% during the past 12 months. Combining the data for physical and sexual violence, 71% of ever-partnered women experienced one or the other form of violence, or both, over their lifetime [12].

Conclusion

Prevalence of sexual violence still is high among female high school students in Dilla town. Life time experience of sexual violence was significantly associated with the income of their family, using of drug or substances and habit of taking alcohol and chat.

Competing Interest

All authors declare that they have no conflict of interest associated with the publication of this manuscript.

Author's Contribution

Desalegn Tarekegn conceived and designed the study and collected data in the field, performed analysis, interpretation of data, and draft the manuscript. Mr. Balcha Berhanu and Mr. Yigrem Ali involved in the design, analysis, and interpretation of data and the critical review of the manuscript. All authors approved and read the final manuscript.

Acknowledgement

This study was funded by Addis Ababa University and it covered all the financial and material support for the research.

We would like to thanks Dilla Educational Office and teachers for their unlimited cooperation during data collection'

We also want to acknowledge participants (female students) for providing us important materials and for giving us relevant information.

References

1. Hamburg Medico-Legal Competence Centre (2008) Sexual and gender based violence in Africa. Population Council Inst.
2. Nikbakht NA, Hossein Abbasi N, Mehrdad N (2015) The Prevalence of Violence against Iranian Women and Its Related Factors. *Glob J Health Sci* 7: 37-45.
3. Shimekaw B, Megabiaw B, Alamrew Z (2013) Prevalence and associated factors of sexual violence among private college female students in Bahir Dar city, North Western Ethiopia. *Biomed Life Sci* 5: 1069-1075.
4. Africa population council (2008) Sexual and gender based violence in Africa.
5. World Health Organization (2012) Understanding and addressing violence against women.
6. Action Aid International (2004) Violence against girls and the right to education, stop violence against girls in school.
7. Cafo MJ, Demisie AS, Abera BB (2014) Assessment of sexual violence and associated factors among high school students in Harari Regional State, Harar Town, Eastern Ethiopia. *Sci Res* 2: 91-97
8. World Health Organization (2013) Intimate partner violence and alcohol fact sheet.
9. Efe SY, Ayaz S (2010) Domestic violence against women and women's opinions related to domestic violence. *Anatolian J Psychiat* 11: 23-29.
10. Lamichhane P, Puri M, Tamang J, Dulal B (2011) Women's status and violence against young married women in rural Nepal. *BMC Womens Health* 11: 19.
11. Sapp MV, Vandeven AM (2005) Update on childhood sexual abuse. *Curr Opin Pediatr* 17: 258-264.
12. Bekele AB, Van Aken MA, Dubas JS (2011) Sexual Violence Victimization Among Female Secondary School Students in Eastern Ethiopia. *Violence Vict* 26: 608-630.
13. Silva MA, Falbo Neto GH, Figueiroa JN, Cabral Filho JE (2010) Violence against women: prevalence and associated factors in patients attending a public healthcare service in the Northeast of Brazil. *Cad Saúde Pública* 26: 264-272.
14. Fawole OI, Dagunduro AT (2014) Prevalence and correlates of violence against female sex workers in Abuja, Nigeria. *Afr Health Sci* 14: 299-313.
15. Swain SN, Saggurti N, Battala M, Verma RK, Jain AK (2011) Experience of violence and adverse reproductive health outcomes, HIV risks among mobile female sex workers in India. *BMC Pub Health* 11: 357.
16. WHO (2005) Multi-country Study on Women's Health and Domestic Violence against Women.
17. Gokler ME, Arslantas D, Unsal A (2014) Prevalence of domestic violence and associated factors among married women in a semi-rural area of western Turkey. *Pak J Med Sci* 30: 1088-1093.
18. Song Y, Ji CY, Agardh A (2014) Sexual coercion and health-risk behaviors among urban Chinese high school students, *Glob Health Action* 7: 24418.
19. Bekel T, Deressa W (2012) Prevalence and experience of sexual coercion among female students of Ambo University in Ethiopia. MPH thesis.