

Prevalence of Anemia among Pregnant Women Attending Antenatal Care at Tikur Anbessa Specialized Hospital, Addis Ababa Ethiopia

Alemayehu Hailu Jufar^{1*} and Tewabech Zewde²

¹Department of Physiology, College of Health Sciences, Mekelle University, P.O.BOX 1871, Mekelle, Ethiopia

²Department of Physiology, School of Medicine, College of Health Sciences, Addis Ababa University, P.O.BOX 16744, Addis Ababa, Ethiopia

Abstract

Background: Anemia impairs cognitive development, reduces physical work capacity and in severe cases increases risk of mortality particularly during prenatal period. There are no previous studies done on anemia in pregnancy in the study area.

Objective: To assess the prevalence and determinant factors of anemia among pregnant women attending antenatal care follow up at Tikur Anbessa specialized hospital.

Methods: Health institutional based cross-sectional study design was conducted from November 23, 2013 - March 8, 2013 among 395 pregnant mothers who had been attending antenatal care. Blood sample was taken to determine the hemoglobin concentration and haematocrit value. Interviewer administered structured questionnaire was used to assess different socio-demographic characteristics, obstetrics and dietary factors related to anaemia. Simple random sampling technique was used to select study subjects. Data were entered with EPI info version 3.5.1. Cleaning and analysis was done using SPSS version 17 and statistical tests were performed at the level of significance of 5%.

Result: The overall prevalence of anemia in this study using a cut off level of hemoglobin <11 g/dl (<33% haematocrit) was 21.3% (84/395). Out of 84 anemic pregnant mothers, 80.95% (68/84) were mildly anemic, 17.86% (15/84) were moderately anemic and 1.19% (1/84) were severely anemic. The following variables were significantly associated with anemia: Age (39-45 yrs (AOR=1.21)), education status (illiterate (AOR=2.12)), family size (greater than four (AOR=2.04), gestational age (third trimester (AOR=2.04)), birth intervals (less than two years (AOR=1.21)), history of blood loss (AOR=1.67)), antenatal care (no ANC (AOR=1.11)), contraception (no contraception (AOR=2.31)), gravidity (multigravid (AOR=1.22)), parity (multiparous (AOR=2.19)).

Conclusion: The present study showed high prevalence of anemia and the majority of them were of the mild type (hemoglobin: 10-10.9 g/dl).

Keywords: Anemia; Pregnancy; Prevalence; Hemoglobin

Introduction

Anemia refers to a condition in which the hemoglobin content of the blood is lower than normal for a person's age, gender and environment, resulting in the oxygen carrying capacity of the blood being reduced [1,2]. The mean minimum acceptable hemoglobin level during pregnancy by WHO criteria is taken to be 11 g/dL [2]. WHO further divides anemia in pregnancy into: mild anemia (hemoglobin 10-10.9 g/dL), moderate anemia (hemoglobin 7.0-9.9 g/dL) and severe anemia (hemoglobin <7 g/dL [2]. It is a global health problem affecting both developing and developed countries with major adverse effects on human health as well as social and economic development [1-5]. In 1993, WHO ranked anemia as the 8th leading cause of disease in girls and women in the developing world [2]. Estimates from the World Health Organization report that 35% to 75% of pregnant women in developing countries, and 18% in developed countries are anemic [2]. The greatest burden of anemia is born by Asia and Africa where it is estimated that 60% and 52% of pregnant women, respectively, are anemic and between 1% and 5% are severely anemic (hemoglobin <7 g/dl) and is associated with women of age less than 20 years, third trimester of pregnancy, rural residents, and multiparous women [6,7]. United Nations expert panel considered severe anemia an associated cause in up to half of the maternal deaths worldwide [8]. Estimates of maternal mortality resulting from anemia range from 34/100,000 live births in Nigeria to as high as 194/100,000 in Pakistan [6,7]. In sub-Saharan Africa it is estimated that 20% of maternal deaths are associated with anemia [2].

In developing countries, the cause of anemia during pregnancy is multi-factorial and includes nutritional deficiencies of iron, folate,

and vitamin B12 and also parasitic diseases, such as malaria and hookworm. The relative contribution of each of these factors to anemia during pregnancy varies greatly by geographical location, season, and dietary practice. In sub-Saharan Africa, iron and folate deficiencies are the most common causes of anemia in pregnant women [1,9]. Vitamin B12 deficiency may be an unrecognized contributor to anemia in this region of the world due to reliance of the population on grains as dietary staples and low consumption of foods of animal origin which are the primary source of dietary vitamin B12 [9].

Although Ethiopia has different climatic conditions and grows a variety of cereals, root crops and vegetables, some of these are not fully utilized [10,11]. There appears to be dependency on a single food crop by region although the specific crop varies in the different regions [10]. The staple crops consumed in the North and Central part of Ethiopia are teff and cereals; in the South and Southwest staple crops are enset, maize, cereals and root crops; and in the East staple crops are sorghum and maize [10]. The lack of dietary diversity results in a shortage of

***Corresponding author:** Alemayehu Hailu Jufar, Department of Physiology, College of Health Sciences, Mekelle University, P.O.BOX 1871, Mekelle, Ethiopia, Tel: +251911172095; E-mail: alemayehuhailu62@yahoo.com

Received December 04, 2013; **Accepted** December 30, 2013; **Published** January 01, 2014

Citation: Jufar AH, Zewde T (2014) Prevalence of Anemia among Pregnant Women Attending Antenatal Care at Tikur Anbessa Specialized Hospital, Addis Ababa Ethiopia. J Hematol Thromb Dis 2: 125 doi: [10.4172/2329-8790.1000125](http://dx.doi.org/10.4172/2329-8790.1000125)

Copyright: © 2014 Jufar AH, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

minerals and vitamins which suggests that the bio-availability of much of the iron in the average Ethiopian diet is restricted and this restriction may have affected the iron status [10]. The present study was designed to see the hemoglobin levels in pregnant women at Tikur Anbessa specialize hospital. The results of this study will help to motivate ante natal care providers towards early detection and management of anemia in pregnancy

Materials and Methods

A cross-sectional study design was conducted at Tikur Anbessa Specialized Hospital from November 23, 2012- March8, 2013. Source population were all pregnant women who live in Addis Ababa and study population were pregnant women attending antenatal care at Tikur Anbessa specialized hospital, Addis Ababa Ethiopia and study subjects were pregnant women attending antenatal care during the study period and who met the inclusion criteria. A total of 395 pregnant mothers were enrolled in this study.

Sampling

The required sample size for this study was calculated using formula for single population proportion based on the prevalence rate of 62.7% reported from the previous study and using the 95% confidence interval and 5% marginal error. By adding 10% for none response the final sample size will be 395.

The study participants were selected based on the inclusion and exclusion criteria by using simple random sampling method. Exclusion criteria were pregnant women who were severely sick due to other medical conditions and those who were on treatment for anemia.

Inclusion criteria

In this study, pregnant mothers who were residents of Addis Ababa, not recently transfused, who had no chronic medical diseases, no diagnosed haemoglobinopathies, and who had no early bleeding or antepartum hemorrhage were included.

Data collection

A structured and interviewer administered questionnaire was used to collect data on the socio-demographic variables, dietary habit, and obstetric factors. The questionnaire of this study was developed from the Ethiopian Demographic and Health Survey (EDHS) guide lines which essentially contains the same information of the WHO guide lines. Data was collected by midwives of the hospital. Venous blood was collected to determine haemoglobin concentration after an informed consent is obtained from the study subjects.

Data analysis

Data were entered with EPI info version 3.5.1. Analysis was done by using SPSS version 17. Mean frequency and p-value has been calculated. After multivariate logistic regression analysis was used variables significant at the p value<0.05 are kept in the final model

Ethical considerations

Ethical approval and clearance was taken from institutional review board of College of Health Sciences, Addis Ababa University and permission letter for data and sample collection has also been received from the Department of Obstetrics and gynecology. Data and blood sample collection was started after written informed consent was obtained and after the objectives, benefits and risks of the study were explained to the study participants. Participation in the study was voluntary, and refusal to participate involved no loss of benefits to

which the mothers were otherwise entitled. Study participants had the right to accept or refuse participation in the study at any time.

Results

Socio-demographic characteristics of study subjects

Total of 395 informed and consented pregnant women who came for their antenatal follow up were enrolled in this study and all were urban dwellers. The mean age of the attendants was 28.8 years old (range from 18-46). Majority of the study groups were in the age range of 25-31 years. Many of the respondents were married 380 (96.6%). One hundred ninety two (48.6%) of the respondents were housewife and 27 (6.8%) were merchants. Forty percent of the respondents had educational status of 9-12 grade and 122 (30.9%) had diploma/degree. One hundred twenty seven of the respondents did not know their income and 125(31.6%) had an average income. Two hundred fifteen (54.4%) respondents had family size of greater than four (Table 1).

Prevalence and severity of anemia

The overall prevalence of anemia using a cut off level of hemoglobin <11 g/dl (<33% haematocrit) was 21.3% (84/395). The mean haematocrit

Variables		Frequency(n)	Percentage (%)
Age	18-24	71	17.97
	25-31	182	46.1
	32-38	108	27.3
	39-45	31	7.8
	>45	3	0.8
Ethnicity	Amhara	179	45.3
	Oromo	96	24.3
	Tigre	40	10.1
	Gurage	52	13.2
	Others	28	7.1
Religion	Orthodox	285	72.2
	Muslim	65	16.5
	Protestant	39	9.9
	others	6	1.5
Marital status	Married	380	96.2
	Single	8	2
	Divorced	5	1.3
	Widowed	2	0.5
Occupation	House wife	192	48.6
	Civil servant	89	22.5
	Merchant	27	6.8
	others	87	22
Education status	Illiterate	20	5.1
	Read and write	14	3.5
	1-8 grade	81	20.5
	9-12 grade	158	40
	Diploma/degree	122	30.9
Income	< 230 ETB	1	0.3
	231-850ETB	55	13.9
	851-2350ETB	125	31.6
	>2350ETB	87	22
	I don't know	127	32.2
Family size	<=4	108	45.6
	>4	215	54.4

ETB: Ethiopian birr

Table 1: Socio demographic characteristics of pregnant women attending antenatal care at Tikur Anbessa Specialized Hospital from November 23, 2012 - March 8, 2013.

Characteristics		Anemic (N=84)	Non anemic (N=311)	Total	AOR(95%CI)	P-value
age	18-24	4 (4.8)	67 (21.5)	71	1	
	25-31	31(36.9)	151(48.6)	182	0.75(0.84,9.12)	0.81
	32-38	35 (41.7)	73 (23.5)	108	0.23(0.43,2.14)	0.51
	39-45	13 (15.5)	18 (5.8)	31	1.21(1.93,11.1)	0.033
	>45	1 (11.9)	2 (0.6)	3	2.12(0.65,1.34)	0.81
Ethnicity	Amhara	41(48.8)	138 (44.4)	179	1	
	Oromo	20 (23.8)	76 (24.4)	96	1.3(0.56,2.76)	0.588
	Tigre	11(13.1)	29 (9.3)	40	0.97(0.35,2.67)	0.946
	Gurage	7 (83.3)	45 (14.5)	52	0.57(0.198,1.6)	0.286
	others	5 (59.5)	23 (7.4)	28	0.54(0.15,1.97)	0.352
Religion	Orthodox	64 (76.2)	221 (71.1)	285	1	
	Muslim	15 (17.9)	50 (16.1)	65	0.76(0.33,1.8)	0.521
	Protestant	4 (4.8)	35 (11.3)	39	0.74(0.21,2.6)	0.641
	Others	1(1.2)	5 (1.6)	6	1.78(0.16,2.2)	0.64
Marital status	Married	81(96.4)	299 (96.1)	380	1	
	Single	1(1.2)	7 (2.3)	8	1.51(0.14,1.7)	0.74
	Divorced	2 (2.4)	3 (0.96)	5	1.2 (0.3 ,7.7)	0.07
	Widowed	0 (0)	2 (0.6)	2	1	
occupation	House wife	43 (51.1)	149 (47.9)	192	0.59(0.47,4.2)	0.584
	Civil servant	15 (17.9)	74 (23.8)	89	0.61(0.23,1.4)	0.234
	Merchant	5 (59.5)	24 (7.7)	27	0.75(0.22,2.6)	0.652
	others	21(25)	66 (31.2)	87	1.13(0.51,2.5)	0.768
Education status	Illiterate	9 (10.7)	11(3.5)	20	2.12(2.47,6.8)	0.037
	Read and write	4 (47.6)	10 (3.2)	14	1	
	1-8 grade	20 (23.8)	61(19.6)	81	1	
	9-12 grade	33 (39.3)	125 (40.2)	158	0.78(0.21,3.45)	0.56
	Diploma/degree	18 (21.4)	104 (33.4)	122	0.89(0.91,1.9)	0.66
Income	< 230 ETB	0 (0)	1(0.32)	1		
	231-850ETB	7(83.3)	45 (14.5)	55	0.66(0.27,1.6)	0.361
	851-2350ETB	28 (33.3)	97 (31.2)	125	0.434(0.15,1.3)	0.134
	>2350ETB	17 (20.2)	70 (22.5)	87	0.63(0.17,2.41)	0.613
	I don't know	31(36.9)	96 (30.9)	127	1.547(0.67,3.5)	0.290
Family size	<=4	16 (19)	164 (52.7)	180	1	
	> 4	68 (81)	147(47.3)	215	1.73(1.13,4.2)	0.043

ETB: Ethiopian birr

Table 2: Anemia and socio-demographic characteristics among pregnant women attending antenatal care follow up at Tikur Anbessa Hospital, from November 23, 2012 – March 8, 2013.

value was 0.36 (36%), ranging from 19-45%. Out of all anemic pregnant women about 80.95% (68/84) were mildly anemic, 17.86% (15/84) were moderately anemic and 1.19% (1/84) were severely anemic (Figure 1).

Anemia and socio-demographic characteristics

About 41.9%, 33.3%, 32.4% of the pregnant women who were in the age range of 39-45, >45 and 32-38 years were anemic respectively. Forty percent of anemic pregnant women were divorced and 21.3% were married. Forty five percent (45%) and 31.6% of anemic pregnant women were illiterate and had family size of greater than four respectively (Table 2).

Obstetrics and other characteristics and anemia

Two hundred sixty two (66.3%) of the respondents were multigravidae and one hundred thirty five (34.2%) were multiparous. About 40.3% of the respondents had birth interval of greater than two years and 42.5% of the multiparous pregnant women delivered their children at health institutions. Two hundred forty one (61%) of them had no history of excess blood loss/abortion in their previous pregnancy and 49.1% of the responded that they use contraceptives.


10.4% of the multigravidae did not follow ANC in their previous

pregnancy. Small number of the respondents, 2.5% and 1.5% had history of blood transfusion and malarial infection in the last one year respectively

The prevalence of anemia in this study was 17.29% and 23.37% for primigravida and multigravida respectively. The prevalence of anemia in multipara was more 24.44% (33/135) than primipara 22.22% (26/117) and in those who were nulliparous 25/143 (17.48%). Anemia was also found to increase as the gestational age increases, showing the highest prevalence in the third trimester 35.48% (33/93) than second 21.64% (29/134) and first trimester 13.09% (22/168). Women with birth interval of less than two years had shown more prevalence of anemia (32/93(34.4%)) than those with an interval of greater than or equal to two years 29/159 (18.23%) and women with no history of delivery 23/143(16.08%). Place of delivery also showed difference in prevalence of anemia (28.9% in the pregnant women who delivered at home and 22.02% in women who delivered at health institution).Contraceptive users showed lower prevalence of anemia (18.04%) compared to none users (24.37%) (Table 3).

Dietary habits and anemia

Out of 395 respondents 146 (36.96%) had the habit of eating meat and animal products once per week, 102 (25.82%) once in a month,



Discussion

The prevalence of anemia in this study population using a cut off level of Hb <11 g/dl (<33% haematocrit) was 21.3% (84/395). This result is almost consistent with across sectional study carried out in Gonder (23%) and in the University of Port Harcourt Teaching Hospital, Port Harcourt in Nigeria which had shown a prevalence rate of 23.2% [12,13]. In contrast to this study, a study conducted at Health Promotion Hospital Nakhonsawan, in Thailand showed a prevalence rate of 14.1% [14]. This may be due to a difference in socio-economic and educational status between the study populations in the two study areas. The result of the present study is much lower than that of Jima (57%), Assendabo (62.7%), Peru (50%), Western Maharashtra, India (92.38%), highlands of Tibet (China) (70%) and most of the pregnant women of developing countries (35-75%) [2,5,15-17]. Other similar studies conducted in east Anatonian province; Turkey, the highlands of Tanzania and in rural areas of Kenya had shown prevalence of 27.1%, 28% and 33% respectively [11,18,19]. This discrepancy might be because of the strengthened health educations given at health institutions during ANC follow up and that there was time difference and this study is done only in the urban women. Additionally, it could be due to the variation of the method where sahli's method was used in previous studies conducted in Jimma and Assendabo. In comparison to the standard method (Fluid based system hematology analyzer) used in this study, the Sahli's technique was reported to be much less satisfactory under operational circumstances, chiefly because of dilution problems involved in the use of ordinary manual pipettes and subjective bias during visual comparison [20,21].

Out of all anemic pregnant mothers, 80.95% (68/84) of them had mild anemia, 17.86% (15/84) had moderate anemia and 1.19% (1/84) had severe anemia according to WHO classification for degree of anemia. In contrast to this study, a study conducted in Kenya and in

83 (21%) every other day and 11 once in a year. One hundred thirty three (33.67%) of the respondents had the habit of eating green leafy vegetables. 127 (32.15%), 90 (22.78%), and 37 (9.36%) had the habit of eating green leafy vegetables every other day, once per week and once in a month respectively. Two hundred eighty eight (72.9%) of the study subjects had the habit of drinking coffee/tea immediately after meal and two hundred sixty three (66.58%) had the habit of eating fruits after meal. Multiple logistic regressions did not show statistically significant association between anemia and any of the dietary habits (Table 4).

Variables		Anemic (N=84) Number (%)	Non-anemic (N=311) Number (%)	Total	AOR(95%CI)	p-value
Gravida	Primi gravida	23(27.38)	110 (35.37)	133	1	
	Multi-gravida	61(72.62)	201(64.63)	262	1.22(1.01,3.42)	0.021
Trimester	<13wks	22 (26.2)	146 (46.95)	168	0.71(0.23,1.63)	0.72
	13-24wks	29 (34.52)	105(33.76)	134	1.1(0.49,1.17)	0.62
	>24wks	33 (39.28)	60 (19.29)	93	2.04(2.65,5.12)	0.41
Parity	Nulli parous	25(29.76)	118 (37.94)	143	1	
	Primipara	26 (30.95)	91(29.26)	117	1	
	Multipara	33 (39.3)	102 (32.8)	135	2.19(3.19,4.32)	0.031
Birth interval	<2years	32(38.09)	61(19.61)	93	1.21(2.16, 5.19)	0.012
	>=2years	29 (34.52)	130 (41.8)	159	1	
Place of delivery	At home	24(28.57)	59 (18.97)	83	0.33(0.11,2.19)	0.576
	At health institution	37 (44.04)	131(42.12)	168	0.88(0.94,3.18)	0.73
Blood loss	Yes	8 (9.52)	13 (4.18)	21	1.67(4.01,20.13)	0.04
	No	53 (63.09)	188 (60.45)	241	1	
Contraception	Yes	35 (41.67)	159 (51.13)	194	0.93(0.73,4.71)	0.069
	No	49 (58.33)	152 (48.87)	201	2.31(1.2,4.41)	0.011
ANC	Yes	57 (67.86)	164 (52.73)	221	0.31(2.32,4.13)	0.07
	No	4 (4.76)	37 (11.9)	41	1.11(1.67,25.1)	0.007
Transfusion	Yes	2 (2.38)	8 (2.57)	10	0.33(0.67,5.19)	0.72
	No	82 (97.62)	303 (97.42)	385	0.59(0.92,4.23)	0.82
Malaria	Yes	2 (2.38)	4 (1.3)	6	1	
	No	82 (97.38)	307 (98.7)	389	0.91(0.18,6.19)	0.71

ANC: antenatal care

Table 3: Distribution of anemia among pregnant women with Obstetric and other variables attending antenatal care at Tikur Anbessa Specialized Hospital from November 23, 2012-March 8, 2013

Eating habit		Anemic (N=84) Number (%)	Non-anemic (N=311) Number (%)	Total	AOR(95%CI)	p-value
Frequency of eating animal foods	I don't eat	0(0)	1(0.32)	1	1.19(0.91,4.32)	0.061
	Daily	7 (8.33)	45 (14.47)	52	1	
	Every other day	14 (16.7)	69 (22.2)	83	0.94(0.23,5.7)	0.072
	Once per week	33 (39.3)	113 (36.33)	146	1	
	Once in a month	25 (29.76)	77 (24.76)	102	1	
Frequency of eating green leafy vegetables	I don't eat	1 (1.19)	6 (1.93)	7	3.21(0.87,6.78)	0.063
	Daily	24(28.57)	66 (21.22)	90	1.12(0.29,3.21)	0.054
	Every other day	29 (34.52)	104 (33.44)	133	1	
	Once per week	26 (30.95)	101(32.48)	127	1.45(0.57,2.19)	0.063
	Once in a month	4 (4.76)	33 (10.61)	37	1	
Taking tea/coffee immediately after meal	Yes	63 (75)	225 (72.35)	288	1	
	No	21(25)	86 (27.65)	107	1.98(0.61,4.32)	0.069
Taking fruit after meal	Yes	53 (63.1)	210 (67.5)	263	1.48(0.97,3.17)	0.076
	No	31(36.9)	101 (32.5)	132	1	

Table 4: Distribution of anemia with dietary habit among pregnant mother attending antenatal care at Tikur Anbessa Specialized Hospital from November 23, 2012- March 8, 2013

Jimma had shown moderate anemia in 68% and 74.3% respectively [11,22]. This inconsistency may be because of the strengthened health education given on risk factors and prevention of anemia and interventions given at health institutions during ANC follow up in an attempt to reduce the prevalence and severity of anemia among pregnant mothers. In addition, it might be because of time and place difference between the present study and the study conducted in Kenya and in Jimma.

Anemia in pregnancy is related to different socio-demographic factors [23]. In different studies, age, educational status, economic position have been found to be significantly associated with anemia during pregnancy [24,25]. This study has assessed socio-demographic variables associated with anemia but only age (39-45 yrs (AOR=1.21)), educational status (illiterate (AOR=2.12)) and family size (greater than four (AOR=1.73)) had shown statistically significant association with anemia which indicates the higher prevalence of anemia in illiterates, large family size and aged pregnant mothers. This high prevalence of anemia in these study participants might be due to inadequate knowledge on factors causing anemia and on how to prevent the risk factors. Similar study conducted on prevalence and risk factors of anemia in rural areas of India and in Jimma showed a statistical significant association between education and anemia and in Turkey, between anemia and large family size which are consistent with this study [16,18,23].

Obstetric factors are known determinants of anemia. In the present study, anemia was 2.04 times more prevalent at third trimester (AOR=2.04). The 26.2% prevalence of anemia at the first trimester increased to 39.28% at the third trimester. This might be due to hemodilution but, it might also indicate poor prenatal care. Anemia was 2.19 and 1.22 times more prevalent in multiparous, multigravidae respectively. This is may be due the fact that multiparity and multigravidity induce anemia by reducing maternal iron reserves at every pregnancy and by causing blood loss at each delivery. There was also a statistically significant association between anemia and history of blood loss (AOR=1.67)), ANC (no ANC (AOR=1.11)) and contraception (no contraception (AOR=2.31)) on multivariate logistic regression analysis which indicates the need for proper follow up of pregnant women starting from the first trimester to the third trimester and the need of attention to multigravidae and multiparous mothers. These findings are consistent with the study conducted in rural areas of Malaysia, in Thailand, in rural areas of Kenya, in the high lands of Tibet and in Jimma which had shown high prevalence of

anemia in multiparous, multigravidae and during the third trimester [11,16,17,26].

Appropriate time after each pregnancy for recuperation and replenishment of nutrient stores and circulating levels is a minimum of three to five years. Pregnancy with a short birth interval leads to iron deficiency anemia as iron requirements are substantially higher than the average [27]. The present study has shown a statistically significant association between birth interval and anemia (birth interval of less than two years (AOR=1.21)). Similar finding has also been documented in a study conducted at Jimma hospital [22]. Thus, this finding indicates the need for strengthening of interventions related to child spacing and awareness [28]. One of the major contributory factors for anemia in developing countries is consumption of plant based food containing insufficient iron, especially insufficient available hem iron from meat [29]. Meat is a good source of high quality protein, iron and zinc and of all the B-vitamins except folic acid. Iron absorption is enhanced when consumed with foods high in vitamin C such as orange juice but substances in coffee and tea inhibit iron absorption [27]. This study has tried to assess different dietary risk factors associated with anemia. Eating animal food, green leafy vegetables, taking fruit after meal and drinking tea/coffee did not show significant association with anemia on multivariate logistic regression which may be due to no difference in eating habit among the study participants.

This study is done only at single institution; hence, further studies have to be conducted in different hospitals of Addis Ababa to have findings representing the whole population. Additionally, further laboratory studies have to be conducted to identify the specific causes of anemia in the pregnant mothers so as to guide the health care givers to work alleviating the existing problems.

Conclusion

The overall prevalence of anemia in this study using a cut off level of haemoglobin <11 g/dl (<33% haematocrit) was 21.3% and the majority of them were of the mild type (haemoglobin: 10-10.9 g/dl). Though the prevalence of anemia in this study is lower than prevalence of anemia in previous studies conducted in other areas of Ethiopia, it still remains higher. The present study has shown a statistically significant association between anemia and age, gestational age, gravidity, parity, antenatal care, birth interval, blood loss, family size and educational status (illiterates). This emphasizes the need for continuing strengthening of interventions on factors associated with anemia.

Acknowledgement

This research was funded by Addis Ababa University Grants for Graduate Research. Therefore, we are grateful to Addis Ababa University, College of Health Sciences for their financial support. Our gratitude also goes to all staff members of Obstetrics and Gynecology of Addis Ababa University for their support and cooperation during the data & sample collection period.

References

1. DeMaeyer E, Adiels-Tegman M (1985) The prevalence of anemia in the world. *World Health Stat Q* 38: 302-316.
2. World Health Organization (1993-2005) Worldwide prevalence of anemia, 49-84.
3. Fairbanks VF (1994) Iron in medicine and nutrition. In *Modern Nutrition in Health and Disease*. (9th edn), 43: 185-213.
4. (1997) UNICEF/South Asia: Malnutrition in South Asia: A Regional Profile. Kathmandu: Rosa Publication. 43: 103-134.
5. Zavaleta N, Berlanga R, Lonnerdal B, Brown H (1993) Prevalence and determinants of iron deficiency anaemia in a representative sample of pregnant women in Lima, Peru, Pan American Health Organization: 54-67.
6. Leenstra T, Kariuki SK, Kurtis JD, Oloo AJ, Kager PA, et al. (2010) Prevalence and severity of anemia and iron deficiency: cross-sectional studies in pregnant women in western Kenya. *Eur J Clin Nutr* 58: 681-91.
7. Bechuram M, Vikal T, Ranjan G (2006) Risk factors of anemia during pregnancy among the Garo of Meghalaya, India. *Human Ecology* 14: S27-S32.
8. World Health Organization (1991) National strategies for overcoming micronutrient malnutrition. Document EB 8.9/27. Executive board, 89th session: 61-83.
9. Baker SJ, DeMaeyer EM (1979) Nutritional anemia: its understanding and control with special reference to the work of the World Health Organization. *Am J Clin Nutr* 32: 368-417.
10. Peters WH (1984) Haematocrit and haemoglobin levels in adult males and in pregnant and non-pregnant females in northern Ethiopia. *Ethiop Med J* 22: 17-27.
11. World Health Organization (1992) The prevalence of anemia in women: a tabulation of available information. (2nd edn), World Health Organization, Geneva, 89-94.
12. Oboro VO, Tabowei TO, Jemikalajah J (2002) Prevalence and risk factors for anaemia in pregnancy in South Southern Nigeria. *J Obstet Gynaecol* 22: 610-613.
13. Singh K, Fong YF, Arulkumaran S (1998) Anemia in pregnancy--a cross-sectional study in Singapore. *Eur J Clin Nutr* 52: 65-70.
14. Sukrat B, Suwathanapisate P, Siritawee S, Pongthong T, Phupongpankul K (2010) The prevalence of iron deficiency anemia in pregnant women in Nakhonsawan, Thailand. *J Med Assoc Thai* 93: 765-770.
15. United Nations Children's Fund Micronutrient Initiative (UNICEF/MI): Vitamin and Mineral Deficiency. A Global Progress Report. Ottawa 2004: 78-83.
16. Desalegn S (1993) Prevalence of anemia in pregnancy in Jima town, southwestern Ethiopia. *Ethiop Med J* 31: 251-258.
17. Xing Y, Yan H, Dang S, Zhuoma B, Zhou X, et al. (2009) Hemoglobin levels and anemia evaluation during pregnancy in the highlands of Tibet: a hospital-based study. *BMC Public Health* 9: 336.
18. Karaoglu L, Pehlivan E, Egri M, Deprem C, Gunes G, et al. (2010) The prevalence of nutritional anemia in pregnancy in an east Anatolian province, Turkey. *BMC Public Health* 10: 329.
19. Hinderaker SG, Olsen BE, Bergsjø P, Lie RT, Gasheka P, et al. (2001) Anemia in pregnancy in the highlands of Tanzania. *Acta Obstet Gynecol Scand* 80: 18-26.
20. Stone JE, Simmons WK, Jutsum PJ, Gurney JM (1984) An evaluation of methods of screening for anaemia. *Bull World Health Organ* 62: 115-120.
21. van Lerberghe W, Keegels G, Cornelis G, Ancona C, Mangelschots E, et al. (1983) Haemoglobin measurement: the reliability of some simple techniques for use in a primary health care setting. *Bull World Health Organ* 61: 957-965.
22. Belachew T, Legesse Y (2006) Risk factors for anemia among pregnant women attending antenatal clinic at Jimma University Hospital, southwest Ethiopia. *Ethiop Med J* 44: 211-220.
23. Federal Ministry of Health. Planning and Programming Department (2004) Health and health related indicators. Addis Ababa: Planning and Programming Department, Federal Ministry of Health, Government of Ethiopia, Ethiopia. 54: 58.
24. Glover-Amengor M, Owusu WB, Akanmori B (2005) Determinants of anaemia in pregnancy in sekere west district, Ghana. *Ghana Med J* 39: 102-107.
25. van den Broek NR, Rogerson SJ, Mhango CG, Kambala B, White SA, et al. (2000) Anaemia in pregnancy in southern Malawi: prevalence and risk factors. *BJOG* 107: 445-451.
26. Haniff J, Das A, Onn LT, Sun CW, Nordin NM, et al. (2007) Anemia in pregnancy in Malaysia: a cross-sectional survey. *Asia Pac J Clin Nutr* 16: 527-536.
27. Woldemariam G, Timotiows G (2002) Determinants of Nutritional Status of Women and Children in Ethiopia, Calverton, Maryland, USA: 49-57.
28. Reproductive Health (2008) New findings on birth spacing: three to five years is the optimal interval, server programs birth spacing optimal interval, 49-58.
29. van den Broek N (2003) Anemia and micronutrient deficiencies. *Br Med Bull* 67: 149-160.

Citation: Jufar AH, Zewde T (2014) Prevalence of Anemia among Pregnant Women Attending Antenatal Care at Tikur Anbessa Specialized Hospital, Addis Ababa Ethiopia. *J Hematol Thromb Dis* 2: 125 doi: [10.4172/2329-8790.1000125](https://doi.org/10.4172/2329-8790.1000125)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>