

Prevalence of Psychiatric Disorders in a Sample of Elderly Residents in Rural and Urban Population of Zulfi Region - Saudi Arabia

Abdul Rahman Al Atram*

Department of Psychiatry, College of Medicine, Majmaah University, Saudi Arabia

Abstract

The population of the world is aging rapidly, because of increasing life expectancy and falling birth rates. This age is also more prone to have chronic diseases and hence there is a need to ensure that the health and social support to these individuals is available. Mental illness represents an important public health problem. Local-level data concerning mental illness in different populations provides the evidence-base for public health authorities to plan, implement and evaluate control programs.

Aims: to estimate the prevalence of psychiatric disorders among the elderly in this region and to analyze for differences between urban and rural areas and also identify and suggest some risk factors for these disorders.

Methodology: The study was done in Zulfi region of Saudi Arabia. The sample size of rural & urban study was 550&392 persons from 65 years and more, respectively.

Instruments and procedures: All the participants of the study were given a general health questionnaire, a symptom check list, a questionnaire to assess the social and economic level and a Structured clinical interview for DSM-IV (SCID)

Results: The psychiatric symptoms among elderly in this study according to SCL 90 were not seen in 15.56% in urban people and 40.4% in rural area. The most prevalent psychiatric disorder in urban area is dysthymia 19.37% while the most prevalent psychiatric disorder in rural area is adjustment disorder with depressed mood at 14%.

Keywords: Psychiatric morbidity; Geriatric; Rural; Urban

Introduction

The global population is aging rapidly, this is due to increasing life expectancy and falling of birth rates, especially in the developed world. The world's older population has a net increase of 1.2 million each month. It is estimated that by the year 2025 the total elderly population will be 976 million [1]. The ageing of the world's population means that health problems associated with the elderly are becoming relatively more important. This growing group has its specific health and social problems which need to be addressed accordingly.

Old age is not a disease, it is a phase of the life cycle characterized by its own developmental issues, many of which are concerned with loss of physical agility, mental acuity, friends and loved ones, status and power [2,3]. This age is also more prone to have chronic diseases and hence, there is a need to ensure that the health and social support to these individuals is available. Mental illness represents an important public health problem. Local-level data concerning mental illness in different populations provides the evidence-base for public health authorities to plan, implement and evaluate control programs [4]. Habib et al. in 2009 reported that 45.5% of prevalence of depressive symptoms were present among Bahraini elders attending local health centers [5].

Saudi Arabia has population growth rates which rank above the general average of 2.37%, registered across the Arab world [6]. As a result of dedicated and intensive efforts towards health care issues the mortality rate has been reduced significantly resulting in increasing percentage of elderly population. Many studies have been done previously to estimate the prevalence of mental disorders with rates varying in different populations, age groups, times, and geographic locations. Psychiatric morbidity in primary care was estimated in 1995 around 30-46% of the visiting patients [7]. In 2002, depression and anxiety disorders were noted about 18% among adults in central Saudi Arabia [8]. Al Ibrahim et al., in 2010 showed an overall prevalence of

41% in a systematic review on depression [9], El Rufaie et al. in 2009 noted a 17% prevalence of depression among residents of Dammam [10]. Al Qahtani et al., in 2008 reported a 27% prevalence of depression in Asir [11]. Abdul Wahid et al. in 2011, reported an overall prevalence of depression nearing 12%, with 6% as severe cases, in the south-eastern region [12]. In Riyadh, Becker et al in 2002 and 2004, found depression prevalence to be 20% in primary care settings [13].

As very few studies have focused on the elderly population [14,15], there is a lacuna in data of mental disorders in elderly in Zulfi region. This study was designed to estimate and compare the prevalence of psychiatric disorders among the elderly between urban and rural areas and also identify some risk factors for these disorders.

Materials and Methods

This study was conducted from October 2012 until October 2013. It was designed as a population based cross-sectional study, in which the city of Zulfi was taken as the urban population and Al-Artaweyah region near Zulfi which is a rural community, represented rural sample. Medical and Dental students of our university were trained regarding the General Health Questionnaires (GHQ) and symptom

*Corresponding author: Abdul Rahman Al Atram, Department of Psychiatry, College of Medicine, Majmaah University, Kingdom of Saudi Arabia, Tel: +966505214617; E-mail: a.atram@mu.edu.sa (or) dr_atram@yahoo.com

Received November 23, 2014; Accepted February 16, 2015; Published February 23, 2015

Citation: Al Atram AR (2015) Prevalence of Psychiatric Disorders in a Sample of Elderly Residents in Rural and Urban Population of Zulfi Region - Saudi Arabia. J Psychol Psychother 5: 170. doi: 10.4172/2161-0487.1000170

Copyright: © 2015 Al Atram AR. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

check list (SCL 90) and these students were instructed to interact with elderly individuals in their respective areas. The data was collected by systematic random sampling technique, 1000 questionnaires were distributed out of which 950 were returned, 58 forms were rejected as they were incomplete. The individuals who rejected the interview or had severe medical or mental illness such as personality disorders, mental retardation etc. which interfered with communication was excluded from the study. Finally a total of 892 elderly of 65 years and older formed the sample size. 392 respondents (262 males & 130 females) formed the urban sample, while rural sample consisted of 500 respondents (278 males & 222 females).

All 892 elderly subjects with scores above cut off point -7 in GHQ and SCL 90 were interviewed with structured clinical interview for DSM -IV to diagnose the disorders. The data was entered and analyzed using SPSS 22.0. Mean ± S.D was reported for quantitative variables. Frequencies and percentages were reported for qualitative variables. Two independent sample t test was applied to compare group mean differences. ANCOVA was applied to compare group means and interaction effects. A p-value of <0.05 was considered as statistically significant.

Results

The demographic data of all 892 subjects reveals that most of the parameters for the urban and rural population are the same however the rural population has a remarkably high percentage of divorcees (35.6%) and the high percentage of low socio economic group (79.2%) (Table 1). However the results of covariance and t test analysis revealed that there are no statistically significant differences between the prevalence of mental disorders among the two groups regarding to the variables such as age, education level, marital status, monthly income, age and gender (Table 2).

Variable	urban		rural		
	n	%	n	%	
Age	65-69	141	35.97	144	28.8%
	70-74	87	22.19	98	19.6%
	75-79	55	14.03	90	18%
	80-85	65	16.58	88	17.6%
	>85	44	11.23	80	16%
Sex	Male	262	66.84	278	55.6%
	Female	130	33.16	222	44.4%
Education	Illiterate	311	79.34	312	62.4%
	Medium	16	4.08	176	35.2%
	High	65	16.58	12	2.4%
work	Working	181	46.17	268	53.6%
	Not working	211	53.83	232	46.4%
Marital status	Married	214	54.59	300	60%
	Divorced	34	8.67	178	35.6%
	Widow	112	28.57	22	4.4%
	Single	32	8.16	0	0
Socio-economic status	low	209	53.32	390	78%
	Average	133	33.93	100	20%
	High	50	12.76	10	2%
Sum	892	392	100	500	100

Socio economic status: monthly income in Saudi riyals
low -less than 5000

medium - 5000 to 15000

high - more than 15000

Education level:

Medium - upto 9th grade ; High - 10th grade and above

Table 1: Demographic and social data of the samples.

Variables	Urban Area		p-value
	Male Mean ± S.D N=262	Female Mean ± S.D N=130	
Gender	3.789 ± 0.61	3.785 ± 0.49	0.967
Work	Yes Mean ± S.D N=181	No Mean ± S.D N=211	0.477
	3.77 ± 0.61	3.81 ± 0.49	
Variables	Rural Area		p-value
	Male Mean ± S.D N=278	Female Mean ± S.D N=222	
Gender	3.41 ± 0.72	3.56 ± 0.77	0.566
Work	Yes Mean ± S.D N=268	No Mean ± S.D N=232	0.424
	3.77 ± 0.61	3.81 ± 0.49	

Table 2: Results of t-test for the variable sex and work.

SCL	Urban		Rural	
	N	%	n	%
Depressive symptom	61	15.56	52	10.4
Somatization	39	9.95	140	28
Psychotic symptom	41	10.46	8	1.6
Paranoid	34	8.67	10	2
Anxiety	39	9.95	50	10
OCD	39	9.95	14	2.8
Phobia	27	6.89	12	2.4
Virtualparanoia	22	5.61	10	2
Sensitiveinteractive	29	7.40	10	2
Negative	61	15.56	202	40.4
Sum	392	100.00	500	100

Table 3: distribution of the sample according to SCL 90.

The GHQ scores were positive for a high percentage of population (79.2%) in rural population as compared to urban (64.54%).

72.07% of the total sample was positive for presence psychiatric symptoms. A high percentage of population (40%) in the rural sample shows no symptoms of psychiatric disorders as compared to 15.56% of urban. Somatization symptom was the most common in rural group (28%) whereas the depressive symptom was common in urban (15.56%) (Table 3).

Psychotic symptoms and paranoid are more in urban population as compared to the rural sample.

According to SCID, dysthymia was the most found disorder in the urban group (19.37%) whereas in rural sample it was adjustment disorder with depressed mood at 14%. Major depressive episode and OCD were more in the urban population at 17.79% and 4.35% than in the rural population; specific phobias were more in the rural group than in the urban group (Table 4).

Discussion

This study showed that the rural population had more percentage of people who were negative for prevalence of mental disorders than the urban population. This was consistent with similar studies done earlier [16], this difference could be due to the variation in lifestyle, social and cultural settings. The demographic data obtained from the study shows that the rural population had more percentage of people of more than 85 years of age; this can be correlated to the less

SCID 1	urban		rural	
	n	%	N	%
schizophrenia	6	2.37	4	1.01
Delusional disorder	7	2.77	8	2.02
Major depressive episode	45	17.79	37	9.34
Adjustment disorder with depressed mood	0	0	70	17.68
Bipolar 1 disorder	6	2.37	5	1.26
Dysthymia	49	19.37	30	7.58
Specific phobias	5	1.98	30	7.58
GAD	22	8.70	40	10.10
PTSD	7	2.77	4	1.01
Panic	5	1.98	3	0.76
Hypomania	2	0.79	5	1.26
OCD	11	4.35	6	1.52
Alcohol & drug abuse disorder	0	0	5	1.26
Negative	88	34.78	149	37.63
Sum	253	100.00	396	100

Table 4: Prevalence of psychiatric disorder.

prevalence of mental disorders in this group and overall healthy living habits seen in rural areas. The socioeconomic status of the rural group shows 78% of the population is in low status and this is consistent with earlier studies [17] which show less prevalence of psychiatric disorders in lower socioeconomic status group. The urban group had 8.67% of people who were divorced but the rural population had a considerably high percentage of people 35.6% who were divorced. This finding is not consistent with earlier studies like M. Mohammadi et al. [16] which have reported that married individuals are less likely to have psychiatric disorders as compared to single and divorced individuals, whereas in our study the rural group which has a high percentage of divorcees also shows less prevalence of psychiatric disorders. Although the differences mentioned above were not statistically significant, there was a large variation in the distribution of psychiatric disorders between the urban and rural groups. This is an interesting finding and it needs to be studied in detail. the lack of statistical significance could be due to small size of sample or could be due to some hidden bias in the sample selection.

Depression was the most common symptom in urban population and was more than that seen in rural group, This was consistent with studies like Al Sehri et al. [18] which have shown that depression is less prevalent in rural population. Anxiety was seen in both the population groups in similar percentage of people (9.95% and 10%), this was considerably less than that seen in previous studies.

Many differences are seen between the urban and rural group in their demographic and social data with respect to the education level, marital status, socio-economic status etc., which is reflected in the difference in prevalence of various psychiatric disorders in the two groups hence these variables could be the possible risk factors in development of psychiatric disorders in elderly and need to be further evaluated as the lack of statistical significance could be because the sample of our study was small, the area targeted was a small region and also there could be a bias in selection of sample as the initial data was collected by medical and dental students and not professionals related to psychiatry. These limitations could also explain the high prevalence of psychiatric symptoms when compared to previously mentioned studies, in spite of this we have obtained a valuable data on the mental disorders in the elderly.

Recommendations

This data shows that the elderly population has more symptoms of psychiatric disorders hence more care should be taken of this population group.

As elderly persons are less likely to seek medical help, more home surveys and camps should be conducted to evaluate the elderly. This data could also be used by the mobile medical units of the university to deliver psychiatric care and implement preventive strategies.

To conduct a more extensive research by psychologists and social workers, with a larger sample size, use of specific scales to detect individual disorders like Hamilton scale for depression, to also correlate with co morbid conditions like diabetes, hypertension etc.

To appoint more psychiatrists in rural areas to make psychiatric care available to all elderly people.

Further investigations are needed to conduct to discuss relationship between elderly and younger population and also how socio-economic supports could the health status of the targeted population.

References

- World Health statistics Annual (1987) World Health Organization, Geneva.
- Hybels CF, Blazer DG (2003) Epidemiology of late life mental disorders. *Clin Geriatr Med* 19: 663-696.
- United Nations (2003) The sex and age distribution of the world populations. United Nations, New York.
- Reeves WC, Lin JM, Nater UM (2013) Mental illness in metropolitan, urban and rural Georgia populations. *BMC Public Health* 13: 414.
- Fatima Habib (2009) Incidence of Depression among Elderly Attending Primary Health Care Centers. *Bahrain Medical Bulletin* 31: 1-7.
- World population prospect: Estimates & projections as assessed in 1982 (1985) New York, United Nation.
- Faris EA, Hamid AA (1995) Hidden and conspicuous psychiatric morbidity in Saudi primary health care. *Arab J Psychiatry* 6: 162-175.
- Al-Khathami AD, Ogbeide DO (2002) Prevalence of mental illness among Saudi adult primary-care patients in Central Saudi Arabia. *Saudi Med J* 23: 721-724.
- Alibrahim O, Al-Sadat N, Elawad N (2010) Gender and risk of depression in Saudi Arabia, a systematic review and meta-analysis. *Journal of Public Health in Africa* 1.
- El-Rufaie OE, Albar AA, Al-Dabal BK (1988) Identifying anxiety and depressive disorders among primary care patients: a pilot study. *ActaPsychiatrScand* 77: 280-282.
- Alqahtani MM, Salmon P (2008) Prevalence of somatization and minor psychiatric morbidity in primary healthcare in Saudi Arabia: a preliminary study in Asir region. *J Family Community Med* 15: 27-33.
- Abdelwahid HA, Al-Shahrani SI (2011) Screening of depression among patients in family medicine in Southeastern Saudi Arabia. *Saudi Med J* 32: 948-952.
- Becker SM (2004) Detection of somatization and depression in primary care in Saudi Arabia. *Socio Psychiatric Epidemiology* 39: 962-966.
- Abolfotouh MA (2001) Psychosocial Assessment of Geriatric Subjects in Abha City, Saudi Arabia. *Eastern Mediterranean Health Journal* 7: 481-491.
- AL-Shammari SA, Al-Subaie A (1999) Prevalence and Correlates of Depression among Saudi Elderly. *International Journal of Geriatric Psychiatry* 14: 739-747.
- Mohammad-Reza Mohammadi, Haratoon Davidian (2005) An epidemiological survey of psychiatric disorders in Iran. *Clinical Practice and Epidemiology in Mental Health* 1: 16.
- Carles Muntaner, William W Eaton (2004) Socioeconomic Position and Major Mental Disorders. *Epidemiologic Rev* 26: 53-62.
- Shaher Z. Al-Shehri, Amr A. Sabra (2012) Depression and Anxiety among Males Attending Primary Health Care Centers, Eastern Saudi Arabia: Prevalence and Predictors. *Life Science Journal* 9: 128-133.