

Prevalence of Toxoplasmosis in Sheep, Goats and Cattle in Southern Tunisia

Ibtissem Lahmar¹, Arwa Lachkhem¹, Darine Slama¹, Wahiba Sakly¹, Najoua Haouas¹, Mohamed Gorcii^{1, 2}, Alexander W Pfaff³, Ermanno Candolfi³ and Hamouda Babba^{1*}

¹Laboratoire de Parasitologie-Mycologie Médicale et Moléculaire (code LR12ES08), Département de Biologie Clinique B, Faculté de Pharmacie de Monastir, Université de Monastir, Tunisia

²Laboratoire de Parasitologie-Mycologie Hôpital Universitaire Fattouma Bourguiba, Monastir, Tunisie

³Institut de Parasitologie et de Pathologie Tropicale, EA 7292, Fédération de Médecine Translationnelle, Université de Strasbourg, France

*Corresponding author: Hamouda Babba, Laboratoire de Parasitologie-Mycologie Médicale et Moléculaire (code LR12ES08), Département de Biologie Clinique B, Faculté de Pharmacie de Monastir, Université de Monastir, Tunisie, Tel: +216 97850853; Fax: + 216 73 496 968; E-mail: hamouda.babba@rns.tn

Received date: September 16, 2015; Accepted date: October 09, 2015; Published date: October 12, 2015

Copyright: © 2015 Lahmar I et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The prevalence of *Toxoplasma gondii* infection in ruminants (sheep, goats, and cattle) in Southern Tunisia is largely unknown.

Antibodies and DNA to *T. gondii* were determined in serum and apex of the heart samples of 261 animals (204 sheep, 32 goats and 25 cattle) using the modified agglutination technique test (MAT) and the PCR respectively.

Antibodies to *T. gondii* (MAT, 1:20) were found in 40.2% (95% CI: 33.4%, 47.2%), 34.5% (95% CI: 19.1%, 53.2%) and 12% (95% CI: 03.15%, 32.13%) in sheep, goats and cattle respectively. Seroprevalence significantly ($p < 0.05$) varied with species, gender, age, and animals' breed. Older animals (>3 years) and females were significantly ($p < 0.05$) more infected than younger and male animals respectively. The seroprevalence was highest in sheep and specifically in the Barbarine breed. The presence of *Toxoplasma* DNA was detected in 11 of 96 samples (11.5%).

The prevalence observed in the present study indicates a widespread exposure to *T. gondii* in South Tunisia. Results indicate, for the first time in Tunisia, that infected sheep and goats represent a potential source of *T. gondii* infection for humans in particular in the South of the country (Gafsa area).

Keywords: *Toxoplasma gondii*; Animals; Prevalence; South Tunisia

Materials and Methods

Introduction

Toxoplasmosis is one of the most common parasitic zoonosis, caused by the obligate intracellular protozoan *T. gondii*, which can infect almost all warm-blooded animals, including humans and domestic animals [1,2]. Toxoplasmosis is also a public health problem owing to its transmission to humans by ingestion of undercooked meat containing tissue cysts, or by consuming food or drink contaminated with oocysts, or through accidental ingestion of sporulated oocysts from the environment [3]. Meat from *T. gondii*-infected mammals is consumed widely in many countries, including Tunisia, and is known to be source of infection for humans [4,5]. Toxoplasmosis leads to great economic losses in ruminants especially in sheep, cattle and goats by causing embryonic death, fetal death, abortion, stillbirth and reduced flock milk production [6]. Worldwide prevalence of *T. gondii* infection in sheep, goats and cattle are largely investigated. However, in Tunisia very few studies have been conducted in all of Tunisia aiming to detect *T. gondii* infection in horses and sheep [2,7,4,8]. Therefore, the knowledge about the seroprevalence of *T. gondii* in domestic ruminants is of interest in order to implement future strategies on public health programs and to clarify the role of ruminants as a source of infection.

The objective of the present study was to investigate the prevalence of *T. gondii* in sheep, goats and cattle in Gafsa, Southern Tunisia.

Animals' blood and heart apex

Blood and heart apex samples were collected from 261 ruminants belonging to three species (sheep n=204, goats n=32 and cattle n=25) at the Gafsa abattoir between December 2010 and October 2012. Each sample is accompanied by a collection of epidemiological data.

The animal population was represented by 102 males and 159 females. The different ages of animals were classified into three groups: the first one including young animals [<1 year-old] (n=152), the second including [1 to 3 years-old] animals (n=37) and the third including the older than 3 years-old (n=72), the oldest animal being nine years old. Two sheep breeds were evaluated: the Barbarine breed (n=15) and the Queue Fine de l'Ouest breed (n=189), while only one breed was tested for each of the other species: the cross breed Bastard for goats and the Frison Pie Noir for cattle. Sera were obtained after 3000 rpm centrifugation for 10 minutes and stored at -20°C until testing for anti-*T. gondii* antibodies.

The apex of the heart (10 g) were removed under sterile conditions with a scalpel blade disposable, then kept in sterile 1.5 ml tubes with the same label as the blood sample from the same animal and kept at -20°C.

All animals were handled in strict accordance with good animal practice and the blood samples were drawn via the jugular vein from

each animal at the time of slaughter by exsanguinations as approved by the Tunisian National Ethics Committee.

Serological examination

Serum samples were analyzed using the Modified agglutination technique test (MAT) (Toxo-Screen[®], BioMerieux) to determine the presence of IgG antibodies against *T. gondii*, according to the manufacturer's protocol for the detection of *T. gondii* antibodies in animals. The test was considered positive when a layer of agglutinated parasites formed in wells at dilution of 1:20, while the control antigen and the negative sera showed sediment as a spot or a small ring respectively. Positive and negative control sera were provided in the kit and were included in each test.

DNA extraction and amplification

DNA was extracted from 25 mg of the homogenized heart apex tissue using the QIAmp DNA tissue Mini Kit isolation kit (Qiagen, France) and stored at -20°C. As positive control, *T. gondii* tachyzoites (RH strain) were obtained from mouse ascites. The presence of *T. gondii* was detected by PCR using the nucleotide sequence of the AF146527 gene as target. Primers were 5'-CTG CAG GGA GGA AGA CGA AAG TTG3' (sense) and 5'-CTG CAG ACA CAG TGC ATC TGG ATT3' (antisense), which amplify a 529 bp fragment of the target gene. PCR reaction was performed in 50 µL reaction mixture containing 10 µL of sample DNA diluted 1:10, 2 µM of each primer, 200 µM dNTP (Invitrogen), 2.5 mM MgCl₂ and 1.25 Units of Hot Start Taq DNA Polymerase (Qiagen). Amplification was performed on a Perkin Elmer thermo cycler (GeneAmpPCR system™ 2400) by 15

min incubation at 94°C, followed by 40 cycles of 30 sec at 95°C, 30 sec at 58.5°C, 30 sec at 72°C and a final 5 min at 72°C. PCR products were analyzed by electrophoresis on 2% agarose gel and visualized on a UV screen using ethidium bromide staining.

Statistical analysis

Statistical analysis and description of *T. gondii* seroprevalence in different species, gender ages and breeds were performed using the software SPSS.v18, M.S. Excel while the Chi square test was calculated with the EPI Info 6.04 software at 95% confidence interval. The differences were considered statistically significant when P<0.05.

Results

Among the 261 animals tested for IgG anti-*T. gondii*, 96 showed seropositivity, leading to an overall seroprevalence of 36.8% (95% CI: 30.9%, 42.9%). Seroprevalence of *T. gondii* varied significantly (p<0.01) among species: 40.2%, 34.5% and 12% in sheep, goats and cattle respectively (Table 1). Moreover, the statistical analysis showed that gender, age and breed affected seroprevalence of *T. gondii* infection. Seroprevalence was higher in females (44%) (95% CI: 36.2%, 52.1%) than in males (25.5%) (95% CI: 17.6%, 35.2%) (p<0.01) (Table 1). In addition, percentage of seropositive animals older than 3 years-old (70.8%) was significantly (p<0.01) greater than those of middle-aged (29.7%) and young (22.3%) animals (Table 1). Considering the sheep breeds, statistical analysis showed a higher infection rate among the Barbarine breed (Table 1). The analysis of 96 DNA samples that had a positive serology showed the presence of Toxoplasma DNA in 11 samples of cardiac apex (11.5%).

Factors	Category	Sample size	Number with anti- <i>T. gondii</i> antibodies	Seroprevalence (%)	95% CI	P value
Species	Sheep	204	82	40.2	[33.4-47.2]	p<0.05
	Goats	32	11	34.50	[19.2-53.2]	
	Cattle	25	3	12	[03.1-32.3]	
Gender	Female	159	70	44	[36.2-52.1]	p<0.001
	Male	102	26	25.50	[17.6-35.2]	
Age (years)	[<1 year]	152	34	22.36	[16.2-29.9]	p<0.0001
	[1-3years]	37	11	29.71	[16.4-47.1]	
	[>3years]	72	51	70.84	[58.8-80.6]	
Sheep Breed	Barbarine	15	12	80	[51.3-94.6]	P=0.0002
	Queue Fine de l'Ouest	188	70	37.23	[30.3-44.6]	

Table 1: Seroprevalence of *Toxoplasma gondii* infection in ruminants in the South of Tunisia.

Discussion

In our study we assessed, for the first time, the prevalence of *T. gondii* infection in ruminants in the Gafsa region, in Southern Tunisia. The overall seroprevalence was about 37%. In sheep, seroprevalence of anti-*T. gondii* antibodies was 40 % which was higher than that found in the north (1.8%) and the center of the country (19%) as reported by Gharbi et al. [4]. This discrepancy may be related to the difference between the number and the age of population studied and the

methods used. Indeed, previous studies were carried out using an ELISA technique which is less sensitive than the agglutination technique used in the present study [9]. The observed seroprevalence in sheep in the current study was higher compared to those reported in other parts of the world, such as Brazil (22.1%) [10], Marrakech (27.6%) [11] and Mexico (29.9%) [12]. However, similar prevalences were reported from the West Indies (44.1%) [13] and Norway (44.3%) [14]. Remarkably, we found a much higher prevalence in Barbarine

sheep (80%, compared to 37%). This result was in agreement with those of Van der Puije et al. [15], who showed a significant difference between breeds. Seropositivity for *T. gondii* in goat (35%) was nearly similar to that reported in Iran (30%) and Brazil (39%) using the indirect immunofluorescence antibody test (IFAT) [16,17]. While seroprevalence is lower in China (14.1%) [18], it is reportedly higher in Ethiopia (74.8%) [19] and in Saudi Arabia (51.7%) [20]. Various risk factors (age, sex, breed, diagnostic test, and climatic variations) may contribute to the difference in seroprevalence in the current study and other studies of the world.

Regarding cattle, the seroprevalence (12%) of toxoplasmosis was higher than that found in West Indies (8.4%) [13], Portugal (7.5%) [21] and Iran (0%) [16], but is comparable to that reported in Poland (12.8%) [22]. Furthermore, this is first report on the seroprevalence of *T. gondii* infection from Tunisian cattle. The difference in prevalence rates between countries is possibly due to different husbandry methods used in these regions. Our findings confirm previous studies stating that seroprevalence of *T. gondii* was highest in sheep, followed by goats and then cattle [15,16,13,21]. Differences in prevalence of toxoplasmosis between animal species might be explained by difference in susceptibility to *T. gondii* infection [23].

Our results showed that older animals (>3 years old) were more likely to be seropositive than younger ones (70% vs 22%, $p < 0.01$). Recently, another Tunisian analysis investigating the seroprevalence of the parasite in ovine meat, reported a similar result (38.2% of young sheep and in 73.6% of adult sheep) [7]. This is probably due to the increasing exposure to the infective oocysts in the environment [10,15]. We also found higher infection rates in female than in male animals, possibly because of higher susceptibility to protozoan parasites of the female than the male [24].

The significant seroprevalence in sheep and goats could be due to the fact that these animals are raised outdoors as grazing animals and could thus have more contact with oocyst shed by cats in the environment. It is also worth mentioning that the oasis of Gafsa was known for the abundance of fresh water, its biodiversity and its traditional irrigation system. In this area, the highest humidity and the vegetation cover protect the oocysts against desiccation which promotes their survival and sporulation. Furthermore, we noted that animals were most affected during the culture months in the time of irrigation water abundance and thus growing herbs yet during winter and summer the animals eat dry feeds which limits parasite transmission.

In addition to the serological study, we determined the presence of toxoplasma DNA in the apex of hearts of different animals studied. Indeed, the heart was considered one of the most prone to the formation of toxoplasma cysts [1] and most accessible collection. In the present study, the overall rate of *Toxoplasma* infection in animals for slaughter was 11.5%. This was close (5.7%) to that found by Khayeche et al. [8] who conducted a study on the hearts of sheep in the center of Tunisia on the occasion of the Muslim feast of sacrifice. However, our prevalence was lower than that showed by recent studies in Tunisia [4,7]. Indeed, these authors found *Toxoplasma* DNA in 50% [7] and 25.5% [4] from the sheep hearts. This discrepancy could be the direct consequence of the difference in the PCR protocols used. In fact, although this technique was highly sensitive, it may give false negative results: (i) the animal may be carrying a low density of parasites in the myocardium, (ii) or the animal hosts the parasite in other organs: the brain, tongue, muscles and retina. However, it could be considered that at least 11.5% of Gafsa slaughtered animals for consumption are

able of transmitting toxoplasmosis infection to humans by ingestion of undercooked or grilled meat or by contamination of hands during meat handling.

In conclusion, the present study showed an enhanced prevalence of *T. gondii* infection in sheep, goats and cattle, for the first time, in the South of Tunisia. Suggesting that consumption of meat of those animals in Gafsa area may represent a potential risk factor for human infection.

Acknowledgments

The authors would like to thank the veterinary Mabrouk Saï for his collaboration in the sampling and supplying information. We also wish to thank Sonia Laabidi and Ahmed Lachkham for their excellent technical assistance. The authors also would like to extend their gratitude to the technical support provided by the laboratory of Parasitology-Mycology of Fattouma Bourguiba Hospital.

References

1. Tenter AM, Heckerroth AR, Weiss LM (2000) *Toxoplasma gondii*: from animals to humans. *Int J Parasitol* 30: 1217-1258.
2. Boughattas S, Bergaoui R, Essid R, Aoun K, Bouratbine A (2011) Seroprevalence of *Toxoplasma gondii* infection among horses in Tunisia. *Parasit Vectors* 4: 218.
3. Alvarado-Esquivel C, Estrada-Martínez S, Pizarro-Villalobos H, Arce-Quiñones M, Liesenfeld O, et al. (2011) Seroepidemiology of *Toxoplasma gondii* infection in general population in a northern Mexican city. *J Parasitol* 97: 40-43.
4. Gharbi M, Zribi L, Jedidi M, Chakkhari H, Hamdi S, et al. (2013) [Prevalence of *Toxoplasma gondii* infection in Tunisian sheep]. *Bull Soc Pathol Exot* 106: 184-187.
5. García-Bocanegra I, Cabezón O, Hernández E, Martínez-Cruz MS, Martínez-Moreno Á, et al. (2013) *Toxoplasma gondii* in ruminant species (cattle, sheep, and goats) from southern Spain. *Journal of Parasitology* 99: 438-440.
6. Dubey JP (2009) Toxoplasmosis in sheep--the last 20 years. *Vet Parasitol* 163: 1-14.
7. Boughattas S, Ayari K, Sa T, Aoun K, Bouratbine A (2014) Survey of the parasite *Toxoplasma gondii* in human consumed ovine meat in Tunis City. *PLoS One* 9: e85044.
8. Khayeche M, Mhadhbi M, Gharbi M, Nasfi I, Darghouth MA (2013) Detection of infection with *Toxoplasma gondii* in sheep slaughtered in the Governorate of Sousse (center-east of Tunisia) during the Muslim Feast of Sacrifice (Eid al-Adha), and analysis of risk factors. *Bulletin of the Society of Exotic Pathology* 107: 60-63.
9. Packham AE, Sverlow KW, Conrad PA, Loomis EF, Rowe JD, et al. (1998) A modified agglutination test for *Neospora caninum*: development, optimization, and comparison to the indirect fluorescent-antibody test and enzyme-linked immunosorbent assay. *Clinical and Diagnostic Laboratory Immunology* 5: 467-473.
10. Andrade MM, Carneiro M, Medeiros AD, Andrade Neto V, Vitor RW (2013) Seroprevalence and risk factors associated with ovine toxoplasmosis in Northeast Brazil. *Parasite* 20: 20.
11. Sawadogo P, Hafid J, Bellele B, Sung RT, Chakdi M, et al. (2005) Seroprevalence of *T. gondii* in sheep from Marrakech, Morocco. *Vet Parasitol* 130: 89-92.
12. Alvarado-Esquivel C, Gayosso-Dominguez EA, Villena I, Dubey JP (2013) Seroprevalence of *Toxoplasma gondii* infection in captive mammals in three zoos in Mexico City, Mexico. *J Zoo Wildl Med* 44: 803-806.
13. Chikweto A, Kumthekar S, Tiwari K, Nyack B, Deokar MS, et al. (2011) Seroprevalence of *Toxoplasma gondii* in pigs, sheep, goats, and cattle from Grenada and Carriacou, West Indies. *J Parasitol* 97: 950-951.

14. Skjerve E, Waldeland H, Nesbakken T, Kapperud G (1998) Risk factors for the presence of antibodies to *Toxoplasma gondii* in Norwegian slaughter lambs. *Prev Vet Med* 35: 219-227.
15. van der Puije WN, Bosompem KM, Canacoo EA, Wastling JM, Akanmori BD (2000) The prevalence of anti-*Toxoplasma gondii* antibodies in Ghanaian sheep and goats. *Acta Trop* 76: 21-26.
16. Sharif M, Gholami Sh, Ziaei H, Daryani A, Laktarashi B, et al. (2007) Seroprevalence of *Toxoplasma gondii* in cattle, sheep and goats slaughtered for food in Mazandaran province, Iran, during 2005. *Vet J* 174: 422-424.
17. Anderlini GA, Mota RA, Faria EB, Cavalcanti EF, Valença RM, et al. (2011) Occurrence and risk factors associated with infection by *Toxoplasma gondii* in goats in the State of Alagoas, Brazil. *Rev Soc Bras Med Trop* 44:157-162.
18. Zhao GH, Zhang MT, Lei LH, Shang CC, Cao DY, et al. (2011) Seroprevalence of *Toxoplasma gondii* infection in dairy goats in Shaanxi Province, Northwestern China. *Parasites & Vectors* 4: 47.
19. Teshale S, Dumètre A, Dardé ML, Merga B, Dorchies P (2007) Serological survey of caprine toxoplasmosis in Ethiopia: prevalence and risk factors. *Parasite* 14: 155-159.
20. Sanad MM, Al-Ghabban AJ (2007) Serological survey on toxoplasmosis among slaughtered sheep and goats in Tabouk, Saudi Arabia. *J Egypt Soc Parasitol* 37: 329-340.
21. Lopes AP, Dubey JP, Neto F, Rodrigues A, Martins T, et al. (2013) Seroprevalence of *Toxoplasma gondii* infection in cattle, sheep, goats and pigs from the North of Portugal for human consumption. *Vet Parasitol* 193: 266-269.
22. Sroka J, Karamon J, Cencek T, Dutkiewicz J (2011) Preliminary assessment of usefulness of cELISA test for screening pig and cattle populations for presence of antibodies against *Toxoplasma gondii*. *Ann Agric Environ Med* 18: 335-339.
23. Fayer R (1981) Toxoplasmosis update and public health implications. *Can Vet J* 22: 344-352.
24. Alexander J, Stinson WH (1988) Sex hormones and the course of parasitic infection. *Parasitology Today* 4: 189-193.