

Prevention of Depression: A Review of Literature

Al Qahtani A* and Al Qahtani N

Department of Psychiatry, Imam Abdulrahman Bin Faisal University, Dammam, Kingdom of Saudi Arabia

Abstract

This article provides a review of recent literature, current and ongoing research in the field of prevention of depression. It highlights the efforts employed in targeting vulnerable and susceptible individuals at higher risk of developing depression. Prevention efforts should target both specific and non-specific risk factors, enhance protective factors, use a developmental approach, and target selective and/or indicated samples. In general, our review suggested that employing specific strategies and interventions that are targeted at high-risk individuals or groups may reduce rates of depression, its associated mortality and morbidity. Overall, it appears that there is a reason for hope regarding the role of interventions in preventing depressive disorders in high-risk groups. Several new directions for future research on the prevention of depression in high-risk groups were outlined.

Keywords: Depression; Prevention; Intervention

Introduction

Depression is a state of low mood and aversion to activity or apathy that can affect a person's thoughts, feelings, behavior and sense of well-being. It is a very serious and disabling psychiatric condition that occasionally leads to suicide or premature death due to unattended physical problems. The annual prevalence of major depressive disorder is 6.6 % and the lifetime prevalence of depression is 16.2% [1]. An estimated 676 million (one in ten people) are affected by depression. Worldwide, 804,000 people committed suicide in 2012, making depression one of the leading causes of death in young adults (15-29 years) second to road traffic accidents [2,3]. Persons with major depression have a 40% greater chance of dying prematurely than the general population [2].

The association between depression and increased risk of death and morbidity is an obvious indicator of the severity of the condition, which is usually clear when reflecting upon the WHO data on life expectancy and the causes of death. However, another extremely important indicator of its severity is the healthy life expectancy (HLE). Healthy life expectancy (HLE), if measured reliably, will reflect both mortality and unhealthy life (in case of disability, years lost due to disability -YLD). Global life expectancy in 2015 was 71.4 years. Globally, HLE in 2015 was estimated to be 63.1 years for both sexes. In general, Healthy life expectancy is 11.7% shorter than life expectancy. Life expectancy is comparatively longer in females (73.8) than males (69.1) in every country worldwide. Mental disorders, especially depression, are the second cause of lost healthy life years in all regions of the world. Years lost due to disability (YLD) is mainly due depression, second to musculoskeletal conditions [2].

The economic burden of bipolar disorders on the individuals as well as society is both direct (e.g. inpatient and outpatient treatment cost) and indirect costs (e.g. lost productivity of the patients and their caregiver). In the United States, in 2009, the total cost was estimated to be \$151 billion, \$30.7 billion in direct and \$120.3 billion in indirect costs [4]. Globally, \$2.5-8.5 trillion in lost output was attributed to mental, neurological and substance abuse disorders [5]. These costs include direct cost of treatment as well as the indirect costs related to unemployment and lost productivity at workplace.

The burden of depression is not limited to the human misery of the depressed person. It extends to the caregiver, the partner, the children, and the whole society in general. The quality of life, which is a measure of subjective well-being in different domains of life, is lower in

persons suffering from depression [6]. Caregivers of depressed patients are more likely themselves to suffer from more depressive symptoms, poorer general health and more chronic medical conditions [1].

In a recent study published in the Lancet, the authors propose a global investment case for scaled-up response to the public health and economic burden of depression [5]. The estimated cost for substantially improving the care for depressed patient, over the period 2016-2030, is US\$147 billion. This figure is huge but the expected return for this great investment is huge. Treating depression will lead to an extra 43 million healthy life years. In addition, there will be a large economic gain, a net present value of US\$ 230 billion and US\$ 169 billion for depression and anxiety, respectively [5]. If policy makers invest in preventing depression in the first place, the return for this investment will be greater than treating depression.

Objectives

The aim of this article is to review the most recent literature related to interventions for depression prevention.

Literature Review

For this review, literature was reviewed using Summon search. This is a powerful search method with a google like interface. It is a unified search of all the databases, e-books and e-journals accessible from the Imam Abdulrahman Alfaisal University. These include BMJ journals, Cochrane, Ovid, ProQuest, Cambridge journals, SAGE journals, ScienceDirect journals, Wiley-Blackwell journals and many related databases [6-9].

The initial search conducted, on 15th November 2016, using the key words Depression AND Prevention yielded 542,435 results. The search was then limited to journal articles and this yielded 269,271 articles.

***Corresponding author:** Al Qahtani A, Department of Psychiatry, Imam Abdulrahman Bin Faisal University, Dammam, Kingdom of Saudi Arabia, Tel: +00966590084004; E-mail: abalqhtani@uod.edu.sa

Received August 31, 2017; **Accepted** September 08, 2017; **Published** September 11, 2017

Citation: Al Qahtani A, Al Qahtani N (2017) Prevention of Depression: A Review of Literature. J Depress Anxiety 6: 292. doi:10.4172/2167-1044.1000292

Copyright: © 2017 Al Qahtani A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Language was limited to English and these yielded 255,830 articles. Because this review is mainly covering the most recent literature, the search was limited to the last 12 months (November 2015- November 2016). This limitation yielded 16,214 articles. The final limitation is using only articles related to the subject of psychiatry. This yielded 2089 articles. Careful search through the titles revealed 58 articles related to prevention of depression [10-65].

Risk factors

Before any discussion on risk factors for depression, it is important to make it clear that there is a considerable confusion over what constitute a risk factor and how they operate. Most of the studies are cross-sectional studies using different correlation regression statistical models to find *possible* risk factors. For a proposed risk factor to be established as *probable* cause, prospective studies must be done where the risk factors are measured before the outcome. This proves to be difficult, and even when this is established; more evidence is needed to prove it as a *causal* risk factor. The matter is even more complicated due to the multidimensional view of depression where these risk factors are not isolated and not static.

There are some risk factors, which are setting factors like race, culture, age, gender and socioeconomic status. In general, there are biological, cognitive and social factors that increase the risk of depression. Biological factors include genetic and familial predisposition, alteration in the neural structures and sleep dysregulation and other regulatory methods. Cognitive factors include cognitive schemas, beliefs, assumptions, pessimism, low self-esteem, ruminative response style and negative cognitive style. Excessive reassurance seeking and negative feedback seeking are also direct risk factors for depression

[66]. Other important risk factors include marriage and relationship issues, low social support and low income [67]. Chronic diseases like cardiovascular problems, diabetes mellitus, cancer, disability, and many other health problems can increase the likelihood of depression [68]. Anxiety and prior depression also increase the risk of depression [69]. There is robust evidence that major life events have strong relationship to depression [70]. Also, childhood adversity and negative core beliefs are a strong predictor of later major depression [71]. It heightens the sensitivity to future stress and makes depression more likely. Figure 1 shows schematic representation of risk factors.

World Health Organization (WHO) produced a comprehensive mental action plan. One of the ambitious objectives is the provision of comprehensive, integrated mental health and social care services in community setting and implementation of strategies for promotion and prevention. One of the major objectives of mental health action plan is to promote mental health and prevent mental disorders. To achieve this, it is necessary to provide technical support to help countries in selecting, formulating and implementing evidence-based and cost-effective best practices [72]. The global target is that by 2020, 80% of countries will have at least two functioning national multi-sectoral mental health promotion and prevention programs. If treatment of depression was effective and available to patients, the prevalence of depression should decrease. However, it has been shown that the prevalence is not changing and is not becoming an “epidemic” [73,74].

Prevention interventions

Most researchers and practitioners define prevention as the process of employing different interventions before the people meet the formal criteria, according to DSM-V, of a depressive disorder. There are many




Figure 1: Schematic representation of risk factors.

interventions for depression prevention in high-risk groups. Many studies have been directed to adolescents at higher risk of depression [8-14,16,29,22,26,30,32]. There are also studies on pregnant women in effort to prevent postpartum depression [17,21,29,31]. In the following paragraphs, these preventive interventions will be discussed.

There are basically three type of intervention program delivery:

1. Universal: Preventive intervention programs that are usually targeting large groups e.g. all school students, all pregnant women attending antenatal clinic or all diabetic patients.

2. Selective: Preventive intervention programs targeting subgroups e.g. children of depressed parents, pregnant women in abusive or difficult relationship or diabetic patients with multiple complications.

3. Indicated: Preventive intervention programs targeting high-risk groups for developing depression e.g. adolescents, pregnant women or diabetic patients with sub-threshold depressive symptoms.

The content and venue of these programs are varied and depending on the characteristics of the targeted group(s). The following are the most studied programs:

School-based intervention programs

These programs are mainly for adolescents because they spend most of their time at schools. The following are the main programs that have been used in many countries and are both feasible and effective. These interventions are summarized in several recent meta-analysis and review articles of prevention programs in schools [16,38,75-77].

i. Universal programs

Resourceful adolescent program: consists of 11 weekly sessions of cognitive behavior therapy (CBT) and interpersonal therapy (IPT). They are delivered by psychologist. There was no improvement in the outcome when three family sessions were added [75].

Beyond-blue program: It is a ten-weekly session program delivered by teachers. The goal is to improve the youth protective factors (social and coping skills) and improve school climate [75].

Problem-solving for life program: It is an eight-weekly session program delivered also by teachers. The goal is to improve the adolescents' ability to modify harmful thoughts and use problem-solving skills. This is achieved by cognitive restructuring and problem-solving training [75].

ii. Selective programs

Penn resiliency program: It is a 12-weekly session program delivered by psychology master students to low-income and racial/ethnic minority students. It uses cognitive behavioral skills to address the link between thoughts and emotions to improve cognition and coping [16].

Aussie optimism program: It is a 20-weekly session program delivered by teachers to low-income students. It relies on improving optimistic thinking skills and social life skills. This in turn enhances cognition and social protective factors [16].

iii. Indicated programs

The feelings club: It is a 12-weekly session program delivered by psychologists to students who screened positive for internalization symptoms [16].

Personal growth class: It is a 20-weekly session program delivered by teachers, counselors or nurses to students who screened positive for

suicide risk. The main component in this program is social support and life-skills training [16].

Family-based intervention programs

In this approach, children of depressed patients, with sub-threshold depressive symptoms receive several sessions of cognitive behavioral intervention with focus on cognition reconstruction, effective communication skills and interpersonal problem-solving skills. Some programs address both adolescents and their parents [19].

Internet and computerized intervention programs

The Internet and computerized cognitive behavioral interventions have been shown to lead to reduction in depressive symptoms among men and adolescents aged between 12-25 years [8,13,27,33,36,42,52,78,79]. These methods were not effective for younger children aged 5-11 years [8].

Text messaging intervention program

A recent study, involving adolescents attending emergency department at risk of depression, has shown that brief in-person discussion in the emergency, followed by 8-week automated messaging intervention is feasible and acceptable [65]. Telephone calls has also been used to deliver these interventions for older patients unable to attend the clinic due to pain [58].

Clinic-based prevention program

Depression prevention programs have been offered to high-risk patients suffering from arthritis in the setting of outpatient clinic. The cognitive behavioral therapy program was feasible and acceptable [58]. This approach is also offered to a large variety of people attending the hospital for various reasons including pregnant women for antenatal care, post-traumatic or geriatric patients [18,37,78,80,81].

Community-based prevention program

Depression prevention intervention programs at a community level aims usually to raise awareness about depression. Education and universal screening leads to selective and indicated intervention in at high-risk groups and their caregivers. There are many studies which has proven that this approach is effective, feasible and acceptable to people [17,50,82]. Other studies are still in progress [83].

Content of the intervention programs

The proposed interventions to prevent depression are as diverse as the groups being targeted and the level at which the intervention is made. For example, the study, conducted in Japan to prevent depression among elderly people, starts with universal interventions in the form of raising awareness through education [50]. Then they offered universal screening and those considered at high risk were offered selective intervention in the form of psychosocial follow-up with community support.

The most common intervention in many studies is cognitive-behavioral therapy (CBT) intervention. CBT comprised the basis of 84% of the programs in different interventions. The programs vary from as simple as having more fun activities to combined cognitive behavioral therapy and interpersonal therapy, social skills programs, creative-expressive experiential therapy, mindfulness-based cognition therapy program, wellbeing therapy program and psychoeducational program [75]. The intervention sessions can be built and delivered to meet the needs of the group. Interestingly, even CB bibliotherapy, in the form of self-help book or brochure, can be effective in those with negative attributional style [48,84]. Any intervention program is

S. No	Psychotherapeutic interventions
1.	Cognitive behavioral therapy (CBT)
2.	CBT bibliotherapy
3.	Interpersonal problem-solving skills
4.	Mindfulness-based CBT
5.	Life problem solving skills
6.	Effective communication skills
7.	Supportive-expressive intervention
8.	Active coping skills
9.	Optimistic thinking skills
10.	Combating demoralization
11.	Relaxation skills
12.	Breathing control training
13.	Social skills program
II	Non-psychotherapeutic interventions
1.	Fun and pleasant activity
2.	Improving home dynamics
3.	Improving school climate
4.	Exercise therapy
5.	Nutrition and nutritional supplements
III	Pharmacotherapy
1.	Hydrocortisone
2.	Aspirin
3.	Propranolol
4.	Morphine
5.	Benzodiazepines
6.	Oxytocin

Table 1: Forms of interventions for depression prevention.

usually delivered in weekly sessions each lasting for 45-120 minutes. The content is tailored to the group's needs. In addition to these interventions, participants are usually advised to monitor themselves at home, practice what they learned and practice self-reinforcement. Table 1 shows the most common forms of interventions for depression prevention that have been evaluated in the literature.

There are other less studied interventions related to food or nutritional supplement [11,21,25,41,44,45]. Omega-3 a highly unsaturated fatty acid (HUFAs), number of studies was concerned with the addition of Omega-3 to the treatment regimen of already depressed patients. Which have been shown it to have beneficial effect to alleviate depressive symptoms in patients already receiving antidepressant treatment [35,83,85]. However, the evidence is not conclusive as the results of different studies are not univocal. There is a current underway randomized, double-blind, placebo-controlled study to look at the effect of fish oil in alleviation of depression in young people [7]. The results are expected very soon.

Very rarely, pharmacotherapy is used for prevention of depression. Many drugs have been used in the acute stage of certain circumstances e.g. military combat injury, post-traumatic events and in disaster regions. The drugs that have been used include hydrocortisone, propranolol, benzodiazepines, aspirin, morphine and oxytocin [39,80,82].

Exercise, Yoga and increasing physical activities in general were evaluated regarding their effectiveness in treating or preventing depression [86]. A systematic review of the literature and meta-analysis of the available trials showed a small number of low-moderate quality trials. Only weak evidence that exercise can prevent depression due to the significant heterogeneity and wide confidence intervals [87]. Larger high quality trials are needed to answer this question.

Conclusion

As a matter of fact, depression is a huge burden on the affected individual, the caregiver, the family, the society and the whole world. The associated morbidity, mortality and the enormous cost of dealing with depression, makes the task of depression prevention an urgent matter. This review article suggests that employing specific strategies and interventions that are targeted at high-risk individuals or groups may reduce rates of depression, its associated mortality and morbidity. The scientists have had their share of work in providing the evidence through very extensive trials, that depression prevention is effective, feasible and acceptable. Finally, more longitudinal studies and clinical trials are required to specify the impact of preventive strategies on incidence, prevalence, mortality and morbidity of depression and its treatment.

Pending such research, clinicians should consider both current active pharmacological and psychological treatments and benefits of preventive strategies and approaches to overcome barriers to accessing treatment for depressive disorders. Now it is the duty of everyone to translate this knowledge into an action. Policy makers, media, clinicians, teachers, nurses and midwives all share the responsibility. It starts with universal intervention as simple as just raising the awareness of the public to the magnitude of the problem and the available prevention programs, all the way to designing the suitable intervention program for each high-risk group.

Limitation of the study

This is not a systematic review of the literature related to depression prevention. Systemic reviews were done in specific high-risk groups as it is not feasible to review all intervention in all groups systematically. It is intended to review the most recent literature and to offer broad-based summary of the prevention interventions to general readers.

Acknowledgment

Our sincere thanks and gratitude goes to Dr. Mukhtar Nasir and Prof. Omer M Alakloby for their support, revision of the manuscript and valuable input.

Future Scope

The evidence is compelling that depression can be prevented. The task is to translate this knowledge into an actual action that will change the life of many people and will eventually save life and money. More studies are needed to answer the questions related to other less studied preventive measures like food, nutritional supplements and exercise.

References

1. Miller S, Dell'Osso B, Ketter TA (2014) The prevalence and burden of bipolar depression. *J Affect Disord* 169S1: S3-S11.
2. World Health Statistics (2016) Monitoring health for the SDGs. World Health Organization, Geneva.
3. Preventing Suicide, A global imperative. World Health Organization, Geneva.
4. Dilsaver SC (2011) An estimate of the minimum economic burden of bipolar I and II disorders in the USA. *J Affect Disord* 129: 79-83.
5. Chisholm D, Sweeny K, Sheehan P, Rasmussen B, Smit F, et al. (2016) Scaling-up treatment of depression and anxiety: A global return on investment analysis. *Lancet Psychiatry* 3: 415-24.
6. Miller CJ, Abraham KM, Bajor LA, Lai Z, Kim HM, et al. (2013) Quality of life among patients with bipolar disorder in primary care versus community mental health settings. *J Affect Disord* 146: 100-105.
7. Rice SM, Hickie IB, Yung AR, Mackinnon A, Berk M, et al. (2016) Youth depression alleviation: The fish oil youth depression study (YoDA-F): A randomized, double-blind, placebo-controlled treatment trial. *Early Interv psychiatry* 10: 290-299.

8. Richards K, Marko-Holguin M, Fogel J, Anker L, Ronayne J, et al. (2016) Randomized clinical trial of an internet-based intervention to prevent adolescent depression in a primary care setting (catch-it): 2.5-year outcomes. *Journal of Evidence - Based Psychotherapies* 16: 113.
9. Shomaker LB, Kelly NR, Pickworth CK, Cassidy OL, Radin RM, et al. (2016) A randomized controlled trial to prevent depression and ameliorate insulin resistance in adolescent girls at risk for type 2 diabetes. *Ann Behav Med* 50: 762-774.
10. Lai ES, Kwok CL, Wong PW, Fu KW, Law YW, et al. (2016) The effectiveness and sustainability of a universal school-based program for preventing depression in Chinese adolescents: A follow-up study using quasi-experimental design. *PLoS One*. 11: e0149854.
11. Morin KH (2016) Dietary influence on depression. *Am J Matern Child Nurs* 41: 311.
12. Duffy S, Brown TM, Katsonga-Phiri T, Bouris A, Grant KE, et al. (2016) Development of an empirically based preventive intervention for depression in preadolescent African American girls. *Prev Sci* 17: 503-512.
13. Gladstone TG, Marko-Holguin M, Rothberg P (2015) An internet-based adolescent depression preventive intervention: Study protocol for a randomized control trial. *Trials* 16: 203.
14. La Greca AM, Ehrenreich-May J, Mufson L, Chan S (2016) Preventing adolescent social anxiety and depression and reducing peer victimization: Intervention development and open trial. *Child & Youth Care Forum* 45: 905-926.
15. Nock MK (2016) Recent and needed advances in the understanding, prediction, and prevention of suicidal behavior: Editorial. *Depression & Anxiety* 33: 460-463.
16. Stockings EA, Degenhardt L, Dobbins T, Lee YY, Erskine HE, et al. (2015) Preventing depression and anxiety in young people: A review of the joint efficacy of universal, selective and indicated prevention. *Psychol Med* 46: 11-16.
17. Brugha TS, Smith J, Austin J, Bankart J, Patterson M, et al. (2015) Can community midwives prevent antenatal depression? an external pilot study to test the feasibility of a cluster randomized controlled universal prevention trial. *Psychol Med* 46: 345-412.
18. Jørstad HT, Minneboo M, Helmes HJM, Fagel ND, Peters JG, et al. (2016) Effects of a nurse-coordinated prevention program on health-related quality of life and depression in patients with an acute coronary syndrome: Results from the RESPONSE randomized controlled trial. *BMC Cardiovasc Disord* 16: 144.
19. Zwaanswijk M, Kösters MP (2015) Children's and parents' evaluations of 'Friends for life', an indicated school-based prevention program for children with symptoms of anxiety and depression. *Behavior Change* 32: 243-254.
20. Stallard P (2016) Long-term benefits of cognitive-behavioral prevention for adolescents at risk of depression but not if parents are depressed at the onset of the program. *Evid-based ment health* 19: e20.
21. Kianpour M, Mansouri A, Mehrabi T, Asghari G (2016) Effect of lavender scent inhalation on prevention of stress, anxiety and depression in the postpartum period. *Iran J Nurs Midwifery Res* 21: 197.
22. Bingol F, Buzlu S (2016) Effect of the cognitive-behavioral prevention program on levels of depression symptoms among working adolescents in turkey. *J Psychosoc Nurs Ment Health Serv* 54: 43-51.
23. Lijja JL, Zellerroth C, Axberg U, Norlander T (2016) Mindfulness-based cognitive therapy is effective as relapse prevention for patients with recurrent depression in Scandinavian primary health care. *Scand J Psychol*. 57: 464-472.
24. Martinsen KD, Kendall PC, Stark K, Neumer S (2016) Prevention of anxiety and depression in children: Acceptability and feasibility of the trans diagnostic Emotion program. *Cogn behav pract* 23: 1-13.
25. Okereke OI, Singh A (2016) The role of vitamin D in the prevention of late-life depression. *J Affect Disord* 198: 1-14.
26. Brent DA, Brunwasser SM, Hollon SD, Weersing VR, Clarke GN (2016) Effect of a cognitive-behavioral prevention program on depression 6 years after implementation among at-risk adolescents: A randomized clinical trial. *JAMA Psychiatry* 72: 1110-11118.
27. Wang J, Lam RW, Ho K, Attridge M, Lashewicz BM, et al. (2016) Preferred features of E-mental health programs for prevention of major depression in male workers: Results from a Canadian national survey. *J Med Internet Res* 18: e132.
28. Dieu K (2016) Prevention and intervention of depression in Asian-American adolescents. *Contemporary School Psychology* 20: 107-117.
29. Osma J, Barrera AZ, Ramphos E (2016) Are pregnant and postpartum women interested in health-related apps? implications for the prevention of perinatal depression. *Cyberpsychol Behav Soc Netw* 19: 412-415.
30. Young J, Mufson L (2016) Interpersonal psychotherapy: Adolescent skills training for the prevention of adolescent depression. *J Am Acad Child Adolesc Psychiatry* 55: S39.
31. Dennis C (2016) 'Time for self' appears to be a proactive strategy for the prevention of postpartum depression. *Evid Based Nurs* 19: 114.
32. Brière FN, Rohde P, Stice E, Morizot J (2016) Group-based symptom trajectories in indicated prevention of adolescent depression. *Depression & Anxiety* 33: 444.
33. Buntrock C, Ebert DD, Lehr D, Smit F, Riper H, et al. (2016) Effect of a web-based guided self-help intervention for prevention of major depression in adults with sub threshold depression: A randomized clinical trial. *JAMA* 315: 1854-1863.
34. Perry Y, Calear AL, Mackinnon A, Batterham JP, Licinio J, et al. (2015) Trial for the prevention of depression (TriPoD) in final-year secondary students: Study protocol for a cluster randomized controlled trial. *Trials* 16: 451.
35. McNamara RK (2016) Role of omega-3 fatty acids in the etiology, treatment, and prevention of depression: Current status and future directions. *J Nutr Intermediary Metabolis* 5: 96-106.
36. Angheliescu I (2016) Online intervention for prevention of major depression. *JAMA* 316: 881.
37. Mejía S (2016) Promoting multilevel primary prevention of depression and diabetes during midlife may protect against dementia. *Evid-based ment health* 19: e4.
38. Weersing VR, Shamseddeen W, Garber J, Hollon SD, Clarke GN, et al. (2016) Prevention of depression in at-risk adolescents: Predictors and moderators of acute effects. *J Am Acad Child Adolesc Psychiatry* 55: 219-226.
39. Berk M, Woods RL, Nelson MR, Shah RC, Reid CM, et al. (2016) ASPREE-D: Aspirin for the prevention of depression in the elderly. *Int Psychogeriatr* 28: 1741-1748.
40. Garber J, Brunwasser SM, Zerr AA, Schwartz KTG, Sova K, et al. (2016) Treatment and prevention of depression and anxiety in youth: Test of Cross Over effects. *Depression & Anxiety* 33: 939-959.
41. Roca M, Kohls E, Gili M, Watkins E, Owens M, et al. (2016) Prevention of depression through nutritional strategies in high-risk persons: Rationale and design of the Mood Food prevention trial. *BMC Psychiatry* 16: 192.
42. Rice S, Gleeson J, Davey C, Hetrick S, Parker A, et al. (2016) Moderated online social therapy for depression relapse prevention in young people: Pilot study of a 'next generation' online intervention: Online depression relapse prevention. *Early Interv Psychiatry*.
43. Muñoz RF, Bunge EL (2016) Prevention of depression worldwide: A wake-up call. *The Lancet Psychiatry* 3: 306-307.
44. Opie RS, Itsiopoulos C, Parletta N, Sanchez-Villegas A, Akbaraly TN, et al. (2015) Dietary recommendations for the prevention of depression. *Nutr Neurosci* 150828083143002.
45. Martínez-González MA, Sánchez-Villegas A (2016) Food patterns and the prevention of depression. *Proc Nutr Soc* 75: 139.
46. Mendelson T, Tandon SD (2016) Prevention of depression in childhood and adolescence. *Child Adolesc Psychiatr Clin N Am* 25: 201-218.
47. Bradley KL, Santor DA, Oram R (2016) A feasibility trial of a novel approach to depression prevention: Targeting proximal risk factors and application of a model of health-behavior change. *Canadian J Community Ment Hlt* 35: 47.
48. Müller S, Rohde P, Gau JM, Stice E (2015) Moderators of the effects of indicated group and bibliotherapy cognitive behavioral depression prevention programs on adolescents' depressive symptoms and depressive disorder onset. *Behav Res Ther* 75: 1-10.
49. Haskins J, Carson JG, Chang CH, Kirshnit C, Daniel PL, et al. (2016) The

- suicide prevention, depression awareness, and clinical engagement program for faculty and residents at the University of California, Davis health system. *Acad Psychiatr* 40: 23-29.
50. Vázquez FL, Torres Á, Blanco V, Otero P, Díaz O, et al. (2016) Long-term follow-up of a randomized clinical trial assessing the efficacy of a brief cognitive-behavioral depression prevention intervention for caregivers with elevated depressive symptoms. *Am J Geriatr Psychiatry* 24: 421-432.
 51. Otero P, Smit F, Cuijpers P, DeRubeis RJ, Torres Á, et al. (2015) Differential response to depression prevention among a sample of informal caregivers: Moderator analysis of longer-term follow-up trial data. *Psychiatry Res* 230: 271.
 52. Patrick IP, Chim D, Chan KL, Tim MH, Tsang A, et al. (2016) Effectiveness of a culturally attuned internet-based depression prevention program for Chinese adolescents: A randomized controlled trial. *Depression & Anxiety* 33: 1123-1131.
 53. Drury SS, Scaramella L, Zeanah CH (2016) The neurobiological impact of postpartum maternal depression: Prevention and intervention approaches. *Child Adolesc Psychiatr Clin N Am* 25: 179-200.
 54. Juangco DNA (2016) Depression prevention in caregivers of patients with dementia: A meta-analysis. *Alzheimer's & Dementia* 12: P785.
 55. Young JF, Benas JS, Schueler CM, Gallop R, Gillham JE, et al. (2016) A randomized depression prevention trial comparing interpersonal Psychotherapy-Adolescent skills training to group counseling in schools. *Prevent Sci* 17: 314-324.
 56. Tak YR, Lichtwarck-Aschoff A, Gillham JE, Zundert, RMP Van Engels, et al. (2016) Universal school-based depression prevention 'Op volle kracht': A longitudinal cluster randomized controlled trial. *J Abnorm Child Psychol* 44: 949-961.
 57. Benas JS, McCarthy AE, Haimm CA, Huang M, Gallop R, et al. (2016) The depression prevention initiative: Impact on adolescent internalizing and externalizing symptoms in a randomized trial. *J Clinical Child Adolescent Psychol* 1-15.
 58. Karp JF, Dew MA, Wahed AS, Fitzgerald K, Bolon CA, et al. (2015) Challenges and solutions for depression prevention research: Methodology for a depression prevention trial for older adults with knee arthritis and emotional distress. *Am J Geriatr Psychiatry* 24: 433-443.
 59. Rohde P, Stice E, Shaw H, Gau JM (2016) Pilot trial of a dissonance-based cognitive-behavioral group depression prevention with college students. *Behav Res Ther* 82: 21-27.
 60. Duong MT, Kelly BM, Haaland WL, Matsumiya B, Huey SJ, et al. (2016) Mediators and moderators of a school-based cognitive-behavioral depression prevention program. *Cognitive Ther Res* 40: 705-716.
 61. Tomy JD, Fuller-Tyszkiewicz M, Richardson B, Colla L (2016) A comprehensive evaluation of a universal school-based depression prevention program for adolescents. *J Abnorm Child Psychol* 44: 1621-1633.
 62. Werner EA, Gustafsson HC, Lee S, Feng T, Jiang N, et al. (2016) PREPP: Postpartum depression prevention through the mother-infant dyad. *Arch Womens Ment Health* 19: 229-242.
 63. Young J, Mufson L, Gillham JE, Benas J, Schueler CM, et al. (2016) 48.1 the depression prevention initiative: Effects on depression symptoms and diagnoses. *J Am Acad Child Adolesc Psychiatr* 55: S335.
 64. Suffoletto B, Aguilera A (2016) Expanding adolescent depression prevention through simple communication technologies. *The Journal of adolescent health: Official publication of the Society for Adolescent Medicine* 59: 373-374.
 65. Ranney ML, Freeman JR, Connell G, Spirito A, Boyer E, et al. (2016) A depression prevention intervention for adolescents in the emergency department. *The Journal of adolescent health: Official publication of the Society for Adolescent Medicine* 59: 401-410.
 66. Dobson KS, Dozois DJA (2008) Risk factors in depression. 2008.
 67. Hölzel L, Härter M, Reese C, Kriston L (2011) Risk factors for chronic depression- A systemic review. *J Affect Disord* 129: 1-13.
 68. Moussavi S, Chatterji S, Verdes E, Tandon A, Patel V, et al. (2007) Depression, chronic diseases, and decrements in health: Results from the World Health Surveys. *Lancet* 370: 851-858.
 69. Almeida OP, Draper B, Pirkis J, Snowdon J, Lautenschlager NT et al. (2012) Anxiety, depression and combined anxiety and depression: risk factors and outcome over two years. *Int Psychogeriatr* 24: 1622-1632.
 70. Heslin M, Desai R, Lappin JM, Donoghue K, Lomas B, et al. (2016) Biological and psychosocial risk factors for major depression. *Soc Psychiatry Psychiatr Epidemiol* 51: 233-245.
 71. Wiersma JE (2015) Childhood adversity and depression. *J Clin Psychiatry* 76: e906-e907.
 72. WHO Statistics (2013-2020) Mental Health Action Plan, World Health Organization, Geneva.
 73. Patten SB, Williams JVA, Lavorato DH, Bulloch AGM, Wiens K, et al. (2016) Why is major depression not changing? *J Affect Disord* 190: 93-97.
 74. Baxter AJ, Scott KM, Ferrari AJ, Rosana EN, Whiteford HA, et al. (2014) Challenging the myth of an "epidemic" of common mental disorders: trends in the global prevalence of anxiety and depression between 1990-2010. *Depression and anxiety* 31: 506-516.
 75. Werna-Seidler A, Perry Y, Calear AL, Newby JM, Christensen H (2017) School-based depression and anxiety preventive programs for young adults: A systemic review and meta-analysis. *Clin Psychol Rev* 51: 30-47.
 76. Stallard P, Buck R (2013) Preventing depression and promoting resilience: feasibility study of school-based cognitive-behavioral intervention. *Br J Psychiatry* 202: s18-s23.
 77. Takagaki K, Okamoto Y, Jinnin R, Mori A, Nishiyama Y, et al. (2016) Behavioral activation for late adolescents with subthreshold depression: a randomized controlled trial. *Eur Child Adolesc Psychiatry* 25: 1171-1182.
 78. Kiosses DN (2016) Innovative research on prevention of common mental disorders in older adults. *Am J Geriatr Psychiatry* 24: 417-420.
 79. Robinson J, Hetrick S, Cox G, Bendall S, Yuen HP, et al. (2016) Can an internet-based intervention reduce suicidal ideation, depression and hopelessness among secondary school students: Results from a pilot study. *Early Intervention in Psychiatry* 59: 28-35.
 80. Qi W, Gevonden M, Shalev A (2016) Prevention of post-traumatic stress disorder after trauma: Current evidence and future directions. *Curr Psychiatry Rep* 18: 20-30.
 81. Gildengers AG, Butters MA, Albert SM, Anderson SJ, Dew MA, et al. (2016) Design and implementation of an intervention development study: Retaining cognition while avoiding late-life depression (ReCall). *Am J Geriatr Psychiatry* 24: 444-454.
 82. Sakashita T, Oyama H (2016) Overview of community-based studies of depression screening intervention among the elderly population in Japan. *Aging Ment Health* 20: 231-239.
 83. Cockayne NL, Duffy SL, Bonomally R, English A, Amminger PG, et al. (2015) The beyond ageing project phase 2-a double-blind, selective prevention, randomized, placebo-controlled trial of omega-3 fatty acids and sertraline in an older age cohort at risk of depression: study protocol for a randomized controlled trial. *Trials* 16: 237.
 84. Rohde P, Stice E, Gau JM (2013) Effects of three depression prevention interventions on risk for depression disorder onset in the context of depression risk factors. *Prev Sci* 13: 584-593.
 85. Bazzatello P, Brignolo E, DeGrandi E, Bellino S (2016) Supplementation with Omega-3 fatty acids in psychiatric disorders: A review of literature data. *J Clin Med* 5: 67.
 86. Velásquez AM, López MA, Quiñonez N, Paba DP (2015) Yoga for the prevention of depression, anxiety, and aggression and the promotion of socio-emotional competencies in school-aged children. *Educational Research and Evaluation* 21: 407-15.
 87. Daley AJ, Foster L, Long G, Palmer C, Robinson O, et al. (2015) The effectiveness of exercise for the prevention and treatment of antenatal depression: systematic review with meta-analysis. *BJOG* 122: 57-63.

Citation: Al Qahtani A, Al Qahtani N (2017) Prevention of Depression: A Review of Literature. *J Depress Anxiety* 6: 292. doi:[10.4172/2167-1044.1000292](https://doi.org/10.4172/2167-1044.1000292)