

Primary Non-Hodgkin's Lymphoma of The Nasal Cavity: Unusual Site and Unusual Presentation

Vishwas D Pai^{1*}, Vidhya Manohar², Kiran Kattimani³ and Sheshachala Lingasur³

¹Department of Surgical Oncology, Kerudi Cancer Hospital, Bagalkot 587101, Karnataka, India

²Department of Pathology, SRL Diagnostics, Banagalore 560076, Karnataka, India

³Department of Medical Oncology, Kerudi Cancer Hospital, Bagalkot 587101, Karnataka, India

Abstract

Non-Hodgkin's lymphoma (NHL) is a heterogeneous subgroup of lymphomas with characteristic morphological and molecular features. Although lymph nodes are the most common organ of involvement, such as liver, stomach, intestine and bone marrow may also be affected on occasions. Primary non-Hodgkin's lymphoma (NHL) of the nasal cavity is an extra nodal form of NHL characterized by association with Epstein-Barr virus (EBV) infection, predominance of T-cell phenotype and a worse prognosis compared to nodal lymphoma. Although relatively common in China, nasal NHL is considered a rare entity in other countries of South East Asia. We are presenting a case of nasal NHL which was initially misdiagnosed as inverted papilloma and treated with surgical resection.

Keywords: Non-Hodgkin's Lymphoma; Paranasal lymphoma; Nasal lymphoma

Introduction

Lymphomas are malignant neoplasm of lympho-reticular system [1]. Non Hodgkin's lymphoma (NHL) is a heterogeneous subgroup of lymphomas with characteristic morphological and molecular features. Although lymph nodes are the most common organ of involvement, such as liver, stomach, intestine and bone marrow may also be affected on occasions [2]. Involvement of the nasal cavities by primary NHL is rare. These are classically associated with Epstein-Barr virus (EBV) infection and hence are common in those regions where EBV infection is endemic [3]. Although relatively common in China, nasal NHL is considered a rare entity in other countries of South East Asia [4,5]. There is no consensus on the management of these rare malignant neoplasm [6]. We are presenting a case of nasal NHL which was initially misdiagnosed as inverted papilloma and treated with surgical resection.

Case Report

A 50 year old gentleman presented with history of epistaxis and nasal blockade of 4 month duration. On examination, he had a polypoidal mass completely filling the left nasal cavity. Contrast enhanced computed tomography (CECT) of the nasal cavity revealed a well-defined, heterogeneously enhancing soft tissue mass occupying the entire nasal cavity with extension into the left maxillary sinus causing mass effect with bulging of nasal septum to the opposite side which was suggestive of inverted papilloma (Figure 1a). Biopsy of the mass was not conclusive of any specific malignancy even after 2 attempts. In view of CECT findings and clinical presentation of the patient, tentative clinical diagnosis was inverted papilloma and hence surgery was planned. Medial maxillectomy was performed. Intra and post-operative course were uneventful and patient was discharged on 3rd post operative day. Histopathological examination revealed monomorphic, large cells with eccentric nucleus and moderate to abundant cytoplasm with plasmablastic differentiation which was suggestive of Non Hodgkin's lymphoma of plasmablastic differentiation (Figure 1b and 1c). CECT thorax and abdomen were done as a part of staging. There was no evidence of disease at any other site and hence patient was staged as IE limited disease according to Ann Arbor staging system. He was treated with CHOP chemotherapy comprising of Cyclophosphamide 750 mg/m², Doxorubicin 50 mg/m², Vincristine 1.4 mg/m², Prednisolone 100 mg/m² for 5 days for a total of 6 cycles. At last follow up, patient was alive and disease free.

Discussion

Primary non-Hodgkin's lymphoma (NHL) of the nasal cavity is an extra nodal form of NHL characterized by association with Epstein-Barr virus (EBV) infection, predominance of T-cell phenotype and a worse prognosis compared to nodal lymphoma. In the west, paranasal sinus NHL is more common. In contrast, in the east, nasal NHL is more common than the NHL of the paranasal sinus [7].

Xiong et al. reported the largest series of primary non-Hodgkin's lymphoma (NHL) of the nasal cavity [8]. Median age of presentation was 44 years (Range: 11-79 years) and males were affected more often than females. Maxillary sinus was the most frequent site of involvement followed by the ethmoid sinus and the nasopharynx. The most common presenting symptoms include nasal obstruction, epistaxis and headache. Diplopia and blurred vision are rare infrequent symptoms associated with nasal NHL. The present patient is a middle aged male who presented with nasal obstruction and extension of the tumour into the maxillary sinus.

Macroscopically, these tumours are sub mucosal in contrast to squamous cell carcinomas which are ulcerative [9]. Angio invasion and necrosis are the two characteristic histological features of sino-nasal NHL [10]. NHL of the nasal cavity are predominantly T-cell lymphomas, whereas NHL of the paranasal sinuses are B-cell lymphomas.

Staging of the patients is according to Ann Arbor system [11]. In addition to CECT of the para nasal sinuses, CECT of the abdomen and thorax are required for adequate staging. Management is according to the stage of the disease. Stage IE is treated with chemotherapy or radiotherapy alone whereas stage IIE/IIIE and IVE require varying

***Corresponding author:** Dr. Vishwas D Pai, Department of Surgical Oncology, Kerudi Cancer Hospital, Bagalkot 587101, Karnataka, India, Tel:+91-9449333502; E-mail: vishpai88@gmail.com

Received: August 18, 2016 ; **Accepted:** October 25, 2016; **Published:** November 01, 2016

Citation: Pai VD, Manohar V, Kattimani K, Lingasur S (2016) Primary Non-Hodgkin's Lymphoma of The Nasal Cavity: Unusual Site and Unusual Presentation. Reconstructive Surgery and Anaplastology 5: 161. doi: [10.4172/2161-1173.1000161](https://doi.org/10.4172/2161-1173.1000161)

Copyright: © 2016 Pai VD, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1A: Contrast enhanced computed tomography (CECT) of the nasal cavity.




Figure 1B: Histopathological examination revealed monomorphic, large cells with eccentric nucleus.

combination of chemotherapy and radiotherapy [12-15]. The present patient had limited stage IE disease and hence was treated with chemotherapy alone.

In general, patients with nasal NHL have worse prognosis compared to other extra nodal sites of NHL. Stage is the most important factor determining prognosis. Patients with Stage IE disease confined to the nasal cavity have improved survival with estimated five-year overall survival (OS) and disease free survival (DFS) of 90% and 89%, respectively. In contrast, patients with Stage IIE, IIIE, and IVE have a worse prognosis with uniformly fatal outcome irrespective of the use of chemotherapy.

Conclusion

The purpose of presenting this case is to highlight the rarity of nasal




Figure 1C: Histopathological examination of cytoplasm with plasmablastic differentiation.

NHL in Indian subcontinent and stress the need for better awareness among treating oncologists, pathologist and radiologists about this disease entity.

References

1. Shohat I, Berkowicz M, Dori S, Horowitz Z, Wolf M, et al. (2004) Primary non-Hodgkin's lymphoma of the sinonasal tract. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 97: 328-331.
2. Wang J, Sun NC, Weinstein SM, Canalis R (2000) Primary T-cell-rich B-cell lymphoma of the ethmoid sinus. A case report with 5 years of follow-up. *Arch Pathol Lab Med* 124: 1213-1216.
3. Mishima K, Horiuchi K, Kojya S, Takahashi H, Ohsawa M, et al. (1994) Epstein-Barr virus in patients with polymorphic reticulosis (lethal midline granuloma) from China and Japan. *Cancer* 73: 3041-3046.
4. Liang R, Todd D, Chan TK, Chiu E, Lie A, et al. (1995) Treatment outcome and prognostic factors for primary nasal lymphoma. *J Clin Oncol* 13: 666-670.
5. Abbondanzo SL, Wenig BM (1995) Non-Hodgkin's lymphoma of the sinonasal tract. A clinicopathologic and immunophenotypic study of 120 cases. *Cancer* 75: 1281-1291.
6. Itami J, Itami M, Mikata A, Tamaru J, Kaneko T, et al. (1991) Non-Hodgkin's lymphoma confined to the nasal cavity: its relationship to the polymorphic reticulosis and results of radiation therapy. *Int J Radiat Oncol Biol Phys* 20: 797-802.
7. Logsdon MD, Ha CS, Kavadi VS, Cabanillas F, Hess MA, et al. (1997) Lymphoma of the nasal cavity and paranasal sinuses: improved outcome and altered prognostic factors with combined modality therapy. *Cancer* 80: 477-488.
8. Li YX, Coucke PA, Li JY, Gu DZ, Liu XF, et al. (1998) Primary non-Hodgkin's lymphoma of the nasal cavity: prognostic significance of paranasal extension and the role of radiotherapy and chemotherapy. *Cancer* 83: 449-456.
9. Chalastros T, Elefteriadou A, Giotakis J, Soulandikas K, Korres S, et al. (2007) Non-Hodgkin's lymphoma of nasal cavity and paranasal sinuses. A clinicopathological and immunohistochemical study. *Acta Otorhinolaryngol Ital* 27: 6-9.
10. Lipford EH, Margolick JB, Longo DL, Fauci AS, Jaffe ES (1988) Angiocentric immunoproliferative lesions: a clinicopathologic spectrum of post-thymic T-cell proliferations. *Blood* 72: 1674-1681.
11. Carbone PP, Kaplan HS, Musshoff K, Smithers DW, Tubiana M (1971) Report of the Committee on Hodgkin's Disease Staging Classification. *Cancer Res* 31:1860-1861.
12. Al-Hakeem DA, Fedele S, Carlos R, Porter S (2007) Extranodal NK/T-cell lymphoma, nasal type. *Oral Oncol* 43: 4-14.
13. Kwong YL (2005) Natural killer-cell malignancies: diagnosis and treatment. *Leukemia* 19: 2186-2194.

14. Hatta C, Ogasawara H, Okita J, Kubota A, Ishida M, et al. (2001) Non-Hodgkin's malignant lymphoma of the sinonasal tract--treatment outcome for 53 patients according to REAL classification. *Int J ORL and HNS* 28: 55-60.
15. You JY, Chi KH, Yang MH, Chen CC, Ho CH, et al. (2004) Radiation therapy versus chemotherapy as initial treatment for localized nasal natural killer (NK)/T-cell lymphoma: a single institute survey in Taiwan. *Ann Oncol* 15: 618-625.

Citation: Pai VD, Manohar V, Kattimani K, Lingasur S (2016) Primary Non-Hodgkin's Lymphoma of The Nasal Cavity: Unusual Site and Unusual Presentation. *Reconstructive Surgery and Anaplastology* 5: 161. doi: [10.4172/2161-1173.1000161](https://doi.org/10.4172/2161-1173.1000161)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>