

Prognosis of Orthostatic Hypotension

Wilbert S Aronow*

Cardiology Division, Department of Medicine, New York Medical College, Valhalla, New York, USA

*Corresponding author: Wilbert S Aronow, MD, FACC, FAHA, Cardiology Division, New York Medical College, Macy Pavilion, Room 138, Valhalla, NY 10595, USA, Tel: (914) 493-5311; Fax: (914) 235-6274; E-mail: wsaronow@aol.com

Received date: September 25, 2016; Accepted date: September 26, 2016; Published date: September 27, 2016

Copyright: © 2016 Aronow WS. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Editorial

Orthostatic hypotension is defined as a fall of 20 mm or more in systolic blood pressure or of 10 mm or more in diastolic blood pressure within 3 minutes of standing [1]. All persons receiving antihypertensive drugs routinely should have their blood pressure measured in the sitting position and within 3 minutes of standing [2]. Blood pressure should not be taken immediately after eating as postprandial hypotension may occur at that time [3]. Orthostatic hypotension and postprandial hypotension may coexist, and both disorders are associated with adverse clinical outcomes [3].

Orthostatic hypotension may be associated with advanced age, disorders associated with hypovolemia, hypertension, diabetes, neurological disorders, antihypertensive medications, antidepressants, antipsychotic drugs, anti-parkinsonian medications, alcohol, cardiovascular conditions, endocrine conditions, and other disorders [4]. The prevalence of orthostatic hypotension in older persons, mean age 82 years, in a long-term health care facility was 13% in 257 persons receiving cardiovascular or psychotropic drugs and 3% in those who did not receive cardiovascular or psychotropic drugs [1]. The prevalence of orthostatic hypotension in 4,733 diabetics, mean age 62.1 years, in the Action to Control Cardiovascular Risk in Diabetes (ACCORD) blood pressure trial was 17.8% at baseline, 10.4% at 1 year, 12.8% at 4 years, and 20.0% at 1 or more visits [5]. The prevalence of orthostatic hypotension in 2,636 persons aged 75 years and older, mean age 79.9-years, in the Systolic Blood Pressure Intervention Trial (SPRINT) was 21.0% in persons randomized to a systolic blood pressure below 120 mmHg versus 21.8% in persons randomized to a systolic blood pressure below 140 mmHg [6]. The prevalence of orthostatic hypotension with dizziness was 1.9% in persons randomized to a systolic blood pressure below 120 mmHg versus 1.3% in persons randomized to a systolic blood pressure below 140 mmHg [6].

At 4-year follow-up of 3,522 older Japanese-American men in the Honolulu Heart program, orthostatic hypotension was significantly associated with increased all-cause mortality by 1.64 times [7]. At 4.4-year follow-up of 2,786 community dwelling older Italians, orthostatic hypotension was significantly associated with increased all-cause mortality by 1.23 times, with cardiovascular mortality by 1.41 times, and by non-cardiovascular mortality by 1.19 times [8]. At 22.7-year follow-up of 33,346 persons, mean age 45.7-years, in the Swedish Malmo Preventive Project, orthostatic hypotension was significantly associated with all-cause mortality by 1.21 times, with coronary events by 1.17 times, with stroke by 1.17 times, and by a composite endpoint of death, coronary event, or stroke by 1.18 times [9].

A meta-analysis included 13 prospective studies with 121,913 persons [10]. At 5-year follow-up of 65, 174 persons, orthostatic hypotension significantly increased all-cause mortality by 1.5 times. At

6.4-year follow-up of 49, 512 persons, orthostatic hypotension significantly increased coronary heart disease by 1.41 times. At 6.8 to 24-year follow-up of 50, 096 persons, orthostatic hypotension significantly increased heart failure by 2.25 times. At 6.8-year follow-up of 58, 300 persons, orthostatic hypotension significantly increased stroke by 1.64 times [10].

At 46.9-month follow-up in the ACCORD blood pressure trial, orthostatic hypotension was significantly associated with all-cause mortality by 1.61 times and with heart failure death or heart failure hospitalization by 1.85 times [5].

At 6-year follow-up of 12,433 black and white middle-aged men and women in the Atherosclerosis Risk in Communities study, orthostatic hypotension was significantly associated with coronary heart disease by 1.85 times [11]. At 7.9-year follow-up of 11,707 persons free of stroke and clinical heart disease at baseline in the Atherosclerosis Risk in Communities study, orthostatic hypotension was significantly associated with ischemic stroke by 2.0 times [12]. At 17.5-year follow-up of 12,363 persons free of heart failure at baseline in the Atherosclerosis Risk in Communities study, orthostatic hypotension was significantly associated with heart failure by 1.54 times [13]. At 13-year follow-up of 5,273 persons free of heart failure at baseline in the Cardiovascular Health Study, we reported that orthostatic hypotension was significantly associated with incident heart failure by 1.24 times [14]. Orthostatic hypotension may also cause syncope [4].

References

1. Aronow WS, Lee NH, Sales FF, Etienne F (1988) Prevalence of postural hypotension in elderly patients in a long-term health care facility. *Am J Cardiol* 62: 336.
2. Aronow WS, Fleg JL, Pepine CJ, Artinian NT, Bakris G, et al. (2011) ACCF/AHA 2011 expert consensus document on hypertension in the elderly: a report of the American College of Cardiology Foundation Task Force on Clinical Expert Consensus Documents. Developed in collaboration with the American Academy of Neurology, American Geriatrics Society, American Society for Preventive Cardiology, American Society of Hypertension, American Society of Nephrology, Association of Black Cardiologists, and European Society of Hypertension. *J Am Coll Cardiol* 57: 2037-2114.
3. Aronow WS, Ahn C (1997) Association of postprandial hypotension with incidence of falls, syncope, coronary events, stroke, and total mortality at 29-month follow-up in 499 older nursing home residents. *J Am Geriatr Soc* 45: 1051-1053.
4. Aronow WS (1999) Dizziness and syncope. In Hazzard WR, Blass JP, Ettinger WH, Jr, Halter JB, Ouslander JG (eds). *Principles of Geriatric Medicine and Gerontology* (fourth edition), New York city, McGraw-Hill Inc: 1519-1534.
5. Fleg JL, Evans GW, Margolis KL, Barzilay J, Basile JN, et al. (2016) Orthostatic Hypotension in the ACCORD (Action to Control

- Cardiovascular Risk in Diabetes) Blood Pressure Trial: Prevalence, Incidence, and Prognostic Significance. *Hypertension* 68: 888-895.
6. Williamson JD, Supiano MA, Applegate WB, Berlowitz DR, Campbell RC, et al. (2016) Intensive vs standard blood pressure control and cardiovascular disease outcomes in adults aged 75 years. A randomized clinical trial. *JAMA* 315: 2673-2682.
 7. Masaki KH, Schatz IJ, Burchfiel CM, Sharp DS, Chiu D, et al. (1998) Orthostatic hypotension predicts mortality in elderly men: the Honolulu Heart Program. *Circulation* 98: 2290-2295.
 8. Veronese N, De Rui M, Bolzetta F, Zambon S, Corti MC, et al. (2015) Orthostatic Changes in Blood Pressure and Mortality in the Elderly: The Pro.V.A Study. *Am J Hypertens* 28: 1248-1256.
 9. Fedorowski A, Stavenow L, Hedblad B, Berglund G, Nilsson PM, et al. (2010) Orthostatic hypotension predicts all-cause mortality and coronary events in middle-aged individuals (The Malmo Preventive Project). *Eur Heart J* 31: 85-91.
 10. Ricci F, Fedorowski A, Radico F, Romanello M, Tataschiere A, et al. (2015) Cardiovascular morbidity and mortality related to orthostatic hypotension: a meta-analysis of prospective observational studies. *Eur Heart J* 36: 1609-1617.
 11. Rose KM, Tyroler HA, Nardo CJ, Arnett DK, Light KC, et al. (2000) Orthostatic hypotension and the incidence of coronary heart disease: the Atherosclerosis Risk in Communities study. *Am J Hypertens* 13: 571-578.
 12. Eigenbrodt ML, Rose KM, Couper DJ, Arnett DK, Smith R, et al. (2000) Orthostatic hypotension as a risk factor for stroke: the atherosclerosis risk in communities (ARIC) study, 1987-1996. *Stroke* 31: 2307-2313.
 13. Jones CD, Loehr L, Franceschini N, Rosamond WD, Chang PP, et al. (2012) Orthostatic hypotension as a risk factor for incident heart failure: the atherosclerosis risk in communities study. *Hypertension* 59: 913-918.
 14. Alagiakrishnan K, Patel K, Desai RV, Ahmed MB, Fonarow GC, et al. (2014) Orthostatic hypotension and incident heart failure in community-dwelling older adults. *J Gerontol A Biol Sci Med Sci* 69: 223-230.