

Radioactivity: An Introduction to Mysterious Science

Nida Tabassum Khan*

Department of Biotechnology, Faculty of Life Sciences and Informatics, Balochistan University of Information Technology Engineering and Management Sciences (BUIITEMS), Quetta, Pakistan

Abstract

Materials that undergo decay process to attain a stable configuration followed by the emission of particles such as alpha, beta and gamma are termed as radioactive elements. Radioactivity is measured in Curie (Ci) and Becquerel (Bq) SI units. Though radioactive elements are abundant on Earth and constantly decays to produce stable nuclei but such nuclear decay process may pose serious health threats to living beings. Therefore, must be regulated when working *in vitro*. Biosafety levels should be strictly observed when engaged in radioactive experiments to minimize contamination exposure.

Keywords: Radioactive decay; Radioactive poisoning; Alpha particles; Beta particles; Half-life

Introduction

In 1896 Henri Becquerel discovered phenomena of radioactivity by exposing potassium uranyl sulfate to sunlight, considering that it will emit neutral X-rays but in actual the emitted radiation was electrically charged since it bent in the magnetic field. So, the experiment was conducted using different elements that emits radiation that showed deflection in different directions in the magnetic field or not at all. Thus three different types of radioactive radiations i.e., negative, positive, and neutral were identified [1]. Marie Curie coined the term radioactivity and along with her husband Pierre, discovered other radioactive elements such as radium and polonium from radioactive ore of uranium for which she won a second Nobel Prize in 1911. Becquerel and the Curies received the Nobel Prize in 1903. The next important step down this road of discovery came from Ernest Rutherford who characterized the defining properties of these radioactive particles and named these as alpha, beta, and gamma particles. Classification of these radiations was based on their ability to penetrate matter. Alpha particles being massive moves slowly as compared to beta and gamma particles [2-4].

Radioactivity in nature

Elements with atomic number (Z) higher than Bi is radioactive in nature and are abundant in Earth. The earth also contains several elemental radioisotopes such as Potassium-40 which undergoes beta decays to form stable ^{40}Ar and ^{40}Ca . Uranium to lead raiton revealed that Earth is approximately 4.5 billion years old. Rn 222, an intermediate product of ^{238}U decay with 3.8 days half-life contributes in background radiation. Radioactive isotopes such as ^7Be and ^{14}C are constantly produced in the upper atmosphere. In order to gain understanding of Earth structural construct, study of radioactivity is crucial [5,6].

Radioactivity units

Becquerel (Bq) after Henri Becquerel is the unit in which radioactivity is measured and which the number of decays per second is.

One day per second equals one Becquerel

Curie another unit, named after Pierre and Marie Curie.

One curie is the activity of 1 gram of radium and equals 3.7×10^{10} Becquerel.

SI units and prefixes

Conference on Weights and Measures recommended the universal use of International System of Units for measuring radioactivity [7-9] (Tables 1 and 2).

Radiation measurements

Radiation measurements: Different related units are used for measuring radioactivity, absorbed dose, exposure and dose equivalent (r-e-a-d).

- Curie (Ci) and Becquerel (Bq) are the units for the measurements of radioactivity which refers to the concentration of emitted particles by a sample. It denotes the number of atoms in the given sample that decayed in a given time period.
- Coulomb/kilogram (C/kg) and Roentgen (R) are the units for exposure which is defined as the amount of radiation traveling through the air.
- Gray (Gy) and radiation absorbed dose (rad) measures the concentration of absorbed (deposited) radiation by a material or person.
- Sievert (Sv) and Roentgen equivalent man (rem) are units to measures dose equivalent also known as effective dose which is defined as the amount of absorbed radiation and its medical effects. Dose equivalent is same as the absorbed dose for beta and gamma radiation but larger than the absorbed dose for alpha and neutron radiation that are more damaging [10].

Types of radiations: Ionizing radiation emitted during radioactive decay are categorized as alpha, beta, gamma, protons, and neutrons.

Radioactive decay: Around 50 naturally occurring radioisotopes


*Corresponding author: Nida Tabassum Khan, Department of Biotechnology, Faculty of Life Sciences and Informatics, Balochistan University of Information Technology Engineering and Management Sciences (BUIITEMS), Quetta, Pakistan, Tel: 03368164903; E-mail: nidatabassumkhan@yahoo.com

Received August 08, 2017; Accepted September 03, 2017; Published September 08, 2017

Citation: Khan NT (2017) Radioactivity: An Introduction to Mysterious Science. J Phys Chem Biophys 7: 254. doi: 10.4172/2161-0398.1000254


Copyright: © 2017 Khan NT. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

are found on earth. Impulsive radioactive fragmentation of an atomic nucleus to achieve a stable form by the emission of radiations is termed as radioactive decay [11]. Based on different types of radiations released during decay process there are three types of radioactive decays which are as follows:



Alpha decay

Alpha particle (4He nucleus) is emitted during the decay process. Such nuclei have high proton to neutron ratio. Alpha particles are quiet stable in their configuration with two protons and two neutrons. Protons to neutrons ratio is reduced in the parent nucleus after decay by altering the Z. so the parent and the daughter atoms are chemically different elements [12].




Beta decay

Beta particles are negatively charged electrons or positively charges electrons (positron) that is emitted from nuclei with too many protons during decay. There are two types of beta decays.

1) Beta minus decay: In which a neutron decays into a proton, an electron, and an antineutrino

2) Beta plus decay: In which a proton decays into a neutron, a positron, and a neutrino [13].

Gamma decay



Through the emission of photons in gamma decay, the atomic mass of the parent and daughter nucleus remains the same. Thus parent and daughter atoms are chemically similar elements [14].

Half life: Half-life is defined as the time taken for one-half of the nuclei in a sample to decay. It is denoted by $t_{1/2}$. Radioactive decay is as an exponential process. N denotes the number of original nuclei remaining after a time t from an original sample of N_0 nuclei.

$$N = N_0 e^{-(t/T)}$$

where T is the mean lifetime of the parent nuclei. Thus $t_{1/2} = 0.693T$ [15,16]. Table 3 provides the half life of few radioactive elements below.

Properties of radioactive radiations

Alpha particles: Alpha particle contains two neutrons and two protons

- Particle carry positive charge
- Mass of each alpha-particle is 4 times that of a proton or H-atom.
- Ionization power of alpha rays is very high.
- Penetration power of is very small
- Produce fluorescence in different substances
- Produce burn and source on human body
- Produce artificial radioactivity is certain nuclei
- They have strong ionizing power because they remove electrons from the atoms of gas through which they pass
- Their velocity range is 3×10^7 m/s to 3×10^6 m/s.

- Highly energetic emits 5 MeV.
- Travel only a few centimeters in air and is easily stopped by a paper sheer or the outer skin layer. Radon, uranium, radium, and thorium are alpha emitters.

Beta particles

- Negatively charged particles i.e., Electron
- Velocity is from 9×10^7 m/sec to 27×10^7 m/sec
- Affect the photo graphic plate
- Ionization power is very small
- Kinetic energy is less than that of alpha - rays.
- Produce fluorescence in different substance
- Produce secondary radiations called bremsstrahlung
- Travel several meters in air and millimeters into the human body and is stopped by an aluminum or plastic sheet. Sulfur-35, hydrogen-3 phosphorus-33, phosphorus-32 and carbon-14 are beta emitters.

Gamma particles

- Electrically neutral
- Travel with the velocity of light that is 3×10^8 m/sec
- Penetration power is very large. It is about hundred times larger than that of alpha rays.
- Produce feeble fluorescence
- Travel at the speed of light
- Could only be stopped by a thick sheet of lead, steel, concrete or several meters of water. Cobalt-60, cesium-137, zinc-65, and radium-22 are gamma emitters [19-22].

Radioactive poisoning

Numerous incidents in the past revealed that if biosafety levels are not strictly observed when working with radioactive isotopes, it increases the possibility of contamination with life threatening consequences. For example in 2006 murder of former Russian intelligence and death of Marie Curie's daughter Irene by radioactive polonium. Similarly in 2003 a Russia journalist, Yuri Shchekochikhin and in 2004 a St Petersburg businessman Romam Tsepov died of radioactive poisoning [23]. Therefore it is important to keep in mind the safety considerations and measures while working in lab to ensure a radioactive free environment is established for protection of an individual from radioactive poisoning.

Safety considerations

Following safety levels should be observed while working with a radioactive material:

Defensive clothing: When working with an open radioactive source wear long length gloves and lab coats, closed toe shoes, safety glasses/goggles, radiation dosimeter, radiation monitor badges etc., to cover maximum parts of your body to avoid radioactive exposure because radioactive incidents frequently involves spills or splashes which can readily contaminate bare parts of the body.

Drinking/Eating: Drinking/eating, smoking etc., should be avoided where open radioactive sources present or stored to reduce the possibility of radioactive content intake in the body.

Oral pipetting: Mouth pipetting of radioactive solutions must not be avoided.

Security: Radioactive resources should be placed in shielded storage containers with proper labelling at secured places with maximum supervision and should not be left in an unattended lab or room.

Radioactive warning labels: Proper signs and symbols should be used to indicate the presence or storage of radioactive materials to avoid mishandling and exposure [24,25]. Labeling of rooms and containers of radioisotopes is mandatory under the following conditions [26,27] (Table 4).

Hoods and biosafety cabinets: Volatile radioelements i-125 or S-35 methionine/cysteine, must be handled inside specifically designed RAM hoods or safety cabinets equipped with alarming flow monitoring device to eliminate the risk of contamination.

ALARA: Radiation exposure must be kept as low as reasonably achievable means all safety measures and monitoring should be attempted to keep radiation doses ALARA [28].

Lab working concerns

- Be aware with the relative properties of the radioactive isotope that is to be used with any specific precautionary concerns specific to that radioisotope.
- Unexperienced nuclear procedures should be rehearsed before engaging with the actual radioactive substance following standard biosafety levels such as protective clothing, use of contamination free safety hoods, etc.

	Radioactivity	Absorbed Dose	Dose Equivalent	Exposure
Common Units	curie (Ci)	Rad	Rem	roentgen (R)
SI Units	becquerel (Bq)	gray (Gy)	sievert (Sv)	coulomb/kilogram (C/kg)

Table 1: SI Units for measuring radioactivity.

To convert from	To	Multiply by
Curies (Ci)	Becquerels (Bq)	3.7×10^{10}
millicuries (mCi)	megabecquerels (MBq)	37
microcuries (μ Ci)	megabecquerels (MBq)	0.037
millirads (mrad)	milligrays (mGy)	0.01
millirems (mrem)	microsieverts (μ Sv)	10
milliroentgens (mR)	microcoulombs/kilogram (μ C/kg)	0.258
Becquerels (Bq)	Curies (Ci)	2.7×10^{-11}
megabecquerels (MBq)	millicuries (mCi)	0.027
megabecquerels (MBq)	microcuries (μ Ci)	27
milligrays (mGy)	millirads (mrad)	100
microsieverts (μ Sv)	millirems (mrem)	0.1
microcoulombs/kilogram (μ C/kg)	milliroentgens (mR)	3.88

Table 2: Conversion Equivalence.

Radioisotope	Half-Life
Hydrogen-3	12.3 years
Carbon-14	5730 years
Phosphorus-32	14.3 days
Phosphorus-33	25.3 days
Sulfur-35	87.6 days
Iodine-125	60.1 days

Table 3: Half-life of radioisotopes [17,18].

Radioactive isotopes	Activity (μ Ci)
H-3	1000
C-14	100
P-32	10
P-33	100
S-35	100
Ca-45	100
Cr-51	1000
Fe-59	10
Zn-65	10
I-125	1
U-238	100

Table 4: Radioisotope activity labelling.

- Avoid direct handling of radioactive entities instead use remote handling tools and instruments.
- Proper storage of radioactive waste in specifically designed labelled storage cans with substantial Plexiglas shielding from external radiation levels.
- Authorized access to individuals to radioactive storage facilities with maximum security and supervision to avoid unauthorized access or breaching.
- Biosafety rules and regulations for working with radioisotopes should be made visible for everyone before the start of an experiment.
- Proper storage/disposal of radioactive waste in shielded cans and containers.
- Contamination monitoring should be done on regular basis after every nuclear experiment conducted.
- Washing sink for contaminated glassware and equipment should be clearly labelled with radiation warning signs [29,30].

Conclusion

Thus, radioisotopes are an abundant source of radioactive emissions that can be used in different biomedical applications but needs strict biosafety regulation to avoid exposure.

References

1. Minder W (1981) History of radioactivity.
2. Gasparini P (1984) History of radioactivity. In: Uranium geochemistry, mineralogy, geology, exploration and resources. Springer, Netherlands, pp: 1-3.
3. Friedlander G (1981) Nuclear and radiochemistry. John Wiley and Sons.
4. Annunziata MF (2007) Radioactivity: Introduction and History.
5. Lieser KH (2013) Radioactivity in Nature. Nuclear and Radiochemistry: Fundament App, pp: 1-4.
6. White JR, Cameron AE (1948) The natural abundance of isotopes of stable elements. Phy Rev 74: 991.
7. Annunziata MF (2012) Handbook of Radioactivity Analysis. Academic Press.
8. Curtiss LF, Evans RD, Johnson W, Seaborg GT (1950) Units of radioactivity.
9. Friedlander G, Kennedy JW (1949) Introduction to Radiochemistry. John Wiley and Sons Inc., New York, USA.
10. US Patent (1953) System for measuring Radioactivity. 2: 643,344.
11. Malik SS, Gupta RK (1989) Theory of cluster radioactive decay and of cluster formation in nuclei. Phys Rev C 39: 5.
12. Westmeier W, Merklin A (1985) Catalogue of alpha particles from radioactive decay.

13. Huck A, Klotz G, Knipper A, Miehé C, Richard-Serre C, et al. (1985) Beta decay of the new isotopes K 52, Ca 52, and Sc 52; A test of the shell model far from stability. *Phy Rev C* 31: 2226.
14. Erdtmann G, Soyka W (1979) The gamma rays of the radionuclides.
15. Browne E, Dairiki JM, Doebler RE (1978) Table of isotopes. National Standard Reference Data System.
16. Houtermans H, Milosevic O, Reichel F (1980) Half-lives of 35 radionuclides. *The Intern J App Rad Isot* 31: 153-154.
17. Woods MJ, Collins SM, Woods SA (2004) Evaluation of half-life data. National Physical Laboratory.
18. Reynolds SA, Emery JF, Wyatt EI (1968) Half-lives of radionuclides-III. *Nucl Sci Eng* 32: 46-48.
19. Rutherford E, Chadwick J, Ellis CD (2010) Radiations from radioactive substances. Cambridge University Press.
20. Knoll GF (2010) Radiation detection and measurement. John Wiley and Sons.
21. Bahr B, Lemmer B, Piccolo R (2016) Alpha, Beta and Gamma Rays. In: *Quirky Quarks*, Springer Berlin Heidelberg, pp: 170-173.
22. Wilson D (1983) Rutherford-simple genius.
23. Perkins AC (2007) The London polonium poisoning: events and medical implications. *World J Nucl Med* 6: 102-106.
24. Faires RA, Parks BH (1973) Radioisotope laboratory techniques.
25. Hendee WR (1970) Medical radiation physics.
26. Janžekovič H, Križman M (1973) Radioisotopes Used in Analytical Chemical Instruments. In *Book of Abstracts*, p: 124.
27. Newbery GR (1960) The Distribution of Radioisotopes Public Safety Aspects. *Royal Soc Heal J* 80: 220-228.
28. Lenhart MJ (1986) US Patent No. 4,581,538. Washington, DC: U.S. Patent and Trademark Office.
29. Woodhead DS (1984) Contamination due to radioactive materials. In: *Marine ecology: A comprehensive, integrated treatise on life in oceans and coastal waters*.
30. Goldstein JA (2008) US Patent No. 7,391,042. Washington, DC: U.S. Patent and Trademark Office.