

Recent Possibilities for Diagnosis and Treatment of Post Parturient Uterine Diseases in Dairy Cow

Ottó Szenci*

MTA-SZIE Large Animal Clinical Research Group and Szent István University, Faculty of Veterinary Science, Department and Clinic for Production Animals, Üllő-Dóra major, H-2225 Hungary

Abstract

The successful genetic selection for higher milk production caused a dramatic decline in the reproductive performance of dairy cows all over the world during the last decades. Achievement of optimum herd reproductive performance (calving interval of 12 months or 13 months with the first calf born at 24 months of age) requires concentrated management activities especially during the first 100 days following calving. The following management activities are needed to pursue during the early postpartum period to reach or approach the optimal reproductive performance such as careful surveillance and assistance at calving, prevention of post parturient metabolic diseases, early diagnosis and treatment of post parturient uterine diseases, accurate detection of oestrus, correct timing of insemination, reducing the effect of heat stress and early pregnancy diagnosis. Among these main activities only early diagnosis and treatment of post parturient uterine diseases and their effects on reproductive performance are discussed. Clinical metritis and clinical endometritis (or recently used purulent vaginal discharge: PVD) can be accurately diagnosed in the field, however it is very important to remark that the presence of PVD alone is not sufficient to diagnose clinical endometritis because cervicitis and/or vaginitis can also cause PVD. Both diseases have detrimental effect on reproduction therefore it is important to diagnose as soon as possible and treat accordingly.

Different treatment protocols that have changed in the recent years for those diseases, as well as recently performed therapies like ozone therapy or vaccination are also discussed. Although subclinical endometritis has also a detrimental effect on reproduction, it is complicated to diagnose in the field.

Keywords: Dairy cow; Clinical metritis; Clinical endometritis; Subclinical endometritis

The successful genetic selection for higher milk production in Holstein cows has nearly doubled the average milk production in the United States since 1960, to over 11.000 kg/year. Over the same time period, there has been a dramatic decline in the reproductive performance of dairy cows. The average number of days open (interval from calving to conception) and the number of services per conception have increased substantially. In order to decrease the longer lactations and the number of cows culled for reproductive reasons it is very important to improve our reproductive management practices [1]. Achievement of optimum herd reproductive performance (calving interval of 12 months or 13 months with the first calf born at 24 months of age) requires concentrated management activities especially during the first 100 days following calving. Early postpartum breeding of dairy cows results in more calves, and higher milk production per lactation [2]. Poor reproductive performance can reduce the number of calves born and milk production and may increase the cost of therapy and semen.

The following management activities such as careful surveillance and assistance at calving, prevention of post parturient metabolic diseases, early diagnosis and treatment of post parturient uterine diseases, accurate detection of oestrus, correct timing of insemination, reducing the effect of summer heat stress and early pregnancy diagnosis are needed to pursue during the early postpartum period to reach or approach the optimal calving interval [3].

Among these main activities only early diagnosis and treatment of post parturient uterine diseases and their effects on milk production and reproductive performance are discussed in the present work. However, this topic has also a great importance because it is generally accepted that up to 40% of dairy cows may have clinical metritis within the first two weeks after calving and infection may persist in 10 to 15% of animals more than 3 weeks after calving causing clinical or subclinical endometritis [4].

Clinical Metritis

Clinical metritis is an acute systemic illness due to infection of the uterus with bacteria, usually within 10 (21) days after parturition.

According to Sheldon et al. [5], clinical metritis can be categorized into three grades:

Grade 1 clinical metritis (CM1) can be characterized by an abnormally enlarged uterus and a purulent uterine discharge detectable in the vagina, within 21 days after calving.

Grade 2 clinical metritis (CM2) or puerperal metritis can be characterized by a fetid red-brown watery uterine discharge, atonic enlarged uterus and, usually pyrexia ($>39.5^{\circ}\text{C}$) [6,7]; in severe cases, reduced milk yield, dullness, inappetence or anorexia, elevated heart rate, and apparent dehydration may also be present. In some cases pyrexia even with daily monitoring of rectal temperature could not be detected [4,8] however an enlarged uterus with a thin wall and atonia used to be present with a fetid discharge. Puerperal metritis is often associated with retained placenta, dystocia, stillbirth or twins, and usually occurs toward the end of the first week after calving, being rare after the second week after calving [7,9,10]. It is important to emphasize

*Corresponding author: Ottó Szenci, MTA-SZIE Large Animal Clinical Research Group and Szent István University, Faculty of Veterinary Science, Department and Clinic for Production Animals, Üllő-Dóra major, H-2225 Hungary, Tel: (36) 29-521301; Fax: (36) 29-521385; E-mail: szenci.otto@aotk.szie.hu

Received December 07, 2015; Accepted February 03, 2016; Published February 10, 2016

Citation: Szenci O (2016) Recent Possibilities for Diagnosis and Treatment of Post Parturient Uterine Diseases in Dairy Cow. JFIV Reprod Med Genet 4: 170. doi:10.4172/2375-4508.1000170

Copyright: © 2016 Szenci O. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

that puerperal metritis (CM2) may occur in around 10%-15% of cows with spontaneous calving and without retained foetal membranes [8].

Grade 3 clinical metritis (CM3) or toxæmic metritis can be characterized by additional signs of toxæmia (such as cold extremities, depression and/or collapse) which has a poor prognosis [5].

Diagnosis of clinical metritis

Diagnosis of clinical metritis (CM1 to CM3) is based on the clinical signs (purulent uterine discharge within 21 days after calving (CM1), fetid watery red-brown uterine discharge, fever $>39.5^{\circ}\text{C}$, dullness and decreased milk yield (CM2), additional signs of toxæmia such as cold extremities, depression and/or collapse (CM3) and abnormally enlarged uterus detected by rectal palpation mainly during the first week (10 days) after calving and rare during the second week [6,10].

Diagnosis of clinical metritis in small farms is based on the presence of vaginal discharge because clinical symptoms are not characteristics. It is important to mention that vaginal discharge can be present only in 24-33% of CM2 cases [11] and this may explain why the farmers used to present CM2 only about 2% of calvings [12] instead of 10% when systematic examination (e.g. measuring rectal temperature) is used [6,13].

In contrast, to diagnose it in large herds may be more complicated. Measuring the body temperature daily during the first 10 days after calving and performing vaginal examination at least once between Days 2 to 10 may help diagnosing clinical metritis more accurately. Measuring only rectal temperature is not enough because some cows may have no elevated rectal temperature ($>39.5^{\circ}\text{C}$) and have CM2 puerperal metritis [4,8]. The rectal measures of body temperature can be influenced "by the procedure itself (up to 0.5°C), type of thermometer (up to 0.3°C), and the penetration depth into the rectum (up to 0.4°C difference between a penetration depth of 11.5 cm and 6.0 cm in one of the experiments). Differences in rectal temperature before and after defecation are minor ($<0.1^{\circ}\text{C}$). These results may indicate that some care is required in generalizing rectal measures of body temperature" [14]. Daily measurement of rectal temperature and the assessment of vaginal discharge (colour, smell and viscosity) may help the diagnostic accuracy of puerperal metritis [15]. Using an electronic nose (DiagNose, C-it, Zutphen, the Netherlands) may increase the accuracy for evaluating vaginal discharge odour however the system needs further development for the field use [16].

It is important to emphasize that this early vaginal examination must perform with great care using adequate lubricant and being as hygienic as possible. The accuracy of our diagnosis can be improved by monitoring milk yield because milk production in some cows is not increasing daily as expected after calving or there is a sudden drop in it. Reduced feeding activity during the pre-partum period can be a significant risk factor for developing puerperal metritis [17]. In contrast, only a moderate accuracy can be reached by measuring the serum concentrations of acute phase proteins (haptoglobin) in dairy cows (sensitivity: 50% to 79%, specificity: 54% to 87%, respectively) [18,19].

Treatment of clinical metritis

Early treatment of clinical metritis (especially puerperal metritis) may decrease the severity of genital disorders (endometritis, cystic ovarian disease), the predisposition of metabolic disorders (left displacement abomasums, ketosis) and other complications like pyelonephritis, arthritis, endocarditis, hepatic and pulmonary abscesses [20-24].

There are a great variety of treatment protocols such as intrauterine antimicrobial agents (oxytetracycline; ampicillin and cloxacillin), antiseptic chemicals (iodine solutions: 500 ml of 2% Lugol's iodine immediately after calving and again 6 hr later as a preventive measure or 200 mL of povidone iodine diluted in 2 L of distilled water [25]), organic certified product (Optimum UterFlush, Van Beek Natural Science, Orange City, IA) in organic dairy farms [25], systemic antibiotics (penicillin or one of its synthetic analogues /amoxicillin/ [26]; ampicillin, oxytetracycline, ceftiofur /third generation cephalosporin/: 2.2 mg/kg of body weight daily for 3 to 5 days; 2 doses of 6.6 mg ceftiofur crystalline free acid sterile suspension (CCFA-SS)/kg of body weight s.c. in the base of the ear at a 72 hr interval [27] or a single dose of CCFA-SS within 24 hr after calving as a preventive measure [28,29], ozone i.u. treatment [30], supportive therapy (nonsteroidal anti-inflammatory drugs such as flunixin meglumine [31], fluid therapy in case of dehydration, therapy with calcium and energy supplements in case of depressed appetite, and hormone therapy (oxytocin: 20 to 40 IU repeated every 3 hr to 6 hr within 48 hr to 72 hr after calving; PGF_{2 α} or its synthetic analogues) have been introduced in the field [32]. The prognosis for recovery from puerperal metritis (CM2) varies with severity of the condition.

According to our present knowledge intrauterine antimicrobial and antiseptic treatments cannot be recommended because of irritating the endometrium [32]. Routine use of hormone therapies (PGF_{2 α}) is also controversial and needs further confirmations. It seems that presently systemic antibiotic (ceftiofur) and supportive therapy can be recommended for the field [6,33]. Reviewing 21 current literature data dealing with the treatment of puerperal metritis, Haimerl and Heuwieser [34] found that most of the studies used ceftiofur (17 of 21) and only 7 studies observed clinical improvement and none of them found improved reproductive performance. It is also important to mention that there is a growing concern of using third and fourth generation cephalosporins in production animals [35] therefore as an alternative therapy amoxicillin (i.m.) and intrauterine oxytetracycline infusion [26] or intrauterine ozone therapy can be recommended [36]. Due to the prevalence of self-recovery of puerperal metritis antibiotic treatment protocol can be started on Day 5 after calving however this hypothesis needs to be confirmed in a field study [37].

Treatment of acute puerperal metritis with a single dose of flunixin meglumine in addition to antibiotic treatment had no beneficial effect on clinical cure, milk yield within 6 d after the first treatment, or reproductive performance [31,38], while according to Amiridis et al. [39], a single dose of flunixin meglumine (2.2 mg/kg BW) administered intravenously to cows with CM2 between Days 5 and 8 after calving accelerated the uterine involution and shortened the calving-to-first-estrous interval.

Retained Placenta

Retained placenta or retained foetal membranes may occur if the placenta has not been shed by 12 (24) hr after calving. Majority of the placenta used to be expelled within 6 hr-9 hr after calving (88.7%) [40]. The average incidence of retained placenta after normal calving used to be 7% between 3 (4) to 11 (12)% [41,42]. After abnormal delivery (e.g., twin pregnancy, Caesarean section, foetotomy, forced extraction of the foetus, abortion, premature calving) and in herds infected with brucellosis its incidence rate can range between 20 and 50% or even more. Several factors like genetic, nutritional, immunological and pathological ones may influence the separation of bovine placenta; however its aetiology is still not fully understood. As retained placenta predisposes the development of uterine infections (clinical metritis, as well as clinical and subclinical endometritis) [43], and causing a

decrease in milk production (decreased milk yield, milk from treated cows withheld) [4,44] and reproductive performance (increases in days open, services per conception, calving to first heat interval, days from calving to first service and culling rate) [45] therefore the aim of its therapy is to prevent its adverse side effects.

Treatment of retained placenta

The aim of the treatment for retained placenta is to reduce the occurrence of puerperal metritis and subsequently clinical and subclinical endometritis, to decrease milk losses, to reduce reproductive inefficiency, and to decrease veterinary expenses [9,44].

There are a great variety of treatment protocols (manual removal of retained membranes, intrauterine treatments with antibiotics or antiseptics /Lugol's iodine/ [46], ozone spray [47,48], hormones / oxytocin, prostaglandin/, ergot derivatives, calcium, injection of collagenase into the umbilical arteries, versus no treatment) recommended for the field [49]. However, all of these methods have some limited values in the treatment of retained placenta [41]. Recent findings confirm that systemic antibiotics (ceftiofur 1 mg/kg) without intrauterine manipulation and treatment can be as effective as conventional treatment [6]. This was also confirmed in a later study in febrile cows [50,51]. It seems that systemic antibiotic is effective if the selection of treatment based on fever which may reduce the use of antibiotics compared with intra-uterine antibiotics [51]. Treatment with oxytocin, PGF_{2α}, or calcium was not effective for the prevention of retained placenta [52] or did not hasten the passage of foetal membranes [53] while injection with a new Chinese herbal medicine (40 mL of Dang Hong Fu) into the standard acupoint GV-1 (located in the depression halfway between the anus and the ventral aspect of the coccygeal vertebrae) decreased the prevalence of retained placenta and the time until expulsion of foetal membranes [54].

Clinical endometritis

Clinical endometritis is characterized by the presence of purulent or mucopurulent uterine discharge detectable in the vagina, 21 days or more after calving, and is not accompanied by systemic signs [45,55]. Due to the fact that mostly there is no endometrial inflammation (endometritis) in case of purulent vaginal discharge (PVD) therefore according to Dubuc et al. [56] the PVD terminology should be used.

Diagnosis of clinical endometritis

There are several methods (transrectal palpation [45]), transrectal ultrasonography [57], histological examination of endometrial biopsies [5,58], manual vaginal examination [5], vaginoscopy [45] available in the field to diagnose clinical endometritis in the cow however each method has some limitations. In a recent study it was confirmed that vaginoscopy was a practical tool to distinguish healthy from diseased cows with clinical endometritis [59].

In contrast, Metrichick^R (Simcro, New Zealand) consisting of a stainless steel rod with a rubber hemisphere can be used to retrieve vaginal contents more easily and precisely. In a recent study three methods (vaginoscopy /reference method/, gloved hand and Metrichick) were compared for diagnosing clinical endometritis between Days 21 to 27 after calving and it was confirmed that somewhat more cows (47.5%) could be diagnosed by Metrichick than by the other two methods (vaginoscopy: 36.9%, gloved hand: 36.8%). At the same time it did not result in improved reproductive performance [60]. On the other hand cytobrush cytology is also a reliable method for diagnosing clinical endometritis in cattle [61] however it is not so practical in the field.

The character and the odour of the vaginal mucus [4] can be scored according to the followings:

Mucus character

Score 0=clear or translucent mucus;

Score 1=mucus containing flecks of white or off-white pus;

Score 2=discharge containing ≤ 50% white or off-white mucopurulent material;

Score 3=discharge containing ≥ 50% purulent material, usually white or yellow, but occasionally sanguineous.

Mucus odour

Score 0: no unpleasant odour

Score 3: Fetid odour

In the absence of a gold standard it seems that vaginoscopy or Metrichick is preferred to use as a cow-side diagnostic tool for diagnosing clinical endometritis in the field [62].

Treatment of clinical endometritis

The general principle of the treatment of clinical endometritis is to reduce the load of pathogenic bacteria and enhance uterine defence and repair mechanisms and hence halt and reverse inflammatory changes that impair fertility [63]. A wide variety of therapy has been reported for clinical endometritis, including systemically or locally administered antibiotics, locally administered antiseptic solution and/or systemically injected PGF_{2α}. Infusion of antimicrobials into the uterus is aimed at achieving high concentrations at the site of infection [64,65]. In contrast to systemic administration, intrauterine administration achieves higher drug concentration in the endometrium, but little penetration to deeper layers of the uterus or other genital tissues.

Intrauterine treatment with 0.5 g cephapirin, first-generation cephalosporin, at 24-42 days before the planned start of mating improved reproductive performance of dairy cattle, especially those that had a history of retained placenta, a calf dead at calving or within 24 hr of calving, or vulval discharge [66]. In an experimental study, systemic administration of cefquinome (1 mg/kg), fourth generation cephalosporin, for three consecutive days was efficient for treatment *E. coli*-induced endometritis [67]. Intrauterine infusion of cephapirin or systemic administration of PGF_{2α} significantly improved the pregnancy rate of cows with clinical endometritis from which *T. pyogenes* was isolated [68]. A field study revealed that odd ratios for pregnancy after treatment of cows with clinical endometritis with PGF_{2α} or cephapirin on Days 28 to 35 postpartum were 1.5 and 1.9 (P<0.05), respectively as compared to the control [33]. In a large field study performed by LeBlanc et al. [45,69], there was no benefit on time to pregnancy of treatment of endometritis before 4 week postpartum. Moreover, administration of PGF_{2α} between 20 and 26 DIM to cows with endometritis that did not have a palpable CL was associated with a significant reduction in pregnancy rate. Cows with endometritis between 27 and 33 DIM, treated with cephapirin i.u. had a significantly shorter time to pregnancy than cows in the untreated groups. In cows with endometritis that had a palpable CL, there was no significant difference in time to pregnancy between those treated by intrauterine infusion of cephapirin or PGF_{2α}. Both groups tended to have a higher pregnancy rate than those in the untreated cows. Numerous reviewers have concluded that PGF_{2α} appears to be at least as effective for clinical endometritis as any available alternative therapy (Lugol's iodine [68], polyvinylpyrrolidone-iodine solution [70], metacresolsulphuric acid

and Lotagen [71]) and presents minimal risk of harm to the uterus or presence of residues in milk or meat [64,72,73].

In the absence of an active corpus luteum, the treatment efficacy of clinical endometritis with only prostaglandin injection is limited however such a treatment according to Lewis [74] may bring certain advantages as well.

It is important to mention that cows with clinical endometritis treated with one or 2 PGF_{2α} before initiation of the timed AI program had the lowest pregnancy rate per AI and the highest pregnancy loss compared with those having no uterine diseases [75]. A recent meta-analysis was also not able to confirm the benefit of PGF_{2α} treatment on reproductive performance of dairy cows with endometritis [76]. Similarly, 2 PGF_{2α} treatments between Days 37 ± 3 and 51 ± 3 after calving as part of an oestrous synchronization protocol and i.u. infusion of ceftiofur (125 mg) given at Day 44 ± 3 did not improve pregnancy per AI following the first postpartum insemination or the rate of pregnancy in the first 300 after calving [77].

Recently, as an alternative treatment of antimicrobials for clinical endometritis, 50% Dextrose [78] or proteolytic enzyme (containing trypsin, chymotrypsin and papain) solutions [79] were used however further investigations are needed to confirm the beneficial effects of such treatments [80].

Pyometra

Pyometra is characterized by the accumulation of purulent or mucopurulent material within the uterine lumen, distension of the uterus in the presence of an active corpus luteum and a closed cervix [5]. There are often an increased number of pathogenic bacteria within the uterine lumen when the corpus luteum forms and pyometra occurs [81]. Although there is a functional closure of the cervix, the lumen is not always completely occluded and some pus may be discharged through the cervix into the vaginal lumen.

The diagnosis of pyometra can be based on transrectal palpation of a distended uterus and/or on transrectal ultrasonography of mixed echodensity fluid and the presence of a persistent corpus luteum, with a history of anoestrus [5].

Treatment of pyometra

The best treatment protocol is to use prostaglandin (PGF_{2α} or its synthetic analogues) injection(s) because of the presence of a persistent corpus luteum. Due to common relapse it is recommended to repeat the prostaglandin treatment 12 to 14 days later. Intra-uterine antibiotic therapy (cephapirin) may be used as well. Complete restoration of the endometrium may need 4 to 8 weeks therefore it is very important to diagnose and treat pyometra as soon as possible after calving to decrease the destructive nature of pyometra on the endometrium [73].

Subclinical Endometritis

Subclinical endometritis can be diagnosed by uterine cytology in the absence of purulent discharge in the vagina. Endometrial and inflammatory cells may be collected by uterine lavage [82,83], cytobrush [82,83] or cytotape techniques [84] to evaluate the presence of PMNs in the uterine sample. If >18% PMNs in uterine cytology samples collected between Days 21 and 33 after calving, or >10% PMNs at Days 34 to 47 after calving can be found, in the absence of clinical endometritis, subclinical endometritis can be defined [82]. If >5% PMNs in uterine cytology samples collected by uterine lavage between Days 40 and 60 after calving can be found, subclinical endometritis is defined [85]. It is important to mention that there are a wide range of cut-off values to diagnose subclinical endometritis in the field [62,86,87].

By comparing the cytobrush technique, cytotape provided higher quality cells and significantly less blood contaminations while the PMNs percentage and the total number of cells did not differ [84].

Subclinical (cytologic) endometritis can also be diagnosed by using urinary test strips (Multistix 10 SG; Bayer Corporation, Elkart, IN, USA) on uterine lavage samples which contains leukocyte esterase, protein, and pH tests, however, in comparison with conventional cytology, 20.6% of cows could not be accurately classified therefore the reagent strip as an alternative test may require further confirmation [20]. It is worth mentioning that besides the difficulties of uterine biopsy in the field it can give low agreement with cytology for the diagnosis of subclinical endometritis [86].

Treatment of subclinical endometritis

Subclinical endometritis can be treated with a prostaglandin i.m. injection (cloprostenol 500 mg) or/and an i.u. antibiotic therapy (cephapirin) at 20–33 DIM to improve the reproductive performance [88]. Intrauterine infusion of ceftiofur hydrochloride reduced the prevalence of *T. pyogenes*, but did not affect fertility of dairy cows already receiving PGF_{2α} [77]. One or 2 treatments with PGF_{2α} before initiation of the timed AI program with subclinical endometritis were unable to improve uterine health, pregnancy rate per AI, and maintenance of pregnancy in lactating dairy cows [75]. Intrauterine lavage with 500-600 ml of sterile physiological saline (35°C-40°C) at Day 30 after calving may significantly decrease the number of PMNs in the uterus and improve pregnancy rate however it requires further large scale confirmations [35].

Prevention of Uterine Diseases

Cows having hypocalcaemia, dystocia, stillbirth, twins or retained placenta in the periparturient period are more likely to contract uterine infections than those cows that calve normally. Thus, management of sanitation, nutrition, population density, stress to prevent or reduce the incidence of these predisposing factors (especially dystocia) should be impeccable. Therefore prevention remains limited to general guidance on hygiene at calving [89], adequate nutrition (Ca, Se, Vitamin E, etc.) and the control of infectious diseases.

Routine systemic or intra-uterine administration of ceftiofur may be beneficial for the prevention of clinical metritis, however its effect on reproductive performance is not significantly different to that of no treatment therefore it cannot be recommended for the field [6,49,51,90]. Similarly controversial results were reported when a single-dose of ceftiofur crystalline free acid sterile suspension was used in dairy cows at high risk of uterine disease (twin, dystocia, or retained placenta) within 24 hr after calving [28,29].

One of the pharmacological approaches to the prevention and treatment of retained placenta can be the administration of prostaglandin immediately after calving [91], however due to controversy results further studies are needed to confirm its efficacy. Similarly the advantage of treatment with a new Chinese herbal aqua-acupuncture formulation Dang Hong Fu [54] needs further confirmation. Repeated administration of PGF_{2α} to cows on Days 7 and 14 or on Days 22 and 35 after calving had no effect on the prevalence of clinical endometritis at Days 22 and 58 after calving, and there was no effect on the probability of pregnancy after insemination at oestrus among cows with a voluntary waiting period of >100 days, or at timed AI at Day 85 when Presynch was performed [92]. Similarly preventive administration of PGF_{2α} at both 5 and 7 wk after calving had no positive effect on reproduction in dairy cows [28]. Preventive ozone intrauterine (spray) treatment [47,93], Sheng Hua Tang (a classical

herbal formula consisting of Radix Angelicae Sinensis, Ligustici Rhizoma, Semen Persicae, Zingiberis Rhizoma, and Radix Glycyrrhizae [94] or herbal tincture containing Herba Leonuri, Angelicae Sinensis Radix, Flos Carthami, Myrrha and Rhizoma Cyperi [95] during early puerperal period may improve the reproductive efficacy in dairy cows. In contrast, homeopathic drugs like Lachesis compositum (Lachesis), Carduus compositum (Carduus), and Traumeel LT (Traumeel) were not effective in preventing bovine endometritis or in enhancing reproductive performance in dairy cows [96].

On the other hand subcutaneous vaccination on Days 230 and 260 of pregnancy with inactivated bacterial components and/or protein subunits of *E. coli*, *E. necrophorum* and *T. pyogenes* may prevent puerperal metritis during the first lactation of dairy cows, leading to improved reproduction however further studies are needed to confirm the benefit of vaccination in the field [97].

References

1. Silva JW (2003) Addressing the decline in reproductive performance of lactating dairy cows: a researcher's perspective. *Veterinary Science Tomorrow* 3: 1-5.
2. Britt JH (1975) Early postpartum breeding in dairy cows. A review. *J Dairy Sci* 58: 266-271.
3. Szenci O (2008) Factors, which may affect reproductive performance in dairy cattle. *Magy Állatorv Lapja* 130: 107-111.
4. Sheldon IM, Dobson H (2004) Postpartum uterine health in cattle. *Anim Reprod Sci* 82-83: 295-306.
5. Sheldon IM, Lewis GS, LeBlanc S, Gilbert RO (2006) Defining postpartum uterine disease in cattle. *Theriogenology* 65: 1516-1530.
6. Drillich M, Beetz O, Pfütznner A, Sabin M, Sabin HJ, et al. (2001) Evaluation of a systemic antibiotic treatment of toxic puerperal metritis in dairy cows. *J Dairy Sci* 84: 2010-2017.
7. Földi J, Kulcsár M, Pécsi A, Huyghe B, de Sa C, et al. (2006) Bacterial complications of postpartum uterine involution in cattle. *Anim Reprod Sci* 96: 265-281.
8. Benzaquen ME, Risco CA, Archbald LF, Melendez P, Thatcher MJ, et al. (2007) Rectal temperature, calving-related factors, and the incidence of puerperal metritis in postpartum dairy cows. *J Dairy Sci* 90: 2804-2814.
9. Konyves L, Szenci O, Jurkovich V, Tegzes L, Tirián A, et al. (2009) Risk assessment and consequences of retained placenta for uterine health, reproduction and milk yield in dairy cows. *Acta Veterinaria Brno* 78: 163-172.
10. Markusfeld O (1984) Factors responsible for post parturient metritis in dairy cattle. *Vet Rec* 114: 539-542.
11. Zhou C, Boucher JF, Dame KJ, Moreira M, Graham R, et al. (2001) Multilocation trial of ceftiofur for treatment of postpartum cows with fever. *J Am Vet Med Assoc* 219: 805-808.
12. Bareille N, Beaudeau F, Billon S, Robert A, Faverdin P (2003) Effects of health disorders on feed intake and milk production in dairy cows. *Livestock Production Science* 83: 53-62.
13. Bareille N, Fourichon C (2006) Facteurs de risque des affections post-partum. *Point Vét* 37: 116-121.
14. Burfeind O, von Keyserlingk MA, Weary DM, Veira DM, Heuwieser W (2010) Short communication: repeatability of measures of rectal temperature in dairy cows. *J Dairy Sci* 93: 624-627.
15. Sannmann I, Heuwieser W (2015) Technical note: Intraobserver, interobserver, and test-retest reliabilities of an assessment of vaginal discharge from cows with and without acute puerperal metritis. *J Dairy Sci* 98: 5460-5466.
16. Sannmann I, Burfeind O, Suthar V, Bos A, Bruins M, et al. (2013) Technical note: Evaluation of odor from vaginal discharge of cows in the first 10 days after calving by olfactory cognition and an electronic device. *J Dairy Sci* 96: 5773-5779.
17. Urton G, von Keyserlingk MA, Weary DM (2005) Feeding behavior identifies dairy cows at risk for metritis. *J Dairy Sci* 88: 2843-2849.
18. Burfeind O, Sannmann I, Voigtsberger R, Heuwieser W (2014) Receiver operating characteristic curve analysis to determine the diagnostic performance of serum haptoglobin concentration for the diagnosis of acute puerperal metritis in dairy cows. *Anim Reprod Sci* 149: 145-151.
19. Huzzey JM, Duffield TF, LeBlanc SJ, Veira DM, Weary DM, et al. (2009) Short communication: Haptoglobin as an early indicator of metritis. *J Dairy Sci* 92: 621-625.
20. Cheong SH, Nydam DV, Galvão KN, Crosier BM, Ricci A, et al. (2012) Use of reagent test strips for diagnosis of endometritis in dairy cows. *Theriogenology* 77: 858-864.
21. Gröhn YT, Erb HN, McCulloch CE, Saloniemi HS (1990) Epidemiology of reproductive disorders in dairy cattle: associations among host characteristics, disease and production. *Preventive Veterinary Medicine* 8: 25-39.
22. Gröhn YT, Rajala-Schultz PJ (2000) Epidemiology of reproductive performance in dairy cows. *Anim Reprod Sci* 60-61: 605-614.
23. Mateus L, da Costa LL, Bernardo F, Silva JR (2002) Influence of puerperal uterine infection on uterine involution and postpartum ovarian activity in dairy cows. *Reprod Domest Anim* 37: 31-35.
24. Opsomer G, Gröhn YT, Hertl J, Coryn M, Deluyker H, et al. (2000) Risk factors for post partum ovarian dysfunction in high producing dairy cows in Belgium: a field study. *Theriogenology* 53: 841-857.
25. Pinedo PJ, Velez JS, Bothe H, Merchan D, Piñeiro JM, et al. (2015) Effect of intrauterine infusion of an organic-certified product on uterine health, survival, and fertility of dairy cows with toxic puerperal metritis. *J Dairy Sci* 98: 3120-3132.
26. Armengol R, Fraile L (2015) Comparison of two treatment strategies for cows with metritis in high-risk lactating dairy cows. *Theriogenology* 83: 1344-1351.
27. McLaughlin CL, Stanisiewski E, Lucas MJ, Cornell CP, Watkins J, et al. (2012) Evaluation of two doses of ceftiofur crystalline free acid sterile suspension for treatment of metritis in lactating dairy cows. *J Dairy Sci* 95: 4363-4371.
28. Dubuc J, Duffield TF, Leslie KE, Walton JS, Leblanc SJ (2011) Randomized clinical trial of antibiotic and prostaglandin treatments for uterine health and reproductive performance in dairy cows. *J Dairy Sci* 94: 1325-1338.
29. McLaughlin CL, Stanisiewski EP, Risco CA, Santos JE, Dahl GE, et al. (2013) Evaluation of ceftiofur crystalline free acid sterile suspension for control of metritis in high-risk lactating dairy cows. *Theriogenology* 79: 725-734.
30. Đuricic D, Lipar M, Samardžija M (2014) Ozone treatment of metritis and endometritis in Holstein cows. *Vet Arhiv* 84: 103-110.
31. Drillich M, Voigt D, Forderung D, Heuwieser W (2007) Treatment of acute puerperal metritis with flunixin meglumine in addition to antibiotic treatment. *J Dairy Sci* 90: 3758-3763.
32. Risco CA, Youngquist RS, Shore MD (2007) Postpartum uterine infections. In: Youngquist RS, Threlfall WR (eds.), *Current therapy in large animal theriogenology*. W.B. Saunders Company, Philadelphia.
33. Földi J, Pécsi A, Szabó J, Pécsi T, Huyghe B, et al. (2009) Use of cephalosporins for the treatment of dairy cows suffering of puerperal metritis and endometritis (in Hungarian with English summary). *Magy Állatorv Lapja* 131: 451-455.
34. Haimel P, Heuwieser W (2014) Invited review: Antibiotic treatment of metritis in dairy cows: a systematic approach. *J Dairy Sci* 97: 6649-6661.
35. Dini P, Farhoodi M, Hostens M, van Eetvelde M, Pascottini OB, et al. (2015) Effect of uterine lavage on neutrophil counts in postpartum dairy cows. *Anim Reprod Sci* 158: 25-30.
36. Samardžija M, Valpotic H, Đuricic D, Szenci O (2015) Intrauterine use of ozone in domestic ruminants. *Magy Állatorv Lapja* 137: 193-206.
37. Sannmann I, Burfeind O, Voigtsberger R, Heuwieser W (2013) Comparison of two monitoring and treatment strategies for cows with acute puerperal metritis. *Theriogenology* 79: 961-969.
38. Königsson K, Gustafsson H, Gunnarsson A, Kindahl H (2001) Clinical and bacteriological aspects on the use of oxytetracycline and flunixin in primiparous cows with induced retained placenta and post-partal endometritis. *Reprod Domest Anim* 36: 247-256.
39. Amiridis GS, Leontides L, Tassos E, Kostoulas P, Fthenakis GC (2001) Flunixin meglumine accelerates uterine involution and shortens the calving-to-first-oestrus interval in cows with puerperal metritis. *J Vet Pharmacol Ther* 24: 365-367.
40. van Werven T, Schukken YH, Lloyd J, Brand A, Heeringa HTJ, et al. (1992) The effects of duration of retained placenta on reproduction, milk production, postpartum disease and culling rate. *Theriogenology* 37: 1191-1203.

41. Eiler H, Fecteau KA (2007) Retained placenta. In: Youngquist RS, Threlfall WR (eds), Current therapy in large animal theriogenology. W.B. Saunders Company, Philadelphia.
42. Grunert E (1986) Etiology and pathogenesis of retained bovine placenta. In: Morrow AD (eds), Current therapy in theriogenology. W.B. Saunders Company, Philadelphia.
43. Dohmen MJ, Joop K, Sturk A, Bols PE, Lohuis JA (2000) Relationship between intra-uterine bacterial contamination, endotoxin levels and the development of endometritis in postpartum cows with dystocia or retained placenta. *Theriogenology* 54: 1019-1032.
44. Konyves L, Szenci O, Jurkovich V, Tegzes L, Tirián A, et al. (2009) Risk assessment of postpartum uterine disease and consequences of puerperal metritis for subsequent metabolic status, reproduction and milk yield in dairy cows. *Acta Vet Hung* 57: 155-169.
45. LeBlanc SJ, Duffield TF, Leslie KE, Bateman KG, Keefe GP, et al. (2002) Defining and diagnosing postpartum clinical endometritis and its impact on reproductive performance in dairy cows. *J Dairy Sci* 85: 2223-2236.
46. Ahmed FO, Elsheikh AS (2013) Intrauterine infusion of Lugol's iodine improves the reproductive traits of postpartum infected dairy cows. *Journal of Agriculture and Veterinary Science* 5: 89-94.
47. Djuricic D, Vince S, Ablondi M, Dobranic T, Samardzija M (2012) Effect of preventive intrauterine ozone application on reproductive efficiency in Holstein cows. *Reprod Domest Anim* 47: 87-91.
48. Zobel R, Tkalčić S (2013) Efficacy of ozone and other treatment modalities for retained placenta in dairy cows. *Reprod Domest Anim* 48: 121-125.
49. Risco CA, Hernandez J (2003) Comparison of ceftiofur hydrochloride and estradiol cypionate for metritis prevention and reproductive performance in dairy cows affected with retained fetal membranes. *Theriogenology* 60: 47-58.
50. Drillich M, Mahlstedt M, Reichert U, Tenhagen BA, Heuwieser W (2006) Strategies to improve the therapy of retained fetal membranes in dairy cows. *J Dairy Sci* 89: 627-635.
51. Drillich M, Reichert U, Mahlstedt M, Heuwieser W (2006) Comparison of two strategies for systemic antibiotic treatment of dairy cows with retained fetal membranes: preventive vs. selective treatment. *J Dairy Sci* 89: 1502-1508.
52. Hernandez J, Risco CA, Elliott JB (1999) Effect of oral administration of a calcium chloride gel on blood mineral concentrations, parturient disorders, reproductive performance, and milk production of dairy cows with retained fetal membranes. *J Am Vet Med Assoc* 215: 72-76.
53. Frazer GS (2005) A rational basis for therapy in the sick postpartum cow. *Vet Clin North Am Food Anim Pract* 21: 523-568.
54. Luo C, Li J, Wang J, Zheng J, Hua L, et al. (2010) Reduction of the incidence of retained placenta in cows treated with a new Chinese herbal medicine Dang Hong Fu used as aqua-acupuncture at GV-1. *AJTVM* 5: 27-34.
55. Sheldon IM, Noakes DE (1998) Comparison of three treatments for bovine endometritis. *Vet Rec* 142: 575-579.
56. Dubuc J, Duffield TF, Leslie KE, Walton JS, LeBlanc SJ (2010) Definitions and diagnosis of postpartum endometritis in dairy cows. *J Dairy Sci* 93: 5225-5233.
57. Kamimura S, Ohgi T, Takahashi M, Tsukamoto T (1993) Postpartum resumption of ovarian activity and uterine involution monitored by ultrasonography in Holstein cows. *J Vet Med Sci* 55: 643-647.
58. Bonnet BN, Martin SW, Meek AH (1993) Associations of clinical findings, bacteriological and histological results of endometrial biopsy with reproductive performance of postpartum dairy cows. *Prev Vet Med* 15: 205-220.
59. Leuter C, von Krueger X, Plöntzke J, Heuwieser W (2012) Evaluation of vaginoscopy for the diagnosis of clinical endometritis in dairy cows. *J Dairy Sci* 95: 206-212.
60. Pleticha S, Drillich M, Heuwieser W (2009) Evaluation of the Metrichk device and the gloved hand for the diagnosis of clinical endometritis in dairy cows. *J Dairy Sci* 92: 5429-5435.
61. Barlund CS, Carruthers TD, Waldner CL, Palmer CW (2008) A comparison of diagnostic techniques for postpartum endometritis in dairy cattle. *Theriogenology* 69: 714-723.
62. de Boer MW, LeBlanc SJ, Dubuc J, Meier S, Heuwieser W, et al. (2014) Invited review: Systematic review of diagnostic tests for reproductive-tract infection and inflammation in dairy cows. *J Dairy Sci* 97: 3983-3999.
63. LeBlanc SJ (2008) Postpartum uterine disease and dairy herd reproductive performance: a review. *Vet J* 176: 102-114.
64. Gilbert RO, Schwark WS (1992) Pharmacologic considerations in the management of peripartum conditions in the cow. *Vet Clin North Am Food Anim Pract* 8: 29-56.
65. Gustafsson BK (1984) Therapeutic strategies involving antimicrobial treatment of the uterus in large animals. *J Am Vet Med Assoc* 185: 1194-1198.
66. McDougall S, Macaulay R, Compton C (2007) Association between endometritis diagnosis using a novel intravaginal device and reproductive performance in dairy cattle. *Anim Reprod Sci* 99: 9-23.
67. Amiridis GS, Fthenakis GC, Dafopoulos J, Papanikolaou T, Mavrogianni VS (2003) Use of ceftiofur for prevention and treatment of bovine endometritis. *J Vet Pharmacol Ther* 26: 387-390.
68. Pécsi A, Földi J, Szabó J, Nagy P, Kulcsár M, et al. (2007) Efficacy of different antimicrobial therapeutic protocols for treatment of puerperal metritis and endometritis in dairy cows (in Hungarian with English summary). *Magy Állatorv Lapja* 129: 590-599.
69. LeBlanc LJ, Duffield T, Leslie K, Bateman K, Keefe G, et al. (2002) The effect of treatment of clinical endometritis on reproductive performance in dairy cows. *J Dairy Sci* 85: 2237-2249.
70. Nakao T, Moriyoshi M, Kawata K (1988) Effect of postpartum intrauterine treatment with 2% polyvinyl-pyrrolidone-iodine solution on reproductive efficiency in cows. *Theriogenology* 30: 1033-1043.
71. Heuwieser W, Tenhagen BA, Tischer M, Lühr J, Blum H (2000) Effect of three programmes for the treatment of endometritis on the reproductive performance of a dairy herd. *Vet Rec* 146: 338-341.
72. Gilbert RO (1992) Bovine endometritis: the burden of proof. *Cornell Vet* 82: 11-14.
73. Paisley LG, Mickelsen WD, Anderson PB (1986) Mechanisms and therapy for retained fetal membranes and uterine infections of cows: A review. *Theriogenology* 25: 353-381.
74. Lewis GS (2004) Steroidal regulation of uterine immune defenses. *Anim Reprod Sci* 82-83: 281-294.
75. Lima FS, Bisinotto RS, Ribeiro ES, Greco LF, Ayres H, et al. (2013) Effects of 1 or 2 treatments with prostaglandin F_{2a} on subclinical endometritis and fertility in lactating dairy cows inseminated by timed artificial insemination. *J Dairy Sci* 96: 6480-6488.
76. Haimerl P, Heuwieser W, Artl S (2013) Therapy of bovine endometritis with prostaglandin F_{2a}: a meta-analysis. *J Dairy Sci* 96: 2973-2987.
77. Galvão KN, Greco LF, Vilela JM, Sá Filho MF, Santos JE (2009) Effect of intrauterine infusion of ceftiofur on uterine health and fertility in dairy cows. *J Dairy Sci* 92: 1532-1542.
78. Brick TA, Schuenemann GM, Bas S, Daniels JB, Pinto CR, et al. (2012) Effect of intrauterine dextrose or antibiotic therapy on reproductive performance of lactating dairy cows diagnosed with clinical endometritis. *J Dairy Sci* 95: 1894-1905.
79. Drillich M, Raab D, Witke M, Heuwieser W (2005) Treatment of chronic endometritis in dairy cows with an intrauterine application of enzymes. A field trial. *Theriogenology* 63: 1811-1823.
80. Sassi G, Ismail S, Bajcsy ÁC, Kiss G, Répási A, et al. (2010) Evaluation of the alternatives of the intrauterine antibiotic treatments in the cow. Literature review (in Hungarian with English summary). *Magy Állatorv Lapja* 132: 516-527.
81. Noakes DE, Wallace LM, Smith GR (1990) Pyometra in a Friesian heifer: bacteriological and endometrial changes. *Vet Rec* 126: 509.
82. Kasimanickam R, Duffield TF, Foster RA, Gartley CJ, Leslie KE, et al. (2005) A comparison of the cytobrush and uterine lavage techniques to evaluate endometrial cytology in clinically normal postpartum dairy cows. *Can Vet J* 46: 255-259.
83. Kasimanickam R, Duffield TF, Foster RA, Gartley CJ, Leslie KE, et al. (2004) Endometrial cytology and ultrasonography for the detection of subclinical endometritis in postpartum dairy cows. *Theriogenology* 62: 9-23.
84. Pascottini OB, Dini P, Hostens M, Ducatelle R, Opsomer G (2015) A novel cytologic sampling technique to diagnose subclinical endometritis and comparison of staining methods for endometrial cytology samples in dairy cows. *Theriogenology* 84: 1438-1446.

85. Gilbert RO, Shin ST, Guard CL, Erb HN, Frajblat M (2005) Prevalence of endometritis and its effects on reproductive performance of dairy cows. *Theriogenology* 64: 1879-1888.
86. Madoz LV, Giuliadori MJ, Migliorisi AL, Jaureguiberry M, de la Sota RL (2014) Endometrial cytology, biopsy, and bacteriology for the diagnosis of subclinical endometritis in grazing dairy cows. *J Dairy Sci* 97: 195-201.
87. Melcher Y, Prunner I, Drillich M (2014) Degree of variation and reproducibility of different methods for the diagnosis of subclinical endometritis. *Theriogenology* 82: 57-63.
88. Kasimanickam R, Duffield TF, Foster RA, Gartley CJ, Leslie KE, et al. (2005) The effect of a single administration of cephalosporin or cloprostenol on the reproductive performance of dairy cows with subclinical endometritis. *Theriogenology* 63: 818-830.
89. Hartigan PJ (1980) Fertility management in the dairy herd: The need to control bacteriological contamination in the environment. *Irish Vet J* 34: 43-48.
90. Scott HM, Schouten MJ, Gaiser JC, Belschner AP, Jordan ER (2005) Effect of intrauterine administration of ceftiofur on fertility and risk of culling in postparturient cows with retained fetal membranes, twins, or both. *J Am Vet Med Assoc* 226: 2044-2052.
91. Stevens RD, Dinsmore RP, Cattell MB (1995) Evaluation of the use of intrauterine infusions of oxytetracycline, subcutaneous injections of fenprostalene, or a combination of both, for the treatment of retained fetal membranes in dairy cows. *J Am Vet Med Assoc* 207: 1612-1615.
92. Hendricks KE, Bartolome JA, Melendez P, Risco C, Archbald LF (2006) Effect of repeated administration of PGF2alpha in the early post partum period on the prevalence of clinical endometritis and probability of pregnancy at first insemination in lactating dairy cows. *Theriogenology* 65: 1454-1464.
93. Đuričić D, Valpotić H, Samardžija M (2015) Prophylaxis and therapeutic potential of ozone in buiatrics: Current knowledge. *Anim Reprod Sci* 159: 1-7.
94. Cui D, Li J, Wang X, Xie J, Zhang K, et al. (2014) Efficacy of herbal tincture as treatment option for retained placenta in dairy cows. *Anim Reprod Sci* 145: 23-28.
95. Cui D, Wang X, Wang L, Wang X, Zhang J, et al. (2014) The administration of Sheng Hua Tang immediately after delivery to reduce the incidence of retained placenta in Holstein dairy cows. *Theriogenology* 81: 645-650.
96. Art S, Padberg W, Drillich M, Heuwieser W (2009) Efficacy of homeopathic remedies as prophylaxis of bovine endometritis. *J Dairy Sci* 92: 4945-4953.
97. Machado VS, Bicalho MLdS, Meira Junior EBdS, Rossi R, Ribeiro BL, et al. (2014) Subcutaneous immunization with inactivated bacterial components and purified protein of *Escherichia coli*, *Fusobacterium necrophorum* and *Trueperella pyogenes* prevents puerperal metritis in Holstein dairy cows. *PLoS ONE* 9(3): e91734. doi:[10.1371/journal.pone.0091734](https://doi.org/10.1371/journal.pone.0091734)

Citation: Szenci O (2016) Recent Possibilities for Diagnosis and Treatment of Post Parturient Uterine Diseases in Dairy Cow. *JFIV Reprod Med Genet* 4: 170. doi:[10.4172/2375-4508.1000170](https://doi.org/10.4172/2375-4508.1000170)

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700 Open Access Journals
- 50,000 editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsgroup.org/journals/submission>