

2-Aminoethyl Diphenylborinate (2-APB) Analogues: Part 3 - Regulators of Huntington Aggregation and Transglutaminase

Shoichiro Ozaki*

The Institute of Physical and Chemical Research, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan

Abstract

Huntington aggregation inhibitory activities and transglutaminase inhibitory activities of 2APB analogues were measured. 2-APB analogues regulated the Huntington aggregation. This fact provided an example that 2-APB analogues can regulate cellular process. Diphenyl (aminoacidonate N,O) boranes, which are effective regulator of Ca²⁺ release and cellular process, were effective regulators of Huntington aggregation. It was also found that many of 2-APB analogues have moderate transglutaminase inhibition activities 2-Aminoethyl di(4-trifluorophenyl) borinate is a good transglutaminase regulator.

Keywords: 2-APB; 2-APB analogue; Regulator of transglutaminase; Regulator of huntington aggregation; Huntington disease

Introduction

Extracellular signal molecules attach to the plasmatic membrane where they are recognized by cell surface receptors. Upon binding of the ligand to the appropriate receptor, activation of G protein activates in turn phospholipase C. Active phospholipase C hydrolyzes phosphatidylinositol 4,5-bisphosphate (PIP₂) giving rise to two products: 1,2-diacylglycerol and inositol 1,4,5-triphosphate (IP₃). IP₃ stimulates the release of Ca²⁺ from the intracellular stores in the endoplasmic reticulum through IP₃ receptor while regulating a wide range of cellular processes [1-20].

In 1997, we identified 2-aminoethyl diphenylborinate (2-APB) as being an IP₃ receptor inhibitor and regulate IP₃-induced calcium release [21,22]. This discovery rose a substantial interest and had a great impact as it gained more than 600 citations and more than 1000 studies on 2-APB [23-37] have been published so far. This was supported by increasing sales of 2-APB by Sigma-Aldrich as membrane-permeable modulator of intracellular IP₃-induced cellular calcium release. We aimed to generate better modulator of calcium release than 2-APB. And we synthesized several 2-APB analogues and measured their inhibitory activities on Store-Operated Calcium Entry (SOCE) and IP₃ Induced Calcium Release (IICR). We found that bis boron compound DBP 161 and DBP 163 were 10 times more effective than 2-APB [38]. Previously, we studied bis- boron compounds in more detail [39,40]. We extended these studies and synthesized 493 2-APB analogues using methods described by us [38-43] and others [44-49] and measured their inhibitory activities on SOCE and IICR [38-43]. We found that these 2-APB analogues regulate Ca²⁺ release and associated cellular processes. We reported these finding on mono-boron compounds [50] and on bis-boron compounds [51].

This time, we report about 2 tests how these 2APB analogues show the ability to regulator Ca²⁺ related enzyme inhibitory activity and regulation of cellular process. One test is a regulation of transglutaminase, Second test is a regulation of Huntington aggregation using 2APB analogs having various Ca²⁺ release-related activities: SOCE. We wish to tell the relations of transglutaminase activity (TG), Huntington aggregation inhibition (x-Fold) and SOCE and chemical structures of compounds.

Transglutaminases (TG) [52-54] are calcium-dependent enzymes that catalyze various post-translational modifications including protein cross-linking, amine incorporation, and deamidation. The protein-crosslink is formed with isopeptide bonds between the carboxamide group of glutamine residues and the β-amino group of lysine residues to form N-(γ-glutamyl)-L-lysine accompanied by loss of the ammonia [55]. TG is implicated in various pathological roles in many neurodegenerative diseases [56-58] including Alzheimer's disease [59], Huntington's disease [60-63]. TG is also implicated in diseases like cataractous lens [64], Psoriasis skin injury [65], liver injury [66], fibrin injury [67,68], immune system injury [69,70] and Celiac disease [71]. Numerous transglutaminase inhibitors are reported [72-75]. We have reported aryl-β-amino ketones [76], dithio β-amino ketones [77] as transglutaminase inhibitors.

We measured transglutaminase (TG) inhibitory activities and Huntington aggregation inhibitory activities (x-Fold) of 276 2APB analogues. And we analyzed the result and relation of TG, x-Fold and SOCE IC₅₀.

Materials and Methods

APB analogues

2-APB was first synthesized by Ronderstvent et al. [44] in 1954 from triphenylboranes and ethanolamine. Later, hydroxy diphenyl borane and ethanolamine methods for 2-APB synthesis were reported by Weidman and Zimmermann [45], Letsinger and Skoog [46], Povlock and Lippincott [47].

We have synthesized 493 2-APB analogues [38-43] using methods described by us [38-43] and others [44-49]. The structures, names and synthetic methods of the compounds are in example 1-493 [43].

*Corresponding author: Shoichiro Ozaki, The Institute of Physical and Chemical Research, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan, Tel: 81-0467670991; E-mail: ozaki-0991@m.jcnnet.jp

Received June 5, 2014; Accepted September 9, 2014; Published September 16, 2014

Citation: Ozaki S (2014) 2-Aminoethyl Diphenylborinate (2-APB) Analogues: Part 3 - Regulators of Huntington Aggregation and Transglutaminase. J Bioengineer & Biomedical Sci 4: 131. doi:10.4172/2155-9538.1000131

Copyright: © 2014 Ozaki S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Methods

We measured transglutaminase (TG) inhibitory activities as described at ref 76 and Huntington aggregation inhibitory activities (x-Fold) [62].

Transglutaminase inhibitory activities measurement

TG (TransGlutaminase) inhibiting activity assay: Inhibition of TG enzyme was determined by assaying the enzyme activity in accordance with an optionally modified version of the enzyme activity in accordance with an optionally modified version of the method of Lorand et al. [53]. An enzyme reaction solution (0.1 ml) (100 mM HEPES-NaOH, pH 7.5, 1 mM CaCl₂, 20 M, monodansyl cadaverine, 0.05 mg/ml N,N-dimethylcasein, 5 µg/ml TGase) was introduced into wells of a 86-well plate (Nunc, 96 well Black Plate with Clear Bottom). A test compound was added in concentration of 100 µM. The plate was set in the fluorescence drug screening system FDSS3000 (hamamatsu Photonics K.K) TGase-inhibiting activity of the compound was calculated by assaying changes in fluorescence wavelength (at 340 nm) per unit time. The assay level at which a fluorescence change was observed with the addition of DMSO (1 µl) used as control instead of the test compound was designated as 100. The assay level at which TGase activity decreased by half in the presence of the test compound was designated as TG 50.

Huntington aggregation inhibitory activities measurement

Truncated N-terminal Huntington 150 Q-EGFP-Neuron 2a cells are cultured in 96 well plates. for 1day, 1 µM Ponasterone A (2 µl), 5 M Dibutyl cyclic AMP (2 µl) are added as to the test compound is to be 20 µM and cultured for 20 hours. The cells are fixed with 4% paraformaldehyde and washed with PBS after 30 minutes, and stained with Hoechst 33342. Aggregated cell numbers and total cell numbers are counted by Array Scan V T1 (Cellomics company, Pittsburg, USA). Ratio of aggregated cell to total cells (x-Fold) was measured. Without test compounds, ratio of aggregated cell to total cell x-Fold is 1. The smaller the x-Fold, the aggregation inhibition is stronger.

SOCE inhibition activities measurement

Inhibitory activities of the 2APB analogues for SOCE were measured using our improved assays as described previously [50].

Results and Discussion

We measured transglutaminase inhibitory activities (TG) and Huntington aggregation inhibition activity (x-Fold), and SOCE inhibition activities IC₅₀ of 2APB analogues. The results are shown in Supplementary Figure S1.

Typical 42 compounds are picked up from Figure S1 and grouped them into three groups and lined up from SOCE smaller value 0.2 to >10. 1) Amino acid adducts compounds, 2) Aminoalcohol adducts compounds & 3) Aminothioli adducts compounds.

Amino acid adduct compounds

Amino acid adduct compounds are majority of effective compounds. The compounds having small IC₅₀ (SOCE) like 0.2 showed strong Huntington aggregation inhibiting activities (x-Fold). TG has no relation with IC₅₀ (SOCE).

TG of 911 Diphenyl (2,6-diaminohexanate-O,N) borane is 136, x-Fold is 0.41, IC₅₀ is 0.2. TG of 919 Diphenyl (2,3-diaminopropionate-O,N) borane is 90 and x-Fold is 0.53, IC₅₀ is 0.2. TG of 855 Diphenyl

asparaginate-O,N- borane is 105 x-Fold is 0.54 and IC₅₀ is 0.2 (Figure 1).

The compounds having big IC₅₀ (SOCE) like 10 showed no transglutaminase activity and no Huntington aggregation activity. TG of 901 Diphenyl (methionate-O,N) borane is 106, x-Fold is 0.90, IC₅₀ is >10. TG of 4129 Diphenyl (2-aminohexanecarboxylate-O,N) borane is 90, x-Fold is 0.97, IC₅₀ is >10.

Aminoalcohol adduct compounds

This group compounds showed moderate transglutaminase inhibition and Huntington aggregation inhibition. 2APB belongs to this group. TG of 2APB is 80, x-Fold is 0.64, IC₅₀ is 3. TG of 1022 Bis-(4,4'-(phenyl aminoethoxy boryl)phenyl)ether is 4 and x-Fold is 0.50, SOCE is 0.2. TG of 424 2-aminoethyl di(4-trifluorophenyl) borinate is 54, x-Fold is 0.69, SOCE is 0.5. TG of 372 2-aminoethyl di(5-chloro-2-methyl-phenyl) borinate is 74, x-Fold is 0.78 and IC₅₀ is 1 (Figure 2).

Aminothioli adduct compounds

Some compounds of this group showed remarkable transglutaminase inhibition, showed no Huntington aggregation activities. TG of 6014 2-aminoethylthio di (4-chloro-2-fluorophenyl) borane is 28, x-Fold is 0.96, SOCE is <1. TG of 1031 2-aminoethylthio diphenyl borane is 33, x-Fold is 0.87, and SOCE is 2. TG of 3115 2-aminoethylthio di (3-chloro-4-methylphenyl) borane is 12, x-Fold is 1.01, SOCE is 2. TG of 6039 2-aminoethylthio di (4-cyanophenyl) borane is 23. X-fold is 0.92, SOCE is >10. This thiol adducts compounds showed toxicity. Cell numbers decreased during assay. Therefore this group compounds would be not suitable for clinical use.

Relations of chemical structures and activities

Active compounds of 2APB analogues, nitrogen atom must come in this order B-O-C-C-N as 2-APB C₆H₅B(OCH₂CH₂NH₂)C₆H₅. The compounds having other order like B-O-C-N, or B-O-C-C-C-N have no activities. When phenyl group is substituted with aliphatic or arylaliphatic group, they lost their activities. When compared mono-boron, bis-boron and poly-boron compounds, mono-boron compounds were best and bis-boron compounds come next.

Relation of SOCE IC₅₀ value and TG and x-Fold

Relation of SOCE IC₅₀ and x-Fold: When look at Figure S1, the compounds having strong Ca²⁺ release activity, low IC₅₀ value (IC₅₀<1) except aminothioli adduct compounds, showed strong Huntington aggregation inhibiting activity. The compounds having weak Ca²⁺ releasing activity (IC₅₀ is >10) showed weak or no transglutaminase inhibiting activity (TG is near 100), and showed weak or no Huntington aggregation inhibiting activity (x-Fold is near 1). These results indicated that 2APB analogues were effective as regulators of cellular process.

Relation of SOCE IC₅₀ and TG. Relation of TG and x-Fold: The compounds having strong Ca²⁺ release activity, low IC₅₀ value (IC₅₀<1) except aminothioli adduct compounds, showed no transglutaminase inhibition activities, x-Fold is near 1.00 even through IC₅₀ is 0.2. Because our assay of transglutaminase inhibition is done in vitro and the assay is not cellular process. Calcium release cannot be expected. Therefore we cannot find any relation between TG and SOCE IC₅₀ value also we cannot find any relation between TG and x-Fold. If we assay transglutaminase inhibition in vivo system, other results would be expected.

ID	Chemical Structure	TG	x-Fold	IC50	ID	Chemical Structure	TG	x-Fold	IC50
911		136	0.41	0.2	5142		51	0.97	1
919		90	0.53	0.2	847		84	0.87	3
855		109	0.54	0.2	916		102	0.41	3
5019		50	1.02	0.2	847		84	0.87	3
1024		24	0.6	0.2	877		94	1.01	>10
2051		140	0.99	0.2	901		106	0.93	>10
1023		66	0.59	0.3	4129		90	0.97	>10
855		109	0.54	0.5	921		94	0.81	>10
907		86	0.91	0.7	897		98	0.88	>10
4124		35	0.98	<1	869		92	0.99	10

Figure 1: Amino acid adducts compounds.




Figure 2: Amino alcohol adduct compounds & Aminothioli adduct compounds.

Regulator of huntington aggregation

Many compounds belong to amino acid adducts especially basic amino acid adduct such as 911 Diphenyl (2,6-diaminohexanoate O,N)borane TG:136, x-Fold:0.41, IC₅₀:0.2 and 919 Diphenyl (2,3-diaminopropionate O,N) borane TG:90, x-Fold:0.53, IC₅₀:0.2 showed strong Huntington aggregation inhibiting activity and no transglutaminase inhibiting activity.. These compound look like to be good and selective regulator of Huntington aggregation inhibitor.

Regulator of transglutaminase

Aminoalcohol adduct, 1022 Bis-(4,4' (phenyl aminoethoxy boryl) phenyl)ether: TG: 4 and x-Fold: 0.50 SOCE:0.2 and 424 2-aminoethyl di(4-trifluorophenyl) borinate: TG: 54, x-Fold :0.69, SOCE : 0.5 are good candidate as transglutaminase inhibitors. I have measured transglutaminase inhibition activities at 100 µM. The activities like these TG:4 or 54 at 100 µM are not so strong as aryl β-aminoethyl ketones IC₅₀: 0.1 µM reported by us [76,77] or thienopyrimidines IC₅₀: 0.13 µM reported by Duval [74]. Transglutaminase is necessary enzyme for our lives. Strong inhibitor will give toxicity. Moderate activity is required for clinical use.. We are reporting many TG inhibitors .differing activities. People will be able to choose most suitably active compounds. Some of these compounds were shown to inhibit the calcium dependent enzyme transglutaminase [43]. Transglutaminase inhibitors block the abnormal cross-link of protein [43,76-78] and they may slow down or even stop the progression of disease caused by over cross-linked proteins, such as Huntington's disease.

I have analyzed the SOCE and TG(transglutaminase inhibition activities and x-Fold (Huntington aggregation inhibition) of 262 2APB analogues . 2-APB analogues regulated the Huntington aggregation This fact provided an example that 2-APB analogues can regulate cellular process. DAB (Diphenyl aminoacidonate (N,O) boranes are best compounds to regulate Huntington aggregation . The 2APB analogues presented in this study could be excellent lead compound for therapeutically intervene in many diseases such as heart disease [26-28,32-36,79], Huntington disease [43,61,62], Celiac disease [71], Alzheimer disease [79] in connection with cross-link of protein.

Conclusion

2-APB analogues regulated the Huntington aggregation has provided an example that 2-APB analogues can regulate cellular process. Diphenyl (aminoacidonate N,O) boranes are effective regulators of Huntington aggregation. It was also found that many of 2-APB analogues had moderate transglutaminase inhibition activities.

References

- Berridge MJ, Dawson RM, Downes CP, Heslop JP, Irvine RF (1983) Changes in the levels of inositol phosphates after agonist-dependent hydrolysis of membrane phosphoinositides. *Biochem J* 212: 473-482.
- Berridge MJ (1983) Rapid accumulation of inositol trisphosphate reveals that agonists hydrolyse polyphosphoinositides instead of phosphatidylinositol. *Biochem J* 212: 849-858.
- Streb H, Irvine RF, Berridge MJ, Schulz I (1983) Release of Ca²⁺ from a nonmitochondrial intracellular store in pancreatic acinar cells by inositol-1,4,5-trisphosphate. *Nature* 306: 67-69.
- Berridge MJ, Heslop JP, Irvine RF, Brown KD (1984) Inositol trisphosphate formation and calcium mobilization in Swiss 3T3 cells in response to platelet-derived growth factor. *Biochem J* 222: 195-201.
- Fein A, Payne R, Corson DW, Berridge MJ, Irvine RF (1984) Photoreceptor excitation and adaptation by inositol 1,4,5-trisphosphate. *Nature* 311: 157-160.
- Brown JE, Rubin LJ, Ghalayini AJ (1984) A biochemical and electrophysiological examination of myo-inositol polyphosphate as a putative messenger for excitation in Limulus ventral photoreceptor cells. *Nature* 311: 160-163.
- Burgess GM, Godfrey PP, McKinney JS, Berridge MJ, Irvine RF, et al. (1984) The second messenger linking receptor activation to internal Ca release in liver. *Nature* 309: 63-66.
- Prentki M, Biden TJ, Janjic D, Irvine RF, Berridge MJ, et al. (1984) Rapid mobilization of Ca²⁺ from rat insulinoma microsomes by inositol-1,4,5-trisphosphate. *Nature* 309: 562-564.
- Irvine RF, Brown KD, Berridge MJ (1984) Specificity of inositol trisphosphate-induced calcium release from permeabilized Swiss-mouse 3T3 cells. *Biochem J* 222: 269-272.
- Irvine RF, Letcher AJ, Heslop JP, Berridge MJ (1986) The inositol tris/tetrakisphosphate pathway - demonstration of Ins 1,4,5P₃ 3-kinase activity in animal cells. *Nature* 320: 631-634.
- Rapp PE, Berridge MJ (1981) The control of transepithelial potential oscillations in the salivary gland of *Calliphora erythrocephala*. *Exp Biol* 93: 119-132.
- Missiaen L, Taylor CW, Berridge MJ (1991) Spontaneous calcium release from inositol trisphosphate-sensitive calcium stores. *Nature* 352: 241-244.
- Berridge MJ, Irvine RF (1984) Inositol trisphosphate, a novel second messenger in cellular signal transduction. *Nature* 312: 315-321.
- Berridge MJ (1987) Inositol trisphosphate and diacylglycerol: two interacting second messengers. *Annu Rev Biochem* 56: 159-193.
- Berridge MJ, Irvine RF (1989) Inositol phosphates and cell signalling. *Nature* 341: 197-205.
- Berridge MJ, Downes CP, Hanley MR (1989) Neural and developmental actions of lithium: a unifying hypothesis. *Cell* 59: 411-419.
- Berridge MJ (1993) Inositol trisphosphate and calcium signalling. *Nature* 361: 315-325.
- Bootman MD, Berridge MJ (1995) The elemental principles of calcium signaling. *Cell* 83: 675-678.
- Berridge MJ (1993) Cell signalling. A tale of two messengers. *Nature* 365: 388-389.
- Decrock E, De Bock M, Wang N, Gadicherla AK, Bol M, et al. (2013) IP₃, a small molecule with a powerful message. *Biochim Biophys Acta* 1833: 1772-1786.
- Maruyama T, Kanaji T, Nakade S, Kanno T, Mikoshiba K (1997) 2APB, 2-aminoethoxydiphenyl borate, a membrane-penetrable modulator of Ins(1,4,5)P₃-induced Ca²⁺ release. *J Biochem* 122: 498-505.
- Iwasaki H, Mori Y, Hara Y, Uchida K, Zhou H, et al. (2001) 2-Aminoethoxydiphenyl borate (2-APB) inhibits capacitative calcium entry independently of the function of inositol 1,4,5-trisphosphate receptors. *Receptors Channels* 7: 429-439.
- Bilmen JG, Michelangeli F (2002) Inhibition of the type 1 inositol 1,4,5-trisphosphate receptor by 2-aminoethoxydiphenylborate. *Cell Signal* 14: 955-960.
- Ma HT, Venkatachalam K, Parys JB, Gill DL (2002) Modification of store-operated channel coupling and inositol trisphosphate receptor function by 2-aminoethoxydiphenyl borate in DT40 lymphocytes. *J Biol Chem* 277: 6915-6922.
- Dobrydneva Y, Blackmore P (2001) 2-Aminoethoxydiphenyl borate directly inhibits store-operated calcium entry channels in human platelets. *Mol Pharmacol* 60: 541-552.
- Bilmen JG, Wootton LL, Godfrey RE, Smart OS, Michelangeli F (2002) Inhibition of SERCA Ca²⁺ pumps by 2-aminoethoxydiphenyl borate (2-APB). 2-APB reduces both Ca²⁺ binding and phosphoryl transfer from ATP, by interfering with the pathway leading to the Ca²⁺-binding sites. *Eur J Biochem* 269: 3678-3687.
- Missiaen L, Callewaert G, De Smedt H, Parys JB (2001) 2-Aminoethoxydiphenyl borate affects the inositol 1,4,5-trisphosphate receptor, the intracellular Ca²⁺ pump and the non-specific Ca²⁺ leak from the non-mitochondrial Ca²⁺ stores in permeabilized A7r5 cells. *Cell Calcium* 29: 111-116.

28. Peppiatt CM, Collins TJ, Mackenzie L, Conway SJ, Holmes AB, et al. (2003) 2-Aminoethoxydiphenyl borate (2-APB) antagonises inositol 1,4,5-trisphosphate-induced calcium release, inhibits calcium pumps and has a use-dependent and slowly reversible action on store-operated calcium entry channels. *Cell Calcium* 34: 97-108.
29. Luo D, Broad LM, Bird GS, Putney JW Jr (2001) Signaling pathways underlying muscarinic receptor-induced $[Ca^{2+}]$ oscillations in HEK293 cells. *J Biol Chem* 276: 5613-5621.
30. Bootman MD, Young KW, Young JM, Moreton RB, Berridge MJ (1996) Extracellular calcium concentration controls the frequency of intracellular calcium spiking independently of inositol 1,4,5-trisphosphate production in HeLa cells. *Biochem J* 314: 347-354.
31. Mackenzie L, Bootman MD, Berridge MJ, Lipp P (2001) Predetermined recruitment of calcium release sites underlies excitation-contraction coupling in rat atrial myocytes. *J Physiol* 530: 417-429.
32. Lipp P, Laine M, Tovey SC, Burrell KM, Berridge MJ, et al. (2000) Functional InsP3 receptors that may modulate excitation-contraction coupling in the heart. *Curr Biol* 10: 939-942.
33. Mackenzie L, Bootman MD, Laine M, Berridge MJ, Thuring J, et al. (2002) The role of inositol 1,4,5-trisphosphate receptors in Ca^{2+} signalling and the generation of arrhythmias in rat atrial myocytes. *J Physiol* 541: 395-409.
34. Proven A, Roderick HL, Conway SJ, Berridge MJ, Horton JK, et al. (2006) Inositol 1,4,5-trisphosphate supports the arrhythmogenic action of endothelin-1 on ventricular cardiac myocytes. *J Cell Sci* 119: 3363-3375.
35. Berridge MJ, Bootman MD, Roderick HL (2003) Calcium signalling: dynamics, homeostasis and remodelling. *Nat Rev Mol Cell Biol* 4: 517-529.
36. Berridge MJ (2006) Remodelling Ca^{2+} signalling systems and cardiac hypertrophy. *Biochem Soc Trans* 34: 228-231.
37. Berridge MJ (2006) Calcium microdomains: organization and function. *Cell Calcium* 40: 405-412.
38. Zhou H, Iwasaki H, Nakamura T, Nakamura K, Maruyama T, et al. (2007) 2-Aminoethyl diphenylborinate analogues: selective inhibition for store-operated Ca^{2+} entry. *Biochem Biophys Res Commun* 352: 277-282.
39. Suzuki AZ, Ozaki S, Goto J, Mikoshiba K (2010) Synthesis of bisboron compounds and their strong inhibitory activity on store-operated calcium entry. *Bioorg Med Chem Lett* 20: 1395-1398.
40. Goto J, Suzuki AZ, Ozaki S, Matsumoto N, Nakamura T, et al. (2010) Two novel 2-aminoethyl diphenylborinate (2-APB) analogues differentially activate and inhibit store-operated Ca^{2+} entry via STIM proteins. *Cell Calcium* 47: 1-10.
41. Mikoshiba K, Ozaki S, Suzuki A, Nakamura T (2007) Preparation of bisboron compounds controlling calcium concentration in cells.
42. Mikoshiba K, Nukina N, Ozaki S, Hamada K, Gota JL, et al. (2010) Protein crosslinking inhibitor.
43. Rondstvedt CS, Scriber RM, Wulffman CE (1995) Alcoholysis of triarylboranes. *J Org Chem* 20: 9-12.
44. Weidmann H, Zimmerman HK (1959) Borsäure-ester von N-substituierten Aminoalkoholen. *Justus Liebigs Ann Der Chemie* 619: 28-35.
45. Letsinger RL, Skoog IJ (1955) Organoboron Compounds. IV. Aminoethyl Diarylborinates. *J Am Chem Soc* 77: 2491-2494.
46. Povlock TP, Lippincott WT (1958) The Reaction of Trimethoxyboroxine with Aromatic Grignard Reagents. *J Am Chem Soc* 80: 5409-5411.
47. Brown HC, Colet TE (1983) A simple preparation of boronic esters from organolithium reagents and selected trialkoxyboranes. *Organometallics* 2: 1316-1319.
48. Mori Y, Kobayashi J, Manabe K, Kobayashi S (2002) Use of boron enolates in water. The first boron enolate-mediated diastereoselective aldol reactions using catalytic boron sources. *Tetrahedron* 58: 8263-8268.
49. Ozaki S, Suzuki AZ, Bauer PO, Ebisui E, Mikoshiba K (2013) 2-Aminoethyl diphenylborinate (2-APB) analogues: regulation of Ca^{2+} signaling. *Biochem Biophys Res Commun* 441: 286-290.
50. Ozaki S (2014) 2-Aminoethyl diphenylborinate (2-APB) analogues: Part 2. Regulators of Ca^{2+} release and consequent cellular processes. *Archives of Physiology* 1: 1-6.
51. Griffin M, Casadio R, Bergamini CM (2002) Transglutaminases: nature's biological glues. *Biochem J* 368: 377-396.
52. Lorand L, Graham RM (2003) Transglutaminases: crosslinking enzymes with pleiotropic functions. *Nat Rev Mol Cell Biol* 4: 140-156.
53. Selkoe DJ, Abraham C, Ihara Y (1982) Brain transglutaminase: in vitro crosslinking of human neurofilament proteins into insoluble polymers. *Proc Natl Acad Sci U S A* 79: 6070-6074.
54. Lorand L (1996) Neurodegenerative diseases and transglutaminase. *Proc Natl Acad Sci U S A* 93: 14310-14313.
55. Hoffner G, Djian P (2005) Transglutaminase and diseases of the central nervous system. *Front Biosci* 10: 3078-3092.
56. Malorni W, Farrace MG, Rodolfo C, Piacentini M (2008) Type 2 transglutaminase in neurodegenerative diseases: the mitochondrial connection. *Curr Pharm Des* 14: 278-288.
57. Norlund MA, Lee JM, Zainelli GM, Muma NA (1999) Elevated transglutaminase-induced bonds in PHF tau in Alzheimer's disease. *Brain Res* 851: 154-163.
58. Jeitner TM, Matson WR, Folk JE, Blass JP, Cooper AJ (2008) Increased levels of gamma-glutamylamines in Huntington disease CSF. *J Neurochem* 106: 37-44.
59. Karpuj MV, Becher MW, Springer JE, Chabas D, Youssef S, et al. (2002) Prolonged survival and decreased abnormal movements in transgenic model of Huntington disease, with administration of the transglutaminase inhibitor cystamine. *Nat Med* 8: 143-149.
60. Mastroberardino PG, Iannicola C, Nardacci R, Bernassola F, De Laurenzi V, et al. (2002) 'Tissue' transglutaminase ablation reduces neuronal death and prolongs survival in a mouse model of Huntington's disease. *Cell Death Differ* 9: 873-880.
61. Bauer PO, Hudec R, Ozaki S, Okuno M, Ebisui E, et al. (2011) Genetic ablation and chemical inhibition of IP3R1 reduce mutant huntingtin aggregation. *Biochem Biophys Res Commun* 416: 13-17.
62. Bauer PO, Hudec R, Goswami A, Kurosawa M, Matsumoto G, et al. (2012) ROCK-phosphorylated vimentin modifies mutant huntingtin aggregation via sequestration of IRBIT. *Mol Neurodegener* 7: 43.
63. Azari P, Rahim I, Clarkson DP (1981) Transglutaminase activity in normal and hereditary cataractous rat lens and its partial purification. *Curr Eye Res* 1: 463-469.
64. Schroeder WT, Thacher SM, Stewart-Galetka S, Annarella M, Chema D, et al. (1992) Type I keratinocyte transglutaminase: expression in human skin and psoriasis. *J Invest Dermatol* 99: 27-34.
65. Tatsukawa H, Fukaya Y, Frampton G, Martinez-Fuentes A, Suzuki K, et al. (2009) Role of transglutaminase 2 in liver injury via cross-linking and silencing of transcription factor Sp1. *Gastroenterology* 136: 1783-1795.
66. Lorand L, Rule NG, Ong HH, Furlanetto R, Jacobsen A, et al. (1968) Amine specificity in transpeptidation. Inhibition of fibrin cross-linking. *Biochemistry* 7: 1214-1223.
67. Stenberg P, Nilsson JL, Eriksson O, Lundén R (1971) Fibrin-stabilizing factor inhibitors. II. The inhibitory properties of various aliphatic amines. *Acta Pharm Suec* 8: 415-422.
68. Fésüs L (1982) Transglutaminase activation: significance with respect to immunologic phenomena. *Surv Immunol Res* 1: 297-304.
69. Piper JL, Gray GM, Khosla C (2002) High selectivity of human tissue transglutaminase for immunoreactive gliadin peptides: implications for celiac sprue. *Biochemistry* 41: 386-393.
70. Hovhannisyan Z, Weiss A, Martin A, Wiesner M, Tollefsen S, et al. (2008) The role of HLA-DQ8 beta57 polymorphism in the anti-gluten T-cell response in coeliac disease. *Nature* 456: 534-538.
71. Watts RE, Siegel M, Khosla C (2006) Structure-activity relationship analysis of the selective inhibition of transglutaminase 2 by dihydroisoxazoles. *J Med Chem* 49: 7493-7501.
72. Pardin C, Pelletier JN, Lubell WD, Keillor JW (2008) Cinnamoyl inhibitors of tissue transglutaminase. *J Org Chem* 73: 5766-5775.
73. Duval E, Case A, Stein RL, Cuny GD (2005) Structure-activity relationship study of novel tissue transglutaminase inhibitors. *Bioorg Med Chem Lett* 15: 1885-1889.

74. Choi K, Siegel M, Piper JL, Yuan L, Cho E, et al. (2005) Chemistry and biology of dihydroisoxazole derivatives: selective inhibitors of human transglutaminase 2. *Chem Biol* 12: 469-475.
75. Ozaki S, Ebisui E, Hamada K, Goto J, Suzuki AZ, et al. (2010) Potent transglutaminase inhibitors, aryl beta-aminoethyl ketones. *Bioorg Med Chem Lett* 20: 1141-1144.
76. Ozaki S, Ebisui E, Hamada K, Suzuki AZ, Terauchi A, et al. (2011) Potent transglutaminase inhibitors, dithio 1²-aminoethyl ketones. *Bioorg Med Chem Lett* 21: 377-379.
77. Berridge MJ (1998) Neuronal calcium signaling. *Neuron* 21: 13-26.
78. Berridge MJ (2010) Calcium hypothesis of Alzheimer's disease. *Pflugers Arch* 459: 441-449.
79. Berridge MJ (2013) Dysregulation of neural calcium signaling in Alzheimer disease, bipolar disorder and schizophrenia. *Prion* 7: 2-13.

Citation: Ozaki S (2014) 2-Aminoethyl Diphenylborinate (2-APB) Analogues: Part 3 - Regulators of Huntington Aggregation and Transglutaminase. J Bioengineer & Biomedical Sci 4: 131. doi:[10.4172/2155-9538.1000131](https://doi.org/10.4172/2155-9538.1000131)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 350 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>

