

Role of Terahertz Radiation on Optical Properties of Laser Pulse in a Double Coupled Quantum Well Nanostructure

Shiri J^{1*} and Malakzadeh A²

¹Young Researchers and Elite Club, North Tehran Branch, Islamic Azad University, Tehran, Iran

²Centre for Laser and Optics, Basic Sciences Department, Imam Hussein University, Tehran, Iran

Abstract

The transient and steady-state behavior of the absorption and the dispersion of a probe pulse laser field propagating through an InGaAs/InP double coupled quantum well are studied. The effect of terahertz radiation excitation, electron tunnelling and incoherent pumping on the optical properties of the probe field is discussed. In the terahertz (30~300 μm or 1~10 THz) intersubband transition, the incoming photon energy is (4~41 meV) and maybe in the order of electron thermal broadening ($KT \sim 6 \text{ meV} \sim 25 \text{ meV}$ for 77 K~300 K). Therefore in the conventional structure, the incoming photon can directly excite the ground state electrons to higher energy levels and this process inhabits the correct optical switching in terahertz applications. We show that the group velocity of a light pulse can be controlled from superluminal to subluminal or vice versa by controlling the rates of incoherent pumping field and tunnelling between the quantum wells. The required switching time is calculated and we find it between 3 to 26 picoseconds.

Keywords: Electro-optical switching; Dispersion and absorption; Group velocity; Terahertz radiation; Tunnelling effects; Incoherent pumping field

Introduction

It is known that the absorption and the dispersion properties of a weak probe field can be modified effectively by atomic coherence and quantum interference [1-6]. Atomic coherence can be achieved by the strong coupling fields, the spontaneous emission and incoherent pumping fields. It is known that atomic coherence due to the coherent laser field has essential roles for modifying the optical properties of atomic systems such as spontaneously generated coherence (SGC) [2], lasing without inversion [3], modifying spontaneous emission [4], coherent population trapping (CPT) [6], optical bistability [7-11] and so on [12-17]. Furthermore, it has been shown that quantum interference arising from SGC [8] and incoherent pumping field [18] can be used for analyse of some interesting phenomena such as lasing without population inversion [5], optical bistability [18], and superluminal/subluminal light propagation [19]. Similar phenomena involving quantum coherence in solid state systems such as semiconductor quantum wells (QWs) and quantum dots (QDs) [20], can also be occurred [21,22]. In the past decade, there has been an increasing interest in optical properties of quantum dot molecules (QDMs) and quantum wells (QWs), due to important role in optoelectronic devices. Recently, investigators have examined the effects of an external field and inter-dot tunnel coupling on the optical properties of QDs and QWs [23-29]. Quantum well semiconductors were chosen because of their advantage in flexible design, controllable interference strength, long dephasing times [30,31] large dephasing rates [$\sim 10\text{ps}^{-1}$] [32] and large electric dipole moment which make them suitable for application in the optoelectronic devices. Quantum coherence in a QW structure can be induced by electron tunnelling or applying a laser field [33,34]. Coherence induced by incoherent field and tunnel coupling in the QW system plays an important role in light-matter interaction and has found numerous implementations in semiconductor optics. On the other hand, coherent control [35-38] over the dispersive and absorptive properties of solid-state media such as photonic crystals and semiconductors has recently attracted a lot of attention [39-43]. Several proposals for quantum coherence and interference in QWs have been performed and analyzed. To utilize the tunnelling effect, an

electron is excited by a laser field, then tunnels to the second QW by controlling the external voltage between the wells [44]. An interesting application of QWs is modification of light pulse to make a fast electro-optical switch by controlling the propagation of a weak light pulse in a semiconductor system, which depends on the dispersive properties of the medium.

In this paper, we introduce a compact four level quantum wells system composed of two QWs. Then, we investigate effect of terahertz radiation, incoherent pumping field and tunnelling between QWs on the absorption, dispersion and the group velocity of a weak probe field. The required switching time when propagation of light changes from subluminal to superluminal and vice versa is also discussed. We find that the dispersion/absorption spectra of the probe pulse can be changed via the effect of terahertz radiation, incoherent pumping field and tunnelling effect.

Model and Equations of Motion

In Figure 1, we consider a compact double coupled quantum well nanostructure which is fabricated using InGaAs/InP nanostructures in material grown by an attractive growth technique i.e., organometallic vapor phase epitaxy (OMVPE). The QWs consist of two periods of alternating InGaAs and InP layers. An incoherent pumping field and weak probe field are applied to first quantum well (QW_1). The quantum well is designed so that its width is narrower in comparison to the other quantum well (QW_2) and provides larger energy difference between the levels. For controlling of the tunnelling rate between QWs, the system is placed between two connected electrodes, as electrodes

*Corresponding author: Shiri J, Young Researchers and Elite Club, North Tehran Branch, Islamic Azad University, Tehran, Iran, Tel: +98 21 7731 8896; E-mail: jalil.shiri@chmail.ir

Received April 23, 2017; Accepted May 15, 2017; Published June 20, 2017

Citation: Shiri J, Malakzadeh A (2017) Role of Terahertz Radiation on Optical Properties of Laser Pulse in a Double Coupled Quantum Well Nanostructure. J Laser Opt Photonics 4: 154. doi: 10.4172/2469-410X.1000154

Copyright: © 2017 Shiri J, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

are in contact to this system. By applying independently tunable gate voltages, electron tunnelling can easily be accomplished between QW_1 and QW_2 . Figure 1 shows the detailed band structure and energy levels of the system. Lower level $|0\rangle$ and upper level $|1\rangle$ are conducting band levels of QW_1 . Level $|2\rangle$ and level $|3\rangle$ are the excited conducting levels of the QW_2 of the right of QW_1 . It is assumed that the energy difference of three excited levels and the lower level is large, so their tunneling couplings can be ignored. By applying a gate voltage the level $|2\rangle$ and the level $|3\rangle$ get closer to the level $|1\rangle$. A weak tunable probe field of the frequency ω_p with Rabi frequency $\Omega_p = E \cdot \vec{\phi} / 2\hbar$ and an incoherent pumping field Λ are applied to the transition $|0\rangle \rightarrow |1\rangle$. Here, $\vec{\phi}$ is electric dipole moment and E is amplitude of the probe field laser. By forming the resonant coupling of the probe field with the QW_1 , an electron is excited from the $|0\rangle$ band to the $|1\rangle$ band of the QW_1 . By providing the tunnelling conditions the electron can be transferred to level $|2\rangle$ in QW_2 and the terahertz radiation of frequency ω_{THz} with Rabi frequency Ω_{THz} prompt the electron from the level $|2\rangle$ to the level $|3\rangle$. The total Hamiltonian in the rotating-wave approximation method [45,46] which represents the interaction of the probe laser field, terahertz radiation and incoherent pumping field with the double coupled QWs system, can be expressed in the form of

$$H = \sum_{j=0}^5 E_j |j\rangle\langle j| + \left[\Omega_p e^{-i\omega_p t} |0\rangle\langle 1| + \phi \varepsilon |0\rangle\langle 1| + T_{12} |1\rangle\langle 2| + \Omega_{THz} e^{-i\omega_{THz} t} |2\rangle\langle 3| + H.C. \right] \quad (1)$$

Where $E_j = \hbar\omega_j$ denotes the energy of state $|i\rangle$. ϕ is the dipole moment of the atomic transition corresponding to the pumping of the electrons from level $|0\rangle$ to level $|1\rangle$, and the electric field \mathcal{E} implies the electrical amplitude of the incoherent pumping field. T_{12} correspond to tunnelling between “ QW_1 ” and “ QW_2 ”. The tunnelling can be described by perturbation theory which can be given by Bardeen’s approach [47]. The density-matrix approach given by

$$\frac{\partial \rho}{\partial t} = -\frac{i}{\hbar} [H, \rho], \quad (2)$$

Can be used to obtaining the density operator in an arbitrary multilevel QWs system. Substituting eqn. (1) in eqn. (2), the density matrix equations of motion can be expressed as

$$\begin{aligned} \dot{\rho}_{01} &= (i\delta_p - \Gamma_{01} - \Lambda)\rho_{01} + iT_{12}\rho_{02} - i\Omega_p(\rho_{00} - \rho_{11}), \\ \dot{\rho}_{02} &= iT_{12}\rho_{01} + (i(\delta_p + \omega_{12}) - \Gamma_{02} - \Lambda/2)\rho_{02} + i\Omega_{THz}\rho_{03} + i\Omega_p\rho_{12}, \end{aligned}$$




Figure 1: Schematic of system, which shows the detailed band structure and quantized energy levels for proposed double coupled QW.

$$\begin{aligned} \dot{\rho}_{03} &= i\Omega_{THz}\rho_{02} + (i(\delta_p + \omega_{12} + \omega_{23}) - \Gamma_{03} - \Lambda/2)\rho_{03} + i\Omega_p\rho_{13}, \\ \dot{\rho}_{12} &= i\Omega_p\rho_{02} - (i\omega_{12} + \Gamma_{12} + \Lambda/2)\rho_{12} + i\Omega_{THz}\rho_{13} + iT_{12}(\rho_{11} - \rho_{22}), \\ \dot{\rho}_{13} &= i\Omega_p\rho_{03} + i\Omega_{THz}\rho_{12} - (i(\omega_{12} + \omega_{23}) + \Gamma_{13} + \Lambda/2)\rho_{13} - iT_{12}\rho_{23}, \\ \dot{\rho}_{23} &= -iT_{12}\rho_{13} - i(\omega_{23} + \Omega_{THz})\rho_{23} + i\Omega_{THz}(\rho_{22} - \rho_{33}), \\ \dot{\rho}_{00} &= -i\Omega_p(\rho_{01} - \rho_{10}) - \Lambda\rho_{00} + (\gamma_{10} + \Lambda)\rho_{11} + \gamma_{20}\rho_{22} + \gamma_{30}\rho_{33}, \\ \dot{\rho}_{11} &= i\Omega_p(\rho_{01} - \rho_{10}) + \Lambda\rho_{00} - (\gamma_{10} + \Lambda)\rho_{11} + iT_{12}(\rho_{12} - \rho_{21}), \\ \dot{\rho}_{22} &= -iT_{12}(\rho_{12} - \rho_{21}) + i\Omega_{THz}(\rho_{23} - \rho_{32}) - \gamma_{20}\rho_{22}, \\ \dot{\rho}_{33} &= -i\Omega_{THz}(\rho_{23} - \rho_{32}) - \gamma_{30}\rho_{33}, \end{aligned} \quad (3)$$

Where $\rho_{mn} = |m\rangle\langle n|$ ($m, n = 0, 1, 2, 3$) and $\rho_{mn} = |m\rangle\langle m|$ ($m = 0, 1, 2, 3$) represent the coherent terms and the population operators for the QWs, respectively. We get $\omega_{12} = \omega_{10} - \omega_{20}$ and $\omega_{23} = \omega_{20} - \omega_{30}$. The probe field detuning with respect to the QW transition frequencies is $\delta_p = \omega_{10} - \omega_p$. The term $\Lambda = 2(\phi^2 / \hbar^2) \Gamma_p$ is the incoherent pumping rate. The spontaneous emission rates for sub-band $|i\rangle$, denoted by γ_{10} , are due primarily to longitudinal optical (LO) phonon emission events at low temperature. The total decay rates Γ_{ij} ($i \neq j$) are given by $\Gamma_{0n} = \gamma_{n0} / 2 + \gamma_{n0}^{dph}$, $\Gamma_{mn} = (\gamma_{n0} + \gamma_{m0}) / 2 + \gamma_{mn}^{dph}$, $m, n = 1, 2, 3$ and $m \neq n$, here $|i\rangle\langle i|$, are the dephasing rates of the quantum coherence of the $|i\rangle \leftrightarrow |j\rangle$ pathway and determined by electron–electron, interface roughness, and phonon scattering processes. Usually, γ_{mn}^{dph} is the dominant mechanism in a semiconductor solid-state system. Eqn. (3) can be solved to obtain the steady state response of the medium. The susceptibility of the compact double QWs to the weak probe field is determined by coherence term ρ_{01}

$$\chi = \frac{2N\phi}{E\varepsilon_0} \rho_{01}, \quad (4)$$

Where N is the carrier density in the proposed QWs system. Susceptibility comprise two parts, real and imaginary ($\chi = \chi' + i\chi''$). Note that the real part of the susceptibility χ' correspond to the dispersion and imaginary part χ'' correspond to absorption. The dispersion slope of the probe field has a major role in the group velocity. The group velocity v_g of the light, which is propagates in the medium, given by Bardeen [48]:

$$v_g = \frac{c}{1 + 2\pi\chi'(\omega_p) + 2\pi\omega_p(\partial\chi'(\omega_p) / \partial\omega_p)}, \quad (5)$$

Eqn. (5) implies that for a negligible real part of susceptibility, the light propagation can be superluminal as a negative slope of dispersion, on the other hand, for positive dispersion slope, the light propagation in the medium can be subluminal.

Results and Discussion

Now, we analyze the numerical results of the above equations and discuss the transient and the steady-state behavior of the absorption and the dispersion. It is assumed that the system is initially in the ground state, i.e., $\tilde{\rho}_{00}(0) = 1$ and $\tilde{\rho}_{ij}(0) = 0$ ($i, j = 0, 1, 2, 3$). We take typically slow dephasing rates $\Gamma_{01} = \gamma = 1$ THz and spontaneous emission $\gamma_{10} = 0.6$ THz and other relevant parameters by the factor of these rates. Introduced rates are equivalent to dephasing times in the order of picoseconds. Here the according to the equation (5), the positive and negative dispersion slope are representing the propagation

of light subluminal and superluminal respectively.

Figure 2(a) shows the dispersion (dashed) and absorption (solid) properties of a probe field vs. probe field detuning δ_p , in the absence of incoherent pumping field Λ and tunnelling effect. We observe that the absorption peak accompanies by a negative dispersion. Thus, superluminal light with large absorption propagates through the medium. In Figure 2(b), we show at the steady state behavior of the probe dispersion (dashed) and absorption (solid) in the presence of the first tunnelling effect T_{12} and absence of an incoherent pumping field. It can easily be seen that the absorption of the probe field reduced around zeros detuning probe field just by applying tunnelling rate $T_{12} = 1\gamma$. We find that the slope of dispersion is very sensitive to the tunnelling effect. When we increase T_{12} from 0 to 1γ , slope of dispersion changes from negative to positive. Figure 2(c) shows the effect of the terahertz radiation on the system, when keeping all other parameters fixed in Figure 2(b), and with applying the terahertz signal, the dispersion slope does not change, but tow windows transparency created that is accompanied by three absorption peaks.

In Figure 3, we investigate the incoherent pumping field effect on the system. In Figure 2(a), we applied an incoherent pumping field to condition of Figure 2(a) with values $\Lambda = 8\gamma$. By increasing the incoherent pumping field Λ from 0 to 8γ , the absorption peak of the probe field becomes broaden, while the slope of the dispersion is still negative around the probe field detuning. Physically, by increasing the incoherent pump rate, the upper levels are populated, for this

condition the probe absorption will reduce in transition $|0\rangle \rightarrow |1\rangle$, thus the peak maximum of the probe field absorption gets reduce. In Figure 3(b), by applying the tunnelling rate T_{12} mid incoherent pumping field, one absorption peak of the probe field is crated around zeros detuning probe field. By applying the incoherent pumping field, when keeping all other parameters fixed in Figure 2(b), the slope of dispersion changes from positive to negative that is shown in Figure 2(b) and 3(b). This indicates that group velocity of light propagates, increasing from subluminal to superluminal through the medium. In Figure 3(c), we simultaneously apply the tunnelling rate T_{12} between QW_1 and QW_2 and the terahertz signal, while the incoherent pump rate is still $\Lambda=8\gamma$. We observe that the probe field absorption convert to probe gain, which is accompanied by positive dispersion slope i.e. the probe field propagation changes from superluminal to subluminal in medium by applying the incoherent field as shown in Figures 2(c) and 3(c). This is an important mechanism in which the probe field absorption and dispersion can be controlled by the incoherent pump field in the quantum well.

Now, we are interested in the dynamical behaviour of the dispersion slope due to this properties can be used as an electro-optical switch of group velocity from subluminal to superluminal or vice versa. We are interested in the required switching time for change of the light propagation from subluminal to superluminal or vice versa. Switching time for subluminal/superluminal light propagation is defined as the time to reach a steady-state from the superluminal state to the subluminal state and vice versa. Figure 4(a) shows the transition






Figure 2: Real (dashed) and imaginary (solid) parts of susceptibility as a probe field wavelength for (a) $T_{12}=0$, TH_2 signal=off, $\Lambda=0$, (b) $T_{12}=1\gamma$, TH_2 signal=of, $\Lambda=0$, (c), $T_{12}=1\gamma$, $\Lambda=0$, TH_2 signal=on Other parameters are, $\omega_{12}=\omega_{23}=0.01\gamma$, $\Gamma_{01}=\gamma=1TH_2$, $\Gamma_{02}=0.1\Gamma_{01}$, $\Gamma_{03}=0.01\Gamma_{01}$, $\Gamma_{12}=0.05\Gamma_{01}$, $\Gamma_{13}=0.025\Gamma_{01}$, $\Gamma_{23}=0.05\Gamma_{01}$, $\gamma_{10}=0.6TH_2$, $\gamma_{20}=0.1\gamma_{10}$, $\gamma_{30}=0.1\gamma_{10}$, $\Omega_p=0.01\gamma$.



behaviour of the dispersion slope by consecutively switching the tunnelling rate from $T_{12} = 0$ to $T_{12} = 1\gamma$ and vice versa. By increasing the normalized time $t\gamma$ ($\gamma = \Gamma_{10} = 1THz$), the dispersion slope takes a steady negative values for $T_{12} = 0$ corresponding to superluminal light propagation, while it changed to negative on applying the terahertz signal which is corresponding to subluminal light propagation. In Figure 4(b), we plot the switching diagram of the dispersion slope for

two different values of T_{12} . The required switching time for propagation of the light from subluminal to superluminal is about 8ps and from superluminal to subluminal is about 12ps. Now, we investigate the effect of the incoherent pumping rate by consecutively switching incoherent pumping rate from $\Lambda = 0$ to $\Lambda = 8\gamma$ and vice versa, while the tunnelling rate $\Lambda = 0$ is fixed (Figure 5(a)). It is clearly find that the slope of the dispersion changes from positive to negative by adjusting




Figure 5: Dynamical behaviour (a) and switching process (b) of dispersion slope for $T_{12}=1\gamma$, TH_z signal=off, $\Lambda=(0, 8\gamma)$. Other parameters are as in Figure 2.




Figure 6: Dynamical behaviour (a) and switching process (b) of absorption for TH_z signal (off, on) $T_{12}=1\gamma$, $\Lambda=0$. Other parameters are as in Figure 2.

the incoherent field. For $\Lambda = 0$ the slope of the dispersion is positive corresponding to subluminal light propagation, while it changes to negative as the incoherent pumping changes from $\Lambda = 0$ to $\Lambda = 8\gamma$ corresponding to superluminal light propagation. The required switching time for change of the propagation light from subluminal to superluminal is about 3ps, and vice versa is 15ps (Figure 5(b)). This approach can be utilized to produce a switch operating only by controlling the tunnelling rate T_{12} and incoherent rate Λ . The transient behaviour of the probe absorption is displayed for two various rate of terahertz signal, while keeping $T_{12} = 1\gamma$ and $\Lambda = 8\gamma$ fixed in Figure 6. We are looking for the required switching time for changing the light absorption case to the probe gain or vice versa by proper manipulating the tunnelling rate T_{12} and terahertz signal. Effect of the terahertz signal leads to a significant probe amplification of steady state probe absorption. Figure 6(a,b) shows that the required switching time from large absorption to probe gain case or vice versa is about 20ps.

Conclusion

We investigated the transient and the steady-state behavior of a weak probe field in a compact double coupled QWs system with applying the tunnelling between QWs, terahertz signal and one incoherent pumping field. It is shown that the absorption and the dispersion of the probe field can be controlled by applying the tunnelling between QWs, terahertz radiation and incoherent pumping fields. It has also

been shown that the medium can be used as an optical switch in which the propagation of the laser pulse can be controlled with tunnelling between QWs and the incoherent pumping field. We obtained the switching time; between 3 to 26 picosecond as a high speed optical switch is an important technique for quantum information, networks and communication.

References

1. Etteh NEI, Harrison P (2001) Carrier scattering approach to the origins of dark current in mid- and far-infrared (terahertz) quantum-well intersubband photodetectors (QWLPs) IEEE J Quant Elec 37: 672-675.
2. Macovei MA, Evers J (2004) Phase dependence of collective fluorescence via interferences from incoherent pumping. Opt Com 240: 379-384.
3. Javanainen J (1992) Effect of State Superpositions Created by Spontaneous Emission on Laser-Driven Transitions. Europhy Let (EPL) 17: 407-412.
4. Hu X, Zhang JP (2004) Coherent population trapping via quantum interference of incoherent pump processes in three-level systems. J Phy B: Atomic, Mol Opt Phy 37: 345-356.
5. Fleischhauer M, Keitel CH, Scully MO, Su C (1992) Lasing without inversion and enhancement of the index of refraction via interference of incoherent pump processes. Opt Com 87: 109-114.
6. Bullock D, Evers J, Keitel CH (2003) Modifying spontaneous emission via interferences from incoherent pump fields. Phy Lett A 307: 8-12.
7. Wu J, Lü XY, Zheng LL (2010) Controllable optical bistability and multistability in a double two-level atomic system. J Phys B: Atom, Mol Opt Phy 43: 161003.

8. Wang Z, Xu M (2009) Control of the switch between optical multistability and bistability in three-level V-type atoms. *Opt Com* 282: 1574-1578.
9. Osman KI, Joshi A (2013) Induced coherence and optical bistability in a four-level system with incoherent pumping. *Opt Com* 293: 86-94.
10. Joshi A, Yang W, Xiao M (2003) Effect of quantum interference on optical bistability in the three-level V-type atomic system. *Phy Rev A* 68: 015806.
11. Hossein Asadpour S, Eslami-Majd A (2012) Controlling the optical bistability and transmission coefficient in a four-level atomic medium *J Lumi* 132: 1477-14782.
12. Antón MA, Carreño F, Calderón OG, Melle S (2008) Tunable all-optical bistability in a semiconductor quantum dot damped by a phase-dependent reservoir. *Opt Com* 281: 3301-3313.
13. Chen A (2014) Coherent manipulation of spontaneous emission spectra in coupled semiconductor quantum well structures. *Opt Exp* 22: 26991.
14. Si LG, Yang WX, Yang X (2009) Ultraslow temporal vector optical solitons in a cold four-level tripod atomic system. *J Opt Soc Am B* 26: 478.
15. Yang WX, Chen AX, Lee RK, Wu Y (2011) Matched slow optical soliton pairs via biexciton coherence in quantum dots. *Phys Rev A* 84: 013835.
16. Yang WX, Hou JM, Lin Y, Lee RK (2009) Detuning management of optical solitons in coupled quantum wells. *Phys Rev A* 79: 033825.
17. Yannopapas V, Paspalakis E, Vitanov NV (2009) Plasmon-Induced Enhancement of Quantum Interference near Metallic Nanostructures. *Phys Rev Lett* 103: 063602.
18. Mahmoudi M, Sahrai M, Tajalli H (2006) Subluminal and superluminal light propagation via interference of incoherent pump fields. *Phys Lett A* 357: 66-71.
19. Sahrai M, Asadpour SH, Mahrami H, Sadighi-Bonabi R (2011) Controlling the optical bistability via quantum interference in a four-level N-type atomic system. *J Lumi* 131: 1682-1686.
20. Shiri J (2016) Propagation of a laser pulse and electro-optic switch in a GaAs/AlGaAs quadruple-coupled quantum dot molecule nanostructure. *Laser Physics* 26: 056202.
21. Su H, Chuang SL (2006) Room-temperature slow light with semiconductor quantum-dot devices. *Opt Lett* 31: 271.
22. Palinginis P, Sedgwick F, Crankshaw S, Moewe M, Chang-Hasnain CJ (2005) Room temperature slow light in a quantum-well waveguide via coherent population oscillation. *Opt Exp* 13: 9909.
23. Schmidt H, Nikonov DE, Campman KL, Gossard AC, Lu AI (1999) Quantum Interference in Semiconductor Quantum Wells. *Laser Physics* 9: 797-812.
24. Chang-Hasnain CJ, Pei-Cheng Ku, Jungho Kim, Shun-Lien Chuang (2003) Variable optical buffer using slow light in semiconductor nanostructures. *IEEE* 9: 1884-1897.
25. Che C, Han Q, Ma J, Zhou Y, Yu S, Tan L (2012) Electron radiation effects on InAs/GaAs quantum dot lasers. *Laser Physics* 22: 1317-1320.
26. Kosionis SG, Terzis AF, Paspalakis E (2007) Optimal control of a symmetric double quantum-dot nanostructure: Analytical results. *Phys Rev B* 75: 193305.
27. Li J, Liu J, Yang X (2008) Controllable gain, absorption and dispersion properties of an asymmetric double quantum dot nanostructure. *Superlattices and Microstructures* 44: 166-172.
28. Tsukanov AV, Openov LA (2004) Electron transfer between semiconductor quantum dots via laser-induced resonance transitions. *Semicond* 38: 91-98.
29. Wang Z (2011) Control of the probe absorption via incoherent pumping fields in asymmetric semiconductor quantum wells. *Ann Phys* 326: 340-349.
30. Joshi A, Xiao M (2004) Optical bistability in a three-level semiconductor quantum-well system. *AppPhys B* 79: 65-9.
31. Wijewardane HO, Ullrich CA (2004) Coherent control of intersubband optical bistability in quantum wells. *App Phys Lett* 84: 3984.
32. Serapiglia GB, Paspalakis E, Sirtori C, Vodopyanov KL, Phillips CC (2000) Laser-Induced Quantum Coherence in a Semiconductor Quantum Well. *Phys Rev Lett* 84: 1019-1022.
33. Sahrai M, Mahmoudi M, Kheradmand R (2007) The impact of the relative phase on the transient optical properties of a four-level EIT medium. *Phys Lett A* 367: 408-414.
34. Dynes JF, Frogley MD, Rodger J, Phillips CC (2005) optically mediated coherent population trapping in asymmetric semiconductor quantum wells. *Phys Rev B* 72: 085323.
35. Kash MM, Sautenkov VA, Zibrov AS, Hollberg L, Welch GR, et al. (1999) Ultraslow Group Velocity and Enhanced Nonlinear Optical Effects in a Coherently Driven Hot Atomic Gas. *Phys Rev Lett* 82: 5229-5232.
36. Joshi A, Xiao M (2004) Optical bistability in a three-level semiconductor quantum-well system. *Appl Phys B* 79: 65-69.
37. Hau LV, Harris SE, Dutton Z, Behroozi CH (1999) No Title *Nature* 397: 594-598.
38. Ghulghazaryan RG, Malakyan YP (2003) Superluminal optical pulse propagation in nonlinear coherent media. *Phys Rev A* 67: 063806.
39. Borri P, Langbein W, Schneider S, Woggon U, Sellin RL, et al. (2002) Rabi oscillations in the excitonic ground-state transition of InGaAs quantum dots. *Phys Rev B* 66: 081306.
40. Cole BE, Williams JB, King BT, Sherwin MS, Stanley CR (2001) No Title *Nature* 410: 60-63.
41. Michler P, Kiraz A, Becher C, Schoenfeld WV, Petroff PM, et al. (2000) A Quantum Dot Single-Photon Turnstile Device *Science*. 290: 2282-2285.
42. Pelton M, Santori C, Vučković J, Zhang B, Solomon GS, et al. (2002) Efficient Source of Single Photons: A Single Quantum Dot in a Micropost Microcavity. *Phys Rev Lett* 89: 233602.
43. Yamamoto Y, Kim J, Benson O and Kan H (1999) No Title *Nature* 397: 500-503.
44. Ginzburg P, Orenstein M (2006) Slow light and voltage control of group velocity in resonantly coupled quantum wells. *Opt Exp* 14: 12467.
45. Mehmannaavaz MR, Nasehi R, Sattari H, Mahmoudi M (2014) Controlling the optical bistability and multistability via tunneling-induced and incoherent pumping field in a triple coupled quantum dots at a wavelength of $\lambda=1.550 \mu\text{m}$. *Superlatt and Microstruc* 75: 27-38.
46. Kapale KT, Scully MO, Zhu SY, Zubairy MS (2003) Quenching of spontaneous emission through interference of incoherent pump processes. *Phys Rev A* 67: 023804.
47. Gardiner CW, Collett MJ (1985) Input and output in damped quantum systems: Quantum stochastic differential equations and the master equation. *Phys Rev A* 31: 3761-3774.
48. Bardeen J (1961) Tunnelling from a Many-Particle Point of View. *Phys Rev Lett* 6: 57-59.

Citation: Shiri J, Malakzadeh A (2017) Role of Terahertz Radiation on Optical Properties of Laser Pulse in a Double Coupled Quantum Well Nanostructure. *J Laser Opt Photonics* 4: 154. doi: [10.4172/2469-410X.1000154](https://doi.org/10.4172/2469-410X.1000154)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submit>