

Sacral Malformations - Implications in Sacral Nerve Stimulation

Ana Povo*

Department of Anatomy, Faculty of Medicine of University of Porto, Portugal

*Corresponding author: Povo A, Department of Anatomy, Faculty of Medicine of University of Porto, Portugal, Tel: +351914296134; Fax: +351225513617; E-mail: anapovo@sapo.pt

Rec date: June 8, 2016; Acc date: June 17, 2016; Pub date: June 23, 2016

Copyright: © 2016 Povo A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Povo A (2016) Sacral Malformations - Implications in Sacral Nerve Stimulation. *Reprod Syst Sex Disord* 5: 175. doi:[10.4172/2161-038X.1000175](https://doi.org/10.4172/2161-038X.1000175)

To Editor,

Sacral Nerve Stimulation (SNS) is a common treatment for various pelvic floor disorders, such as urinary and faecal incontinence and constipation [1-5].

This technique consists of the introduction of electrodes by a percutaneous approach through the posterior sacral foramina for placement in the proximity of the target sacral spinal nerve for therapeutic low frequency stimulation [1-5].

One of the most important steps of this technique is the correct topographic anatomic identification of the posterior sacral foramina on the body surface of the sacrum [6,7]. Although the sacrum's anatomy is recognized to be highly variable. In the literature there are several studies regarding the variable morphology of the sacrum that also shows high incidence of sacral malformations, ranging between 10% and 58% [8]. A variety of the anatomic malformations are clinically imperceptible. SNS can be difficult or impossible to be performed in these patients with an unknown or undetected malformation of the sacrum [1,3,4].

Recently we reviewed 998 consecutive MRI scans performed to investigate low back pain in patients who had undergone CT and/or X-ray. Sacral malformations were found in almost one quarter of our 998 cases and may represent an under-reported cause of inadequate (or impossible) electrode placement. We therefore recommend a sacrum X-ray before SNS. In all patients with minimal or major symptoms of unknown cause, such as lower back pain, perianal or sciatic pain or radiculopathy, an MRI before SNS may be helpful to exclude sacral malformations and/or guide the surgical procedure. In all patients in whom appropriate foramina placement is difficult or impossible, we recommend a CT scan or MRI to exclude sacral malformation [8].

We believe that, in the future, a three-dimensional anatomical model based on radiologic studies of the sacral malformations will ease the navigation and placement of electrodes in a variety of conditions that are currently contraindications for SNS [8].

References

1. Matzel KE (2007) Sacral Nerve Stimulation. In: Ratto C, Doglietto GB (eds.) *Fecal Incontinence: diagnosis and treatment*, pp: 211-217.
2. Wexner SD, Coller JA, Devroede G, Hull T, McCallum R, et al. (2012) Sacral Nerve Stimulation for Fecal Incontinence – Results of a 120-Patient Prospective Multicenter Study. *Ann Surg* 251: 441-449.
3. Dudding TC, Hollingshead JR, Nicholls RJ, Vaizey CJ (2011) Sacral nerve stimulation for faecal incontinence: patient selection, service provision and operative technique. *Colorectal Dis* 13: e187-195.
4. Matzel KE, Kamm MA, Stösser M, Baeten CG, Christiansen J, et al. (2004) Sacral spinal nerve stimulation for faecal incontinence: multicentre study. *Lancet* 363: 1270-1276.
5. Madoff RD, Laurberg S, Leher P, Matzel KE, Mellgren AF, et al. (2013) Surgery for Faecal Incontinence. In Abrams, L Cardozo, S Khoury, A Wein (eds.) *INCONTINENCE*, 5th Edition. ICUD-EAU.
6. Hasan TS, Shanahan DA, Pridie AK, Neal DE (1996) Surface localization of sacral foramina for neuromodulation of bladder function- an anatomical study. *Eur Urol* 29: 90-98.
7. Prapasivorakul S, Gorissen KJ, Gosselink MP, Curran K, Jones OM, et al. (2014) Temporary sacral neuromodulation under local anaesthesia using new anatomical reference points. *Tech Coloproctol* 18: 1093-1097.
8. Povo A, Arantes M, Matzel KE, Barbosa J, Ferreira MA (2016) Sacral malformations: use of imaging to optimise sacral nerve stimulation. *Int J Colorectal Dis* 31: 351-357.