

Santilli's Prime Chains: $P_j + 1 = aP_j \pm b$

Chun-Xuan Jiang*

Institute for Basic Research, Palm Harbor, P.O. Box 3924, Beijing 100854, P.R. China

Abstract

Santilli's prime chains: $P_j + 1 = aP_j \pm b, j=1, \dots, k-1, (a,b)=1, 2|ab$. If $a-1 = P_1^{\lambda_1} \dots P_n^{\lambda_n}, P_1 \dots P_n | b$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$. There exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k . It is the Book proof. This is a generalized Euclid-Euler proof for the existence of infinitely many primes. Therefore Euclid-Euler-Jiang theorem in the distribution of primes is advanced. It is the Book theorem.

Keywords: Generalized; Arithmetic; Prime; Euclid; Arbitrary; Integer

Introduction

A new branch of number theory: Santilli's additive isoprime theory is introduced. By using the arithmetic function $J_n(\omega)$ the following prime theorems have been proved.

It is the Book proof. [1-10].

1. There exist infinitely many twin primes.
2. The Goldbach's theorem. Every even number greater than 4 is the sum of two odd primes.
3. There exist finitely many Mersenne primes, that is, primes of the form $2^p - 1$ where P is prime.
4. There exist finitely many Fermat primes, that is, primes of the form $2^{2^n} + 1$.
5. There exist finitely many repunit primes whose digits (in base 10) are all ones.
6. There exist infinitely many primes of the forms: $x^2+1, x^4+1, x^8+1, x^{16}+1, x^{32}+1, x^{64}+1$.
7. There exist infinitely many primes of the forms: $x^2+b, x^3+2, x^5+2, x^7+2$
8. There exist infinitely many prime m -chains, $P_{j+1} = mP_j \pm (m-1), m=2,3, \dots$, including Cunningham chains.
9. There exist infinitely many triplets of consecutive integers, each being the product of k distinct primes. (Here is an example: $1727913=3 \times 11 \times 52361, 1727914=2 \times 17 \times 50821, 1727915=5 \times 7 \times 49369$.)
10. There exist infinitely many k -tuples of consecutive integers, each being the product of m primes, where $k>3, m>2$.
11. Every integer m may be written in infinitely many ways in the form.
$$m = \frac{P_2 + 1}{P_1^k - 1}$$
Where $k=1,2,3, \dots, P_1$ and P_2 are primes.
12. There exist infinitely many Carmichael numbers, which are the product of three primes, four primes, and five primes.
13. There exist infinitely many prime chains in the arithmetic progressions.
14. In a table of prime numbers there exist infinitely many k -tuples of primes, where $k=2, 3, 4, \dots, 10^5$.

15. Proof of Schinzel's hypothesis.

16. Every large even number is representable in the form $P_1 + P_2 \dots P_n$. It is the n primes theorem which has no almost-primes.

17. Diophantine equation

$$P_{n+1}^{\lambda_{n+1}} = \frac{P_{n+2} + \dots + P_{2n+1} + b}{P_1^{\lambda_1} + \dots + P_n^{\lambda_n} + b}$$
, has infinitely many prime solutions.

18. There are infinitely many primes of the forms: $x^2 + y^n, n \geq 2$ and $mP_1^3 + nP_2^3 (m,n)=1, 2|mn, n \neq \pm b^3$.

19. There are infinitely many prime 5-tuples represented by $P^6 - 42^6 = (P-42)(P+42)(P^2+42P+1764)(P^2-42P+1764)$

20. There are infinitely many prime k -tuples represented by $P^m \pm A^m$.

In this paper by using the arithmetic function $J_2(\omega)$ Santilli's prime chains: $P_{j+1} = aP_j \pm b$ are studied. It is a generalization of Santilli's isoprime m -chains: $P_{j+1} = mP_j \pm (m-1)$ [6].

Santilli's Prime Chains: $P_{j+1} = aP_j \pm b$

Theorem 1

An increasing sequence of primes P_1, P_2, \dots, P_k is called a Santilli's prime chain of the first kind of length k if

$$P_{j+1} = aP_j + b$$

for $j=1, \dots, k-1, (a,b)=1, 2|ab$.

We have the arithmetic function [6]

$$J_2(\omega) = \prod_{3 \leq P \leq \omega} (P - \chi(P))$$

Where $\omega = \prod_{2 \leq P \leq P_i} P$ is called the primorials, P_i the last prime of the primorials.

We now calculate $\chi(P)$. The smallest positive integer such that,

$$a^s \equiv 1 \pmod{P}, (a, b)=1.$$

*Corresponding author: Chun-Xuan Jiang, Institute for Basic Research, Palm Harbor, P.O. Box 3924, Beijing 100854, P.R. China, Tel: +1-727-688 3992; E-mail: jcxuan@sina.com

Received March 10, 2017; Accepted July 29, 2017; Published July 31, 2017

Citation: Jiang CX (2017) Santilli's Prime Chains: $P_j + 1 = aP_j \pm b$. J Generalized Lie Theory Appl 11: 280. doi: 10.4172/1736-4337.1000280

Copyright: © 2017 Jiang CX. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

$$\chi(P) = k \text{ if } k < s; \chi(P) = s \text{ if } k \geq s; \chi(P) = 1 \text{ if } P|ab$$

If $J_2(\omega) = 0$, there exist finitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k . If $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k . It is the Book proof. This is a generalization of the Euclid-Euler proof for the existence of infinitely many primes.

We have the best asymptotic formula of the number of primes $P_1 \leq N$,

$$\pi_k(N, 2) = \frac{J_2(\omega)\omega^{k-1}}{\phi^k(\omega)} \frac{N}{\log^k N} (1 + O(1)),$$

Where $\phi(\omega) = \prod_{2 \leq P \leq \omega} (P-1)$ is called the Euler function of the primorials.

The $P_{j+1} = aP_j - b$ is called a Santilli's prime chain of the second kind of length k . Both $P_{j+1} = aP_j \pm b$ have the same arithmetic function $J_2(\omega)$. If $a = m$ and $b = m - 1$, it is Santilli's isoprime m -chains [6].

Theorem 2

$$P_{j+1} = 2P_j \pm b, j=1, \dots, k-1, b \text{ is an odd number.}$$

We have the arithmetic function [6].

$$J_2(\omega) = \prod_{3 \leq P \leq \omega} (P - \chi(P)) \neq 0.$$

We now calculate $\chi(P)$. The smallest positive integer s such that,

$$2^s \equiv 1 \pmod{P},$$

$$\chi(P) = k \text{ if } k < s; \chi(P) = s \text{ if } k \geq s; \chi(P) = 1 \text{ if } P|b.$$

Since $J_3(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k . This is the Book proof.

We have the best asymptotic formula of the number of primes $P_1 \leq N$.

$$\pi_k(N, 2) = \frac{J_2(\omega)\omega^{k-1}}{\phi^k(\omega)} \frac{N}{\log^k N} (1 + O(1)).$$

The $P_{j+1} = 2P_j \pm 1$ are Cunningham prime chains [6].

Example 1: $P_{j+1} = 2P_j + 7, j=1, 2, 3, 4, 5.$

We have the arithmetic function,

$$J_2(\omega) = 6 \prod_{11 \leq P \leq \omega} (P - 6 - \chi(P)) \neq 0.$$

Where $\chi(31) = -1, \chi(p) = 0$ otherwise.

Since $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_6 are primes.

We have the best asymptotic formula of the number of primes $P_1 \leq N$,

$$\pi_6(N, 2) = \frac{1}{8} \left(\frac{35}{8}\right)^5 \prod_{11 \leq P \leq \omega} \frac{P^5(P-6-\chi(P))}{(P-1)^6} \frac{N}{\log^6 N} (1 + O(1))$$

Theorem 3

$$P_{j+1} = 3P_j \pm b, j=1, \dots, k-1, (3, b) = 1, 2|b.$$

We have the arithmetic function,

$$J_2(\omega) = \prod_{3 \leq P \leq \omega} (P - \chi(P)) \neq 0.$$

We now calculate $\chi(P)$. The smallest positive integer s such that,

$$3^s \equiv 1 \pmod{P},$$

Since $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

We have the best asymptotic formula of the number of primes $P_1 \leq N$,

$$\pi_k(N, 2) = \frac{J_2(\omega)\omega^{k-1}}{\phi^k(\omega)} \frac{N}{\log^k N} (1 + O(1)).$$

Example 2: $P_{j+1} = 3P_j \pm 4, j=1, 2, 3, 4, 5.$

We have the arithmetic function,

$$J_2(\omega) = 96 \prod_{17 \leq P \leq \omega} (P - 6) \neq 0.$$

Since $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_6 are primes.

We have the best asymptotic formula of the number of primes $P_1 \leq N$.

$$\pi_6(N, 2) = \frac{1}{60} \left(\frac{1001}{192}\right)^5 \prod_{17 \leq P \leq \omega} \frac{P^5(P-6-\chi(P))}{(P-1)^6} \frac{N}{\log^6 N} (1 + O(1)).$$

Theorem 4

$$P_{j+1} = 4P_j \pm b, j=1, \dots, k-1, b \text{ is an odd number.}$$

(1) $3|b$, we have the arithmetic function,

$$J_2(\omega) = \prod_{3 \leq P \leq \omega} (P - \chi(P)) \neq 0.$$

We now calculate $\chi(P)$. The smallest positive integer s such that,

$$4^s \equiv 1 \pmod{P}.$$

$$\chi(P) = k \text{ if } k < s; \chi(P) = s \text{ if } k \geq s; \chi(P) = 1 \text{ if } P|b.$$

Since $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

We have the best asymptotic formula of the number of primes $P_1 \leq N$,

$$\pi_k(N, 2) = \frac{J_2(\omega)\omega^{k-1}}{\phi^k(\omega)} \frac{N}{\log^k N} (1 + O(1)).$$

(2) $b \neq 3c, k=3$, we have $J_2(3) = 0$.

(3) $b \neq 3c, k=2$, we have $P_2 = 4P_1 \pm b$. Since $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2 is a prime.

Theorem 5

$$P_{j+1} = 5P_j \pm b, j=1, \dots, k-1, (5, b) = 1, 2|b \text{ is an odd number.}$$

We have the arithmetic function,

$$J_2(\omega) = \prod_{3 \leq P \leq \omega} (P - \chi(P)) \neq 0.$$

We now calculate $\chi(P)$. The smallest positive integer s such that

$$5^s \equiv 1 \pmod{P}.$$

$$\chi(P) = k \text{ if } k < s; \chi(P) = s \text{ if } k \geq s; \chi(5) = 1; \chi(P) = 1 \text{ if } P|b.$$

Since $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

We have the best asymptotic formula of the number of primes $P_1 \leq N$,

$$\pi_k(N, 2) = \frac{J_2(\omega)\omega^{k-1}}{\phi^k(\omega)} \frac{N}{\log^k N} (1 + O(1)).$$

Theorem 6

$P_{j+1} = 6P_j \pm b, j=1, \dots, k-1, (3, b)=1, b$ is an odd number.

(1) $5|b$, we have the arithmetic function

$$J_2(\omega) = \prod_{3 \leq P \leq \omega} (P - \chi(P)) \neq 0$$

We now calculate $\chi(P)$. The smallest positive integer s such that $6 \equiv 1 \pmod{P}$.

$\chi(P) = k$ if $k < s; \chi(P) = s$ if $k \geq s; \chi(3) = 1; \chi(P) = 1$ if $P|b$.

Since $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

We have the best asymptotic formula of the number of primes $P_1 \leq N$,

$$\pi_k(N, 2) = \frac{J_2(\omega) \omega^{k-1}}{\phi^k(\omega)} (1 + O(1)).$$

(2) $b \neq 5c, k=5$, we have $J_2(5)=0$.

(3) $b \neq 5c, k \leq 4$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

Theorem 7

$P_{j+1} = 7P_j \pm b, j=1, \dots, k-1, (7, b)=1, 2|b$.

(1) $6|b$, we have $J_2(\omega) \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

(2) $6|b, k=3$, we have $J_2(3)=0$.

(3) $6|b, k=2$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

Theorem 8

$P_{j+1} = 8P_j \pm b, j=1, \dots, k-1, b$ is an odd number.

(1) $7|b$, we have $J_2(\omega) \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

(2) $b \neq 7c, k=7$, we have $J_2(7)=0$.

(3) $b \neq 7c, k \leq 6$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

Theorem 9

$P_{j+1} = 9P_j \pm b, j=1, \dots, k-1, (3, b)=1, 2|b$.

We have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

Theorem 10

$P_{j+1} = 10P_j \pm b, j=1, \dots, k-1, b$ is an odd number. $(5, b)=1$.

(1) $3|b$, we have $J_2(\omega) \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

(2) $b \neq 3c, k=3$, we have $J_2(3)=0$.

(3) $b \neq 3c, k=2$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

Theorem 11

$P_{j+1} = 11P_j \pm b, j=1, \dots, k-1, 2|b, (11, b)=1$.

(1) $5|b$, we have $J_2(\omega) \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

(2) $b \neq 5c, k=5$, we have $J_2(5)=0$.

(3) $b \neq 5c, k \leq 4$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

Theorem 12

$P_{j+1} = 12P_j \pm b, j=1, \dots, k-1, (3, b)=1, b$ is an odd number.

(1) $11|b$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

(2) $b \neq 11c, k=11$, we have $J_2(11)=0$.

(3) $b \neq 11c, k \leq 10$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

Theorem 13

$P_{j+1} = 16P_j \pm b, j=1, \dots, k-1, b$ is an odd number.

(1) $15|b$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

(2) $b \neq 3c, k=3$, we have $J_2(3)=0. k=2$, we have $J_2(\omega) \infty$ as $\omega \rightarrow \infty$.

(3) $3|b, k=5$, we have $J_2(5)=0. k=4$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

Theorem 14

$P_{j+1} = 17P_j \pm b, j=1, \dots, k-1, 2|b, (17, b)=1$ is an odd number.

We have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

Theorem 15

$P_{j+1} = (2^{\lambda} + 1)P_j \pm b, j=1, \dots, k-1, 2|b, ((2^{\lambda} + 1), b)=1$.

We have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

Theorem 16

$P_{j+1} = (3^{\lambda} + 1)P_j \pm b, j=1, \dots, k-1, 2|b, ((3^{\lambda} + 1), b)=1$ b is an odd number..

(1) $3|b$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

(2) $3|b, k=3$, we have $J_2(3)=0. k=2$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

Theorem 17

$P_{j+1} = (2^{\lambda} \cdot 3^{\lambda_2} + 1)P_j \pm b, j=1, \dots, k-1, 2|b, ((2^{\lambda} \cdot 3^{\lambda_2} + 1), b)=1$.

(1) $6|b$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

(2) $b \neq 6c, k=3$, we have $J_2(3)=0. k=2$, we have $J_2(\omega) \infty$ as $\omega \rightarrow \infty$.

Theorem 18

$P_{j+1} = (3^{\lambda_1} \cdot 5^{\lambda_2} + 1)P_j \pm b, j=1, \dots, k-1, b$ is an odd number.

$$((3^{\lambda_1} \cdot 5^{\lambda_2} + 1), b) = 1$$

(1) $15|b$, we have $J_2(\omega) \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

(2) $b \neq 3c, k=3$, we have $J_2(3)=0. k=2$, we have $J_2(\omega) \infty$ as $\omega \rightarrow \infty$.

(3) $b \neq 3c, k=5$, we have $J_2(5)=0. k \leq 4$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

Theorem 19

$P_{j+1} = (3^{\lambda_1} \cdot 5^{\lambda_2} \cdot 7^{\lambda_3} + 1)P_j \pm b, j=1, \dots, k-1, b$ is an odd number.

$$((3^{\lambda_1} \cdot 5^{\lambda_2} \cdot 7^{\lambda_3} + 1), b) = 1$$

(1) $105|b$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

(2) $3|b, k=5$, we have $J_2(5)=0$. $k \leq 4$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

(3) $3|b, k=3$, we have $J_2(3)=0$. $K=2$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

Theorem 20

$P_{j+1} = (3^{\lambda_1} \cdot 5^{\lambda_2} \cdot 7^{\lambda_3} \cdot 11^{\lambda_4} + 1)P_j \pm b, j=1, \dots, k-1, b$ is an odd number.

$$(3^{\lambda_1} \cdot 5^{\lambda_2} \cdot 7^{\lambda_3} \cdot 11^{\lambda_4} + 1, b) = 1.$$

(1) $1155|b$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

(2) $3|b, k=5$, we have $J_2(5)=0$. $k \leq 4$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

(3) $b3c, k=3$, we have $J_2(3)=0$. $K=2$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

Theorem 21

$P_{j+1} = (7^{\lambda_1} \cdot 19^{\lambda_2} \cdot 31^{\lambda_3} + 1)P_j \pm b, j=1, \dots, k-1, b$ is an odd number.

$$(7^{\lambda_1} \cdot 19^{\lambda_2} \cdot 31^{\lambda_3} + 1, b) = 1$$

(1) $4123|b$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k .

(2) $7|b, k=19$, we have $J_2(19)=0$. $k \leq 18$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

(3) $b7c, k=7$, we have $J_2(7)=0$. $K \leq 6$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$.

Theorem 22

$P_{j+1} = aP_j \pm b, j=1, \dots, k-1, (a, b) = 1, 2|ab$.

If $a-1 = P_1^{\lambda_1} \dots P_n^{\lambda_n}, P_1 \dots P_n | b$, we have $J_2(\omega) \rightarrow \infty$ as $\omega \rightarrow \infty$. There exist infinitely many primes P_1 such that P_2, \dots, P_k are primes for arbitrary length k [6].

Euclid-Euler-Jiang Theorem

Around 300BC by using the equation

$$(\omega+1, \omega) = 1 \text{ as } \omega \rightarrow \infty,$$

Euclid proved that there are infinitely many primes.

In 1748 by using the equation

$$\frac{\omega}{\phi(\omega)} = \prod_{i=1}^{\infty} \left(1 + \frac{1}{P_i}\right) = \sum_{n=1}^{\infty} \frac{1}{n} \rightarrow \infty \text{ as } \omega \rightarrow \infty,$$

Euler proved that there are infinitely many primes.

By using the equation [1-10].

$$J_2(\omega) \rightarrow \infty \text{ as } \omega \rightarrow \infty.$$

Jiang has proved that there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes [1-10]. It is a generalization of Euclid-Euler theorem. Therefore Euclid-Euler-Jiang theorem in the distribution of primes is advanced. It is the Book theorem.

From ref. [6] we have:

$$\frac{\omega}{\phi(\omega)} \prod_{2 \leq P \leq N} \left(1 + \frac{1}{P}\right) \sim \frac{C_1}{\log N}$$

Therefore we have the prime number theorem.

$$\pi(N) \sim \frac{N}{\log N}$$

Where $\pi(N)$ denotes the number of primes $\leq N$.

From ref. [6] we have:

$$\frac{J_2(\omega)}{\omega} \sim B_1 \prod_{k \leq P \leq N} \left(1 + \frac{k}{P}\right) \sim \frac{B_2}{\log^k N}$$

Therefore we have the prime k -tuples theorem:

$$\pi_k(N, 2) \sim C_k \frac{N}{\log^k N},$$

Where $\pi_k(N, 2)$ denotes the number of primes $P_1 \leq N$.

If the arithmetic constant $C_k \frac{J_2(\omega)\omega^{k-1}}{\phi^k(\omega)} \neq 0$, that is $J_2(\omega) \neq 0$, there exist infinitely many primes P_1 such that P_2, \dots, P_k are primes. $\pi_k(N, 2)$ have the same form $\frac{N}{\log^k N}$ but differ in C_k .

References

- Jiang CX (1996) On the Yu-Goldbach prime theorem. Guangxi Sciences (in Chinese) 3: 9-12.
- Jiang CX (1998) Foundations of Santilli's isonumber theory 1. Algebras, Groups and Geometries 15: 351-393. MR2000c: 11214.
- Jiang CX (1998) Foundations of Santilli's isonumber theory 2. Algebras, Groups and Geometries 15: 509-544.
- Jiang CX (1999) Foundations of Santilli's isonumber theory. In: Fundamental open problems in sciences at the end of the millennium, T. Gill, K. Liu and E. Trel (Eds) 13 Hadronic Press, USA pp: 105-139.
- Jiang CX (2001) Proof of Schinzel's hypothesis, Algebras Groups and Geometries 18: 411-420.
- Jiang CX (2002) Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture. International Academic Press. www. i-b-r.org
- Jiang CX, Disproofs of Riemann hypothesis. To appear.
- Jiang CX, Prime theorem in Santilli's isonumber theory. To appear.
- Jiang CX, Prime theorem in Santilli's isonumber theory (II). To appear.
- Jiang CX, Diophantine equation $P_{n+1}^{\lambda_{n+1}} = \frac{P_{n+2} + \dots + P_{2n+1} + b}{P_1^{\lambda_1} + \dots + P_n^{\lambda_n} + b}$ has infinitely many prime solution. To appear.