

# Schiff Base N-(5-Chlorosalicylidene) Aniline, a Novel Antifungal Agent: Insights from Crystallographic Analysis, Semi Empirical and Molinspirations Calculations

Shalini Suresh<sup>1</sup>, Girija CR<sup>2\*</sup>, Sathish CD<sup>3</sup> and Venkatesha TV<sup>4</sup>

<sup>1</sup>Chemistry Research Centre (Affiliated to Kuvempu University), SSMRV Degree College, Jayanagar 4th T-Block, Bengaluru-560 041, Karnataka, India

<sup>2</sup>Government Science College, Nrupatunga Road, Bengaluru-560 001, Karnataka, India

<sup>3</sup>Department of Chemistry, PESIT, BSK III Stage, Bengaluru-560 085, Karnataka, India

<sup>4</sup>Department of Chemistry, Jnana Sahyadri, Kuvempu University, Shankaraghatta-577 451, Karnataka, India

## Abstract

The single crystals of a Schiff base, N-(5-chlorosalicylidene) aniline suitable for X-ray diffraction were grown by slow evaporation from its ethanol solution. The geometrical, supramolecular features, energetics and biological activity of the title compound were analysed through a combined experimental and theoretical approach. The compound crystallizes in orthorhombic system with the space group  $Pca2_1$ . The unit cell parameters  $a=12.391(6)$  Å,  $b=4.5223(18)$  Å,  $c=19.514(8)$  Å,  $\alpha=\beta=\gamma=90.00^\circ$ .  $V=1093.5(8)$  Å<sup>3</sup>,  $Z=4$ ,  $D_x=Mg\ m^{-3}$  and  $\mu=0.32\ mm^{-1}$ . The crystal packing is mainly stabilized by dispersion forces from short H...H contacts. Also an intramolecular N-H...O hydrogen bonding is observed. For the optimised geometry, the HOMO-LUMO energy gap (7.071 eV) reflects the chemical activity. The compound is expected to have moderate hydrophilicity and hydrophobicity. Hence, have moderate permeability across cell membrane. Bioactivity scores suggest weakly bioactive nature of the compound.

**Keywords:** Schiff base; Non-covalent interactions; MOPAC; Semi-empirical calculations; Mollog calculations

## Introduction

Schiff bases are the compounds carrying imine or azomethine ( $-C=N-$ ) functional group. These are the condensation products of primary amines with carbonyl compounds. Schiff base ligands contain a variety of substituents with different electron-donating or electron-withdrawing groups, and therefore may have interesting chemical properties. They have attracted particular interest due to their biological activities such as radiopharmaceuticals for cancer targeting. They have also been used as model systems for biological macromolecules. Schiff bases have gained importance in medicinal and pharmaceutical fields due to a broad spectrum of biological activities like anti-inflammatory, analgesic, antimicrobial, anticonvulsant, anti-tubercular, anticancer, antioxidant, antihelminthic, and so forth. The nitrogen atom of azomethine may be involved in the formation of a hydrogen bond with the active centres of cell constituents and interferes in normal cell processes. Concisely, Schiff bases are among the molecules which have therapeutic potential for the treatment of various human diseases. These properties are greatly influenced by the topochemistry of the Schiff base molecules which in turn is highly affected by the crystal structure. Therefore, the study of crystal packing and the intermolecular interactions in the crystal structures of various Schiff bases can lead to valuable data for the design and synthesis of new materials [1].

In view of this importance, the crystal structure analysis of title compound (I) and semi-empirical quantum chemical calculations were performed to investigate the role of intermolecular interactions in controlling the crystal packing and thereby influencing the biological activity.

## Materials and Methods

### X-ray structure analysis

The IUPAC name of the compound (I), N-(5-chlorosalicylidene) aniline is 4-chloro-2-[(E)-(phenylimino) methyl] phenol and is shown in the Figure 1. Single crystals of the compound (I),  $C_{13}H_{10}ClNO$

were grown by slow evaporation method using ethanol as solvent at room temperature. A suitable crystal was selected and mounted on a Bruker Smart [2] CCD Area Detector System using Mo-K $\alpha$  (0.71073 Å) radiation for the crystal. The crystal was kept at 296(2) K during data collection. Intensity data were collected up to a  $\theta_{max}$  of 25.0° for the compound in the  $\omega$ - $\phi$  scan mode. A total of 3489 reflections were collected, resulting in 1561 independent reflections of which the number of reflections satisfying  $I > 2\sigma(I)$  criteria were 1500, and these were treated as observed. Corrections for Lorentz and polarisation effects were applied. Using Olex2 [3], the structure was solved with the olex2.solve [4] structure solution program using Charge Flipping and refined with the olex2.refine [5,6] refinement package using Gauss-Newton minimisation. The final R factor is 0.034. The details of crystal data, data collection and the refinement are given in Table 1.

### Computational studies

**MOPAC calculations:** Despite the interesting properties of N-salicylideneaniline derivatives, have been scarcely studied from a computational point of view [7]. Our goal was to compute quantum chemical derived properties that would be useful as starting points for understanding the properties of this type of ring system. Moreover, the other main task of conformational analyses of isolated molecule (I) was to examine the stable conformation and a global energy minimum for the studied molecule. If there was considerable energy difference

**\*Corresponding author:** Girija CR, Chemistry Research Centre (Affiliated to Kuvempu University), SSMRV Degree College, Jayanagar 4th T-Block, Bengaluru-560 041, Karnataka, India, E-mail: [girija.shivakumar@rediffmail.com](mailto:girija.shivakumar@rediffmail.com)

Received March 07, 2016; Accepted March 08, 2016; Published March 16, 2016

**Citation:** Suresh S, Girija CR, Sathish CD, Venkatesha TV (2016) Schiff Base N-(5-Chlorosalicylidene) Aniline, a Novel Antifungal Agent: Insights from Crystallographic Analysis, Semi Empirical and Molinspirations Calculations. Chem Sci J 7: 122. doi:[10.4172/2150-3494.1000122](http://dx.doi.org/10.4172/2150-3494.1000122)

**Copyright:** © 2016 Suresh S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: Schematic diagram of the title compound(I), (C<sub>13</sub>H<sub>10</sub>ClNO).

between the lowest energy types of conformer, then we concluded that theoretical calculations predicted one type of geometric molecule. Since, the size and the variety of heteroatoms in the N-(5-chlorosalicylidene) aniline are considerable, a full semi-empirical geometrical optimization is computationally very demanding. This work is simplified, if we use a realistic structure as starting geometry for the MM conformational search and quantum-chemical calculation. Therefore, the structure obtained by X-ray diffraction analysis is suitable to this end. The X-ray structure of the compound (I) is used as input to MOPAC which was optimized using molecular mechanics (MM) incorporated in Gabedit (GUI) software [8] using Newton Raphson method. Semi-empirical Quantum Chemical Calculations were performed on the refined parameters using MOPAC 2012 program [9] to optimize the structure with Parametrization Model 7 (PM7) approximation. The minimizations were terminated at r.m.s. gradient of less than 0.01 kJ·mol<sup>-1</sup> Å<sup>-1</sup>. Electronic structure calculations provide useful estimates of the energetic properties of chemical systems, including molecular structures, spectroscopic features and probable reaction pathways. The PM7 Hamiltonian in MOPAC package was used to calculate the bond lengths, bond angles, heats of formations, core-core repulsion energies, ionization potentials etc.

### Molinspiration calculations

Quantitative Structure Activity Relationship (QSAR) studies were applied to find correlation between different calculated molecular descriptors of the compound (I) and biological activity. Structure of the title compound (I) was drawn in the online molinspiration software version 2014.06 (www.molinspiration.com) [10] for calculation of molecular properties and prediction of bioactivity score for drug targets.

## Results and Discussion

### X-ray crystal structure analysis

**Geometrical features:** The structure of 4-chloro-2[(E)-(phenylimino) methyl] phenol, C<sub>13</sub>H<sub>10</sub>ClNO, has orthorhombic *Pca*2<sub>1</sub> symmetry. The compound (I), C<sub>13</sub>H<sub>10</sub>ClNO, is planar with a dihedral angle of 9.49° between the planes of the two benzene rings. The compound exists in a trans-configuration about the central C=N bond [C7-N1=1.278 (3)Å] with a torsion angle C5-N1-C7-C8=-178.9(2)°. All other selected bond distances [11], bond angles and torsional angles are given in Tables 2-4, respectively have the normal values. The ORTEP [12] plot of the compound (I) is as shown in the Figure 2.

**Supramolecular features:** Intramolecular hydrogen bonding is observed between the oxygen atom of keto group and the H-atom of imine group, resulting in a keto-imine tautomeric form. A strong intramolecular N1—H1...O6 hydrogen bond generates an S(6) ring motif [13] as shown in the Figure 3a and 3b. A short H4...H7 contact [2.0984(5) Å] is also observed. The hydrogen bond geometry is listed

in Table 5.

### MOPAC calculations

**Geometry optimization:** The significant difference in C-N, C-O bond distances of atoms involved in intramolecular hydrogen bond interactions are observed between refined X-ray data and optimized geometrical data. The comparison of these results along with bond distances, angles and torsion angles are given in Table 6 and that of hydrogen bond geometries are given in Table 5. The change in torsion angles indicates how the crystal packing effect significantly influences these interactions.

**HOMO-LUMO studies:** The heat of formation is equal to -12.66715 kcal mol<sup>-1</sup> for the molecule in the asymmetric unit. The ionization potential, dipole moment and self-consistency field (SCF) factor are calculated as 8.516512 eV, 4.945 Debye and 40 respectively. The geometry optimization using MOPAC2012 results in HOMO-LUMO energies as -8.517 and -1.446 eV respectively. LUMO as an electron

Crystal Data	
Molecular formula, Mol. mass	C <sub>13</sub> H <sub>10</sub> ClNO, 231.67
Unit Cell Parameters	a=12.391 (6)Å, b=4.5223 (18) Å, c=19.514 (8) Å, α=β=γ=90.00°
	V=1093.5 (8) Å <sup>3</sup> , Z=4, D <sub>x</sub> =Mg m <sup>-3</sup> and m=0.32 mm <sup>-1</sup>
Crystal System, space group	Orthorhombic, <i>Pca</i> 2 <sub>1</sub>
<b>Data collection</b>	
Diffractometer	Bruker SMART CCD area detector
Data collection method	ω / φ, mode
Absorption correction	multi-scan (SADABS)
No. of measured reflections	3489
No. of independent reflections	1561
No. of observed reflections	1500
Criterion for observed reflections	I > 2σ (I)
R <sub>int</sub>	0.034
θ <sub>max</sub> (°)	4.2, 25.0°
Range of h, k, l	h=-12→14 k=-5→5 l=-19→23
<b>Refinement</b>	
Refinement on	F <sup>2</sup>
R[F <sup>2</sup> >2σ(F <sup>2</sup> )]	0.033
w/R	0.089
S	1.05
No. of reflections used in refinement	1561
No. of parameters used	149
No. of restraints applied	1
Weighting scheme W=1/[σ <sup>2</sup> (F <sub>o</sub> ) + (0.0894P) <sup>2</sup> + 2.0000P] Where P=(F <sup>2</sup> +2F <sub>o</sub> <sup>2</sup> )/3	
Δρ <sub>max</sub> (e Å <sup>-3</sup> )	0.25 e Å <sup>-3</sup>
Δρ <sub>min</sub> (e Å <sup>-3</sup> )	-0.30 e Å <sup>-3</sup>
Friedel pairs	480
Absolute structure parameter [Flack parameter] [6]	-0.01 (8)

Table 1: Crystal data, data collection and refinement details of compound(I).

O -C12 1.355(3)	N1 -C5 1.409(3)
N1 -C7 1.278(3)	
<b>Short Contacts (Å)</b>	
H4-H7 2.1000	



Table 2: Selected Bond Distances (Å).

C5-N1-C7	122.75(19)	N1-C5-C6	116.97(19)
N1-C5-C4	24.0(2)	N1-C7-C8	121.6(2)
C10-C11-C12	120.9(2)	O6-C12-C11	119.14(19)
O6-C12-C8	121.2(2)		

Table 3: Selected Bond Angles (°).

C7-N1-C5-C6	-177.0(2)	C5-N1-C7-C8	-178.9(2)
C7-N1-C5-C4	2.7(3)	C3-C4-C5-N1	-179.4(2)
N1-C5-C6-C1	179.5(2)	N1-C7-C8-C9	177.1(2)
N1-C7-C8-C12	-3.9(4)	C7-C8-C12-O6	1.0(4)
C7-C8-C12-C11	-180.0(2)	C8-C9-C13-C11	78.41(18)
C10-C11-C12-O6	179.2(2)		

Table 4: Selected Torsion Angles (°).



acceptor (electrophilic) represents the ability to obtain an electron; HOMO represents the ability to donate an electron (nucleophilic). The HOMO-LUMO energy gap of 7.071 eV reflects the chemical activity of molecule, resulting in charge transfer interaction from  $\pi$ -bonding MO of a phenyl ring from the salicylidene moiety in HOMO to a  $\pi^*$ -anti-bonding MO of the aniline moiety extended to imine group in LUMO (Figure 4). According to the Pearson's maximum hardness principle (PMHP) the molecules arrange themselves with maximum hardness [14-17]. The conformation has the maximum hardness value, which is the minimum energy conformation. In the finite different approximation, the ionization energy and electron affinity can be replaced by the HOMO and LUMO energy, respectively, using Koopmans' theorem, [15] within a Hartree-Fock and/or density functional theory schemes, with

$$I = -\epsilon_{\text{HOMO}} \text{ and } A = -\epsilon_{\text{LUMO}} \text{ yielding}$$

$$\mu = -1/2(\epsilon_{\text{HOMO}} + \epsilon_{\text{LUMO}}) \text{ and } \eta = 1/2(\epsilon_{\text{HOMO}} - \epsilon_{\text{LUMO}})$$

The calculated chemical hardness  $\eta$  is found to be 3.535 eV and chemical potential  $\mu$  4.9815 eV. The energy values and dipole moments resulting from Semi-empirical Quantum Chemical Calculations are given in Table 7. A larger value of the energy gap indicates low reactivity to a chemical species because the energy gap is related to the softness or hardness of a molecule. A soft molecule is more reactive than a hard molecule because a hard molecule has a larger energy gap. Dipole moment ( $\mu$ ) is the measure of polarity in a bond and is related to the distribution of electrons in a molecule. Schiff bases are an important class of organic compounds which, has a high dipole moment and is capable of hydrogen bonding, which could be favourable in the binding of biomolecular targets.

### Packing coefficient

70.2% as calculated using the program PLUTON [18]. The crystallographic symmetry elements present in the compound are 2-fold screw axis and glide plane. The crystal structure is thermodynamically favoured as the packing efficiency is relatively high.

### Molinspiration calculations

The molecular properties (miLog P, Total polar surface area, number of hydrogen bond donors and acceptors, molecular weight, number of atoms, number of rotatable bonds etc.) and the bioactivity scores for drug targets (GPCR ligands, kinase inhibitors KI, ion channel modulators ICM, enzymes inhibitor EI, Protease inhibitor PI and nuclear receptors NRL) of the title compound (I) are valued in Table 8. The miLogP value of the compound (I) is found to be 4.07 which clearly obeys Lipinski's rule of five [19]. The compound (I) is expected to have moderate hydrophilicity and also moderately lipophilic. This indicates that the compound will have moderate permeability across cell membrane. The title compound (I) having low molecular mass (231.68) [drug molecules (<500)] is known to be easily transported, diffuse and absorbed as compared to heavy molecules. Total polar surface area (TPSA) is found to be 32.59 (<160 Å limit), closely related to the hydrogen bonding potential of a molecule and is a very good predictor of drug transport properties such as intestinal absorption, bioavailability, blood brain barrier penetration etc.

Number of rotatable bonds is a simple topological parameter that measures molecular flexibility and is considered to be a good descriptor of oral bioavailability of drugs. The screened compound (I) is flexible with (2 rotatable bonds). As a general rule, larger is the bioactivity score, higher is the probability that investigated compound will be active. Therefore, a molecule having bioactivity score more than 0.00 is most likely to possess considerable biological activities, while values -0.50 to 0.00 are expected to be moderately active and if score is less than -0.50 it is presumed to be inactive. The results of the present study demonstrated that the investigated compound is biologically inactive molecule. The bioactivity score for GPCR ligand is found to be -0.87, for ion channel modulator activity is -0.81. Similar results are obtained for kinase inhibition showed score -0.59, indicates the molecules were inactive towards this drug target. Bioactivity scores for nuclear receptor ligand, protease inhibitor and enzyme inhibition was found to be in the range of -0.82, -0.99 and -0.43 respectively, suggest weakly bioactive nature of molecules. The interaction between the HOMO drug with the LUMO receptor and vice-versa play important role in intermolecular interactions [20], the interactions stabilised inversely with energy gap between the interacting orbitals. Increasing HOMO energy and




Figure 3b: Crystal packing diagram of the title compound(I) showing intermolecular interactions

D-H...A	D-H		H...A		D...A		D-H...A	
	XRD	MOPAC	XRD	MOPAC	XRD	MOPAC	XRD	MOPAC
N1-H1...O6	0.8600	1.061	1.9100	1.804	2.610(3)	2.607	137.00	129.11

Symmetry codes: (i) x, y, z

Table 5: Hydrogen Bond geometry (Å, °).

Parameters	XRD refined	MOPAC2012
<b>Bond length (Å)</b>		
C7-N1	1.278(2)	1.349
C12-O6	1.354(3)	1.229
<b>Bond angle (°)</b>		
C8-C12-C11	119.6(2)	118.21
N1-C7-C8	121.5(2)	121.80
<b>Dihedral angle (°)</b>		
C5-N1-C7-C8	-178.9(2)	-179.742
C4-C5-N1-C7	2.7(3)	0.046
N1-C7-C8-C9	177.1(2)	-179.92


Table 6: Optimized geometrical parameters for the compound(I).

<b>Final heat of formation</b>	-12.66715 kcal/mol=-52.99936 kJ/mol
<b>Total energy (eV)</b>	-2479.37797
<b>Electronic energy (eV)</b>	-14419.00509
<b>Point group</b>	Cs
<b>Core-core repulsion (eV)</b>	11939.62711
<b>Ionization potential (eV)</b>	8.516512
<b>HOMO LUMO energies (eV)</b>	-8.517 , -1.446
<b>No. of filled levels</b>	40
<b>Dipole moment (in D)</b>	4.945

Table 7: Energy values obtained from optimised geometry of the compound(I).

Compound	Molinspiration calculations						Drug-likeness					
	MW (g/mol)	miLogP	TPSA	nOH-NH	nrotb	Volume	GPCRL	ICM	KI	NRL	PI	EI
(I)	231.68	4.07	32.59	1	2	200.27	-0.87	-0.81	-0.59	-0.82	-0.99	-0.43

Table 8: Molecular properties and bioactivity score of compound(I).



decreasing LUMO energy in the drug molecule lead to enhanced stabilising interactions and hence, binding with the receptor.

## Conclusion

The title compound (I) is mainly stabilized by strong intra molecular N-H...O hydrogen bond and also dispersion forces from H...H contacts. The HOMO-LUMO energy gap of 7.071 eV reflects the chemical activity of molecule, resulting in charge transfer interaction from  $\pi$ -bonding MO of a phenyl ring from the salicylidene moiety in HOMO to a  $\pi^*$ -anti-bonding MO of the aniline moiety extended to imine group in LUMO. The dipole moment of 4.945D indicates moderately polar nature. The compound (I) is expected to have moderate hydrophilicity and lipophilicity. Hence, have moderate permeability across cell membrane. The title compound (I) having low molecular mass (231.68) known to be easily transported, diffuse and absorbed as compared to heavy molecules. Total polar surface area (TPSA) is found to be 32.59, closely related to the hydrogen bonding potential of a molecule and is a very good predictor of drug transport properties such as intestinal absorption, bioavailability and blood brain barrier penetration. Bioactivity scores for nuclear receptor ligand, protease inhibitor and enzyme inhibition suggest weakly bioactive nature of the title compound (I).

Computational analysis suggests that the dipole moment and geometry of molecule may be good descriptors for the QSAR studies of novel N-salicylideneanilines as potent antibacterial agents.

## Acknowledgements

The authors thank Solid State Chemistry Unit (SSCU), Indian Institute of Science (IISc), Bengaluru, Karnataka, India, for the X-ray intensity data collection. SS thank Rashtriya Sikshana Samsthe Trust (RSST) and the Principal, SSMRV Degree College, Bengaluru, for their constant support and encouragement for carrying out research work.

## Competing Interest

Authors have declared that no competing interests exist.

## Supplementary Data

The CIF file was deposited at the Cambridge Crystallographic Data Centre, The deposition number is CCDC- 1444547. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), by e-mailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK.

## References

1. Blagus A, Dominik C, Friscic T, Kaitner B, Stilinovic V (2010) Schiff bases derived from hydroxyaryl aldehydes: molecular and crystal structure, tautomerism, quinoid effect, coordination compounds. Macedonian Journal of Chemistry and Chemical Engineering 29: 117-138.

2. Bruker APEX2, SAINT and SADABS (2009) Bruker AXS Inc., Madison, Wisconsin, USA.
3. Dolomanov OV, Bourhis LJ, Gildea RJ, Howard JAK, Puschmann H (2009) OLEX2: a complete structure solution, refinement and analysis program. J Appl Cryst 42: 339-341.
4. Bourhis LJ, Dolomanov OV, Gildea RJ, Howard JA, Puschmann H (2015) The anatomy of a comprehensive constrained, restrained refinement program for the modern computing environment - Olex2 dissected. Acta Crystallogr A Found Adv 71: 59-75.
5. Flack HD (1983) On enantiomorph-polarity estimation. Acta Crystallography A39: 876-881
6. Al-Swaidan IA, SEI-Azab A, Alanazi AM, Abdel-ziz AAM (2014) Synthesis and Conformational Analysis of Sterically Congested (4R)-(-)-1-(2,4,6-Trimethylbenzenesulfonyl)-3-n-butyl-4-tertbutyl-2-imidazolidinone: X-Ray Crystallography and Semi empirical Calculations. Journal of Chemistry 2014: 15.
7. Allouche AR (2011) Gabedit—a graphical user interface for computational chemistry softwares. J Comput Chem 32: 174-182.
8. MOPAC2012 (2012) James JP Stewart, Stewart Computational Chemistry, Version 14.083W.
9. <http://www.molinspiration.com/>
10. Allen FH, Kennard O, Watson DG, Brammer L, Orpen AG, et al. (1987) Tables of bond lengths determined by X-ray and neutron diffraction. Part 1. Bond lengths in organic compounds. Journal of Chemical Society. Perkin Transactions, 2: S1-19.
11. Farrugia LJ (2012) WinGX and ORTEP for Windows: an update. Journal of Applied Crystallography 45: 849-854.
12. Etter MC, MacDonald JC, Bernstein J (1990) Graph-set analysis of hydrogen-bond patterns in organic crystals. Acta Crystallogr B 46: 256-262.
13. Kar T, Scheiner S, Sannigrahi AB (1998) Hardness and Chemical Potential Profiles for Some Open-Shell HAB  $\rightarrow$  HBA Type Reactions. Ab Initio and Density Functional Study. J Phys Chem A 102: 5967-5973.
14. Koopmans TA (1934) Über die Zuordnung von Wellenfunktionen und Eigenwerten zu den Einzelnen Elektronen Eines Atoms. Physica 1: 104-113.
15. Pople JA, Beveridge DL (1970) Approximate Molecular Orbital Theory, Advanced Chemistry.
16. Arrindell WA, Eisemann M, Richter J, Oei TP, Caballo VE, et al. (2003) Masculinity-femininity as a national characteristic and its relationship with national agoraphobic fear levels: Fodor's sex role hypothesis revitalized. Behav Res Ther 41: 795-807.
17. Ghosh DC (1988) A study of correlation of the order of chemical reactivity of a sequence of binary compounds of nitrogen and oxygen in terms of frontier orbital theory. J Indian Chem Soc 65: 554-556.
18. Spek AL (2009) Structure validation in chemical crystallography. Acta Crystallogr D Biol Crystallogr 65: 148-155.
19. Lipinski CA (2004) Lead- and drug-like compounds: the rule-of-five revolution. Drug Discov Today Technol 1: 337-341.
20. Fukui K (1982) Role of frontier orbitals in chemical reactions. Science 218: 747-754.

**Citation:** Suresh S, Girija CR, Sathish CD, Venkatesha TV (2016) Schiff Base N-(5-Chlorosalicylidene) Aniline, a Novel Antifungal Agent: Insights from Crystallographic Analysis, Semi Empirical and Molinspirations Calculations. Chem Sci J 7: 122. doi:10.4172/2150-3494.1000122