

# Signal or To Sink: The Long-Sought-After Cytokinin Transporter

Shabana Shams<sup>1</sup> and Muhammad Naseem<sup>2,3\*</sup>

<sup>1</sup>Department of Animal Sciences, Quaid-i-Azam University Islamabad, Pakistan

<sup>2</sup>Department of Bioinformatics, Functional Genomics and Systems Biology Group, Biocenter, University of Würzburg, Am Hubland, Würzburg, Germany

<sup>3</sup>Department of Molecular Biology and Genetics, Systems Biology of Plant-Microbes Interaction Group, Kuzey Park, Istanbul


## Description

An interesting mechanism of trade-off between cytokinin signaling and cellular cytokinin transport through Purine Permease 14 (PUP14) protein is very recently unfolded by Zürcher et al. [1]. The perception of the signal of cytokinin and its cellular phosphorelay circuitry [2] that result in target genes expression are well understood (Figure 1). However, the patchy distribution of bioactive cytokinin ligands, which has vast implications for tissue patterning and multicellularity in plants, is still under research [2,3]. When and where not to signal; is a fine-tuned spatiotemporal regulation that needs clearance of the active cytokinins-ligands away from the cytokinin signaling receptors histidine kinases (Figure 1). In addition to the transport of cytokinins to distal tissues, cells also can achieve the clearance of active cytokinins from apoplast by importing them into the cytosol. Inside the cytosol, cytokinins may also interact with various soluble proteins. Consistent with this notion, and besides membrane-bound cytokinin receptors, many other cytokinin binding proteins (CBPs) have been detected in plants [4]. However, the cellular mechanism of the entry of cytokinins from the apoplast to cytosol was not fully understood until very recently. The findings of Bruno group [1] concerning the transport of apoplastic cytokinins through PUP14 transporter not only fills this knowledge gap but also has opened new avenues for research in cytokinin biology.

Taking the bioactive cytokinin ligands away from the CHASE-domain of sensor AHKs, the activity of PUP14 enables the import of apoplastic cytokinin inside the cytosol (Figure 1). This influx of bioactive cytokinins exhausts apoplastic space of active ligands for AHKs, and that culminates in reduced cytokinin signaling as well as diminished cellular cytokinin responses (Figure 1). It is worth noting that PUP14 mediated uptake of the active cytokinins was not interfered by cytokinin-ribosides, which suggests that only active ligands are cleared from the apoplast. Zürcher et al. [1] have meticulously exemplified this phenomena on various morphogenetic contexts; for instance, heart-stage embryo, lateral root primordia as well root and shoot apical meristems. The clearance of active ligands away from cell surface cytokinin receptors has no doubt spatiotemporal implications for plant morphogenesis; however, below I highlight some structural biology prospects of this new finding.

Either signal or to sink; for an active cytokinin ligand, the possibility to remain in the apoplast or sink-down into the cytosol is highly determined by the morphogenetic dynamics of a tissue [1]. However, what are those biochemically competing circumstances that propel active cytokinin species away from a functional cytokinin receptor and let them pass through the barrel of the PUP 14 transporter at the expense of adenosine tri-phosphate (ATPs), merits further investigation. Likewise, what is the biochemical explanation of selectivity in ligand-clearance where bioactive cytokinin ligands are imported into the cell while riboside forms are left behind in the apoplast [1]. Therefore, future efforts addressing these aspects with structural biology approaches will further decipher the understanding of this important finding.

One may tempt to speculate that besides external morphogenetic dynamics, the process of cytokinin influx might also be triggered by inside-out signaling. Reason being, the plant cell also contains cytosolic cytokinin receptors [2] which are located on endoplasmic reticulum (ER) and need active ligands for their activity. Although the bulk of



**Left:** In the absence of a functional cytokinin transporter (PUP14: cylindrical shape with cross), the plasma-membrane localized cytokinin signaling receptors sensor Histidine Kinases (AHKs: green rectangles) perceive active cytokinin ligands (green filled circles), transmit strong phosphorelay signal (more green arrow) and strongly activate cytokinin response genes (more red arrows).

**Right:** During the influx of cytokinin, the PUP14 (with blue arrow) transporter channelizes active ligands to the cytosol and leaving behind ribosides forms of cytokinins (orange filled circles). This results in attenuated cytokinin signaling and response (less number of green arrows).

**Figure 1:** Cellular cytokinin transport through Purine Permease 14 (PUP14) transporter impedes cytokinin signaling and responses.

transported (through PUP14) cytokinins are converted into inactive ligands but the possibility of interaction between freshly transported cytokinins and ER-based AHKs still exists (Figure 1). Genetic studies focusing on mutation in the CHASE-domain of cytoplasmic cytokinin receptors and PUP14 transporter would be helpful in addressing this hypothesis. Lastly, plant apoplast is a seat of many other biological functions such as immunity to pathogens [5,6] as well as long-distance communication [2,7], it would be much intriguing to explain these and alike processes under the auspices of this new development in cytokinins biology.

## References

- Zürcher E, Liu J, Di Donato M, Geisler M, Müller B (2016) Plant development regulated by cytokinin sinks. *Science* 353: 1027-1030.
- Hwang I, Sheen J, Müller B (2012) Cytokinin signaling networks. *Annu Rev Plant Biol* 63: 353-380.
- Takatashi K, Kentaro T, Kojima M, Sakakibara H (2013) Side-chain modification of cytokinins controls shoot growth in Arabidopsis. *Developmental Cell* 2: 452-461.
- Wang Y, Letham DS, John PC, Zhang R (2016) Synthesis of a cytokinin linked by a spacer to dexamethasone and biotin: Conjugates to detect cytokinin-binding proteins. *Molecules* 21: 576.

\*Corresponding author: Muhammad Naseem, Department of Bioinformatics, Functional Genomics and Systems Biology Group, Biocenter, University of Würzburg, Am Hubland, Würzburg, Germany. Tel: +49931 310; E-mail: [muhammad@biozentrum.uni-wuerzburg.de](mailto:muhammad@biozentrum.uni-wuerzburg.de)

Received August 01, 2017; Accepted August 05, 2017; Published August 12, 2017

Citation: Shams S, Naseem M (2017) Signal or To Sink: The Long-Sought-After Cytokinin Transporter. *Transcriptomics* 5: 141. doi:10.4172/2329-8936.1000141

Copyright: © 2017 Shams S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

5. Naseem M, Wolfling M, Dandekar T (2014) Cytokinins for immunity beyond growth, galls and green islands. *Trends in Plant Sci* 8: 481-484.
6. Yamada K, Saijo Y, Nakagami H, Takano Y (2016) Regulation of sugar transporter activity for antibacterial defense in *Arabidopsis*. *Science* 354: 1427.
7. Sasaki T, Suzaki T, Soyano T, Kojima M, Sakakibara H, et al. (2014) Shoot-derived cytokinins systemically regulate root nodulation. *Nat Commun*.19: 4983

**Citation:** Shams S, Naseem M (2017) Signal or To Sink: The Long-Sought-After Cytokinin Transporter. *Transcriptomics* 5: 141. doi:[10.4172/2329-8936.1000141](https://doi.org/10.4172/2329-8936.1000141)

### OMICS International: Open Access Publication Benefits & Features

**Unique features:**

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

**Special features:**

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>